

EXPRESS MAILING CERTIFICATE"EXPRESS MAIL" Mailing Label No.: EL084652896USDate of Deposit : January 31, 2001

I hereby certify that this paper or fee is being deposited
with the United States Postal Service "Express Mail Post Office to
Addressee" service under 37 CFR 1.10 on the date indicated above
and is addressed to the Assistant Commissioner for Patents, Box
Patent Application, Washington, D.C. 20231

Typed or printed name of person signing this certificate:

Signed: Pat Simpson
PAT SIMPSON

PATENT**Detection of Methylated CpG Rich Sequences Diagnostic for Malignant Cells**

This invention was conducted, at least in part, with government support under National
Institutes of Health Grants No: P30 CA16058 and CA80912 awarded by the National Cancer
Institute. The U.S. government has certain rights in the invention.

Background of the Invention

Diagnosis of cancer, classification of tumors, and cancer-patient prognosis all depend on
detection of properties inherent to cancer, or malignant cells, that are absent in normal,
nonmalignant cells. Since cancer is largely a genetic disease, resulting from and associated with
changes in the DNA of cells, one important method of diagnosis is through detection of related
changes within the DNA of cancer cells. Such changes can be of two types. The first type of
change is a genetic change that occurs when the sequence of nucleotide bases within the DNA is
changed. Base changes, deletions and insertions in the DNA are examples of such genetic
changes. The second type of change in the DNA is an epigenetic change. Epigenetic changes do
not result in nucleotide sequence changes, but rather, result in modification of nucleotide bases.
The most common type of epigenetic change is DNA methylation.

In mammalian cells, DNA methylation consists exclusively of addition of a methyl group
to the 5-carbon position of cytosine nucleotide bases. In the process, cytosine is changed to 5-
methylcytosine. Cellular enzymes carry out the methylation events. Only cytosines located 5' to

guanosines in CpG dinucleotides are methylated by the enzymes in mammalian cells. Such CpG dinucleotides are not distributed randomly throughout the genome. Instead, there are regions of mammalian genomes which contain many CpG dinucleotides, while other areas of the genome contain few CpG dinucleotides. Such CpG-rich areas of the genome are called "CpG islands."

5 Most often, CpG islands are located in the transcriptional promoter regions of genes.

Not all CpG islands are methylated. However, the methylation status of CpG islands (i.e., whether the CpG dinucleotides within a particular CpG island are methylated or not) is relatively constant in cells. Nevertheless, the pattern of CpG island methylation can change and, when it does, often a new, relatively stable methylation pattern is established. Such changes in
10 methylation of CpG islands can be either increases or decreases in methylation.

Methylation of CpG islands in the promoter region of a few specific genes has been observed in some types of human cancer. However, at present it is still uncertain whether the methylation status of multiple CpG islands in the genomic DNA of subjects suspected of having cancer can be used as a diagnostic tool for determining whether or not tissue obtained from such
15 subjects contain malignant cells.

Summary Of The Invention

The present invention relates to methods for identifying CpG islands which are diagnostic of one or more cancers in a subject. The method employs restriction landmark genomic scanning (RLGS) techniques and comprises separately digesting genomic DNA which has been obtained
20 from malignant cells derived from a particular tumor tissue and genomic DNA which has been obtained from control cells derived from healthy tissue with an infrequently cutting restriction enzyme that is not capable of cleaving methylated recognition sites to provide a first set of DNA restriction fragments from the tumor tissue, referred to hereinafter as "malignant cell restriction
25 fragments", and a first set of DNA restriction fragments from the healthy tissue, referred to hereinafter as "control cell restriction fragments"; attaching a detectable label to the ends of the malignant and control cell restriction fragments; digesting the labeled malignant and control cell restriction fragments with a second restriction enzyme; separating each set of restriction fragments on a gel; digesting the restriction fragments in each of the gels with a third more
30 frequently cutting restriction enzyme; electrophoresing each set of restriction fragments in a direction perpendicular to the first direction to provide a first pattern of detectable malignant cell

restriction fragments and a second pattern of detectable control cell restriction fragments; and comparing the second pattern to the first pattern to identify control cell restriction fragments, hereinafter referred to as "diagnostic fragments", which are absent, or exhibit an decreased intensity of label in the first pattern. Such fragments comprise CpG islands that are methylated in the malignant cells. Such patterns are useful for characterizing tissue which is suspected of containing malignant cells. Preferably, each of the diagnostic fragments is then isolated and sequenced, at least in part. In one preferred embodiment, the first restriction enzyme is NotI. In another preferred embodiment, the first restriction enzyme is AscI. Advantageously, the present method permits the detection of numerous methylation sites within the entire genome. In accordance with the present method, applicants have determined that particular CpG islands are preferentially methylated in DNA obtained from tumor tissues of subjects diagnosed as having breast cancer, glioma, acute myeloid leukemia, primitive neuroectodermal tumors of childhood, colon cancer, head and neck cancer, testicular cancer, and lung cancer.

The present invention also provides isolated polynucleotides, referred to hereinafter as "CpG diagnostic polynucleotides", and isolated oligonucleotides referred to hereinafter as "CpG diagnostic oligonucleotides", which are useful for characterizing tissue samples obtained from a subject suspected of having gliomas, acute myeloid leukemia, primitive neuroectodermal tumors of childhood, or cancer of the breast, colon, head and neck, testicle or lung. The CpG diagnostic polynucleotides and oligonucleotides both comprise a sequence which contains CpG islands that have been shown to be preferentially methylated in DNA that has been obtained from malignant cells of subjects diagnosed as having breast cancer, glioma, acute myeloid leukemia, primitive neuroectodermal tumor of childhood, colon cancer, head and neck cancer, testicular cancer or lung cancer. The CpG diagnostic polynucleotides are from 35 to 3000, preferably, 35 to 100 nucleotides in length, and comprise from 15 to 34, preferably 18 to 25 of the consecutive nucleotides contained with the sequences depicted in the accompanying DNA sequence listing, or sequences which are complementary thereto. The CpG diagnostic polynucleotides comprise two or, preferably, more CpG dinucleotides or dinucleotides which are complementary thereto. The CpG diagnostic oligonucleotides are from 15 to 34 nucleotides in length and comprise from 15 to 34 consecutive nucleotides contained within the sequences depicted in the sequence listing, or sequences which are complementary thereto. The CpG oligonucleotides comprises two or more CpG dinucleotides, or dinucleotides which are complementary thereto.

The present invention also relates to methods which employ the CpG diagnostic polynucleotides and oligonucleotides of the present invention to characterize tissue from patients suspected of having cancer. Such methods are based on the methylation status of CpG islands that have been shown to be preferentially methylated in DNA that has been obtained from tumor tissues of subjects diagnosed as having breast cancer, glioma, acute myeloid leukemia, primitive neuroectodermal tumor of childhood, colon cancer, head and neck cancer, testicular cancer and lung cancer. In one method, DNA which is isolated from suspected tumor tissue from a subject is digested into smaller fragments and reacted with a CpG diagnostic polynucleotide under stringent hybridization conditions. The reaction products are then assayed to determine the size or the sequence of the DNA fragment with which the CpG diagnostic polynucleotide has hybridized. The size or the sequence of the DNA fragment to which the CpG diagnostic polynucleotide has hybridized, hereinafter referred to as the "target DNA fragment", indicates whether the target DNA fragment comprises methylated or non-methylated CpG islands. The presence of methylated CpG islands in the target DNA fragment indicates that the DNA has been obtained from a tumor or neoplasm for which the diagnostic CpG polynucleotide serves as a diagnostic marker.

In another method the DNA from the suspected tumor tissue is treated with a chemical compound which converts non-methylated cytosines to a different nucleotide base. An example of such a compound is sodium bisulfite which converts non-methylated cytosines to uracil. The DNA is then reacted with at CpG diagnostic oligonucleotides under conditions which permit the CpG diagnostic oligonucleotide to hybridize with a complementary sequence in the DNA, referred to hereinafter as the "target sequence". The DNA is also reacted with a modified CpG diagnostic oligonucleotide. The modified CpG diagnostic oligonucleotide comprises a sequence that is complementary to a modified target sequence, i.e., a sequence in which the non-methylated cytosines in the target sequence are converted to a different nucleotide base, e.g. uracil, when treated with a chemical compound. The reaction products are then assayed to determine whether the DNA contains sequences which have hybridized with the CpG diagnostic oligonucleotide or with the modified CpG diagnostic oligonucleotide. Hybridization of the sample DNA with the CpG diagnostic oligonucleotide, as opposed to the modified CpG diagnostic oligonucleotide, indicates that the cytosines in the target sequence are methylated and that the DNA sample has been obtained from a tumor or neoplasm for which the CpG

oligonucleotide has been shown to serve as a diagnostic marker.

The present invention also relates to a method of identifying genes whose expression is increased or decreased in cancer cells.

5

Brief Description of the Figures

Fig. 1. Methylation detection in restriction landmark genomic scanning (RLGS) profiles. A, Diagram of the RLGS procedure showing the quantitative nature of methylation detection on NotI fragments displayed on RLGS profiles. Methylation detection in RLGS profiles depends on the methylation sensitivity of the endonuclease activity of NotI. Differences in digestion are assessed by radiolabelling the DNA at cleaved NotI sites. Following further endonuclease digestion, two-dimensional electrophoretic separation and autoradiography, the intensity of a DNA fragment on the resultant RLGS profile quantitatively reflects the copy number and methylation status of the NotI fragment. A priori, this allows NotI fragments containing single-copy CpG islands to be distinguished from the abundant NotI fragments present in repeat elements and rDNA sequences. B, A portion of an RLGS profile from normal peripheral blood lymphocyte DNA displaying nearly 2,000 single-copy NotI fragments and 15-20 high copy-number fragments. First-dimension separation of labeled NotI/EcoRV fragments extends from right to left horizontally. Following in-gel digestion with HinfI, the fragments are separated vertically downward into a polyacrylamide gel and autoradiographed. To allow uniform comparisons of RLGS profiles from different samples and different laboratories, each fragment is given a three-variable designation (Y coordinate, X coordinate, fragment number). The central region of the RLGS profile used for all comparisons described in this invention has 28 sections (1-5 vertically and B-G horizontally; the 4G and 5G sections were excluded due to high density and lower resolution of fragments). C, Enlarged view of profile section 2D, showing the numbers assigned to each NotI fragment. D, Analysis of the GC content and CpG ratio $\{(\text{number of CpGs})/(\text{number of guanines})(\text{number of cytosines})\}(\text{number of nucleotides analyzed})$ of 210 non-redundant NotI/EcoRV clones containing the NotI/HinfI fragments seen in B and in other portions of the RLGS profile. Of 210 clones, 184 clones were randomly chosen and 26 corresponded to fragments which were frequently lost from tumor profiles. CpG islands have a GC content of greater than 50% and a CpG value of 0.6 or greater, relative to bulk DNA (average CG content of 40% and CpG ratio of 0.2). Nucleotide sequences were determined with

greater than 99% accuracy overall. An average of 377nt/clone were analyzed (not indicative of actual CpG island size). The average NotI/EcoRV clone size was approximately 2 kb.

Fig. 2. Fragment loss from RLGS profiles is due to methylation. Top, portions of the RLGS profiles obtained from normal tissue and from two tumors having NotI fragments with either decreased intensity or no change in intensity. Bottom, Southern-blot analysis of EcoRV (NotI: -) and EcoRV/NotI (NotI: +) restriction digested DNAs from a larger number of samples, including the samples at top. In samples without methylation in the NotI site, the probe detects a smaller fragment on double digestion with NotI and EcoRV. The quantitation from multiple Southern blots using a phosphorimager allowed the determination of a lower limit of reliable detection in RLGS profiles of 30% decreased intensity of the diploid NotI/EcoRV fragments. Presence (+) or absence (-) of the corresponding NotI fragment is indicated. N, normal tissue DNA; T, tumor tissue DNA. A, CpG-island locus 3C1 methylation in low-grade gliomas. B, CpG island locus 2C40 methylation in leukemias. C, CpG-island locus 3E24 methylation in PNETs of childhood. *, EcoRV fragment of approximately 13 kb with homology to the probe. BLAST searches using the NotI-EcoRV clone sequence identified a homologous BAC clone sequence lacking an internal NotI site, which accounts for the 13-kb fragment on the Southern blot.

Fig. 3. Heterogeneity in CpG-island methylation across tumors. RLGS profiles were generated from 98 primary human tumors and compared with profiles of either matched normal DNA (58 of 98 cases) or to multiple profiles of tissue type-matched normal DNA from unrelated individuals. Loss or decreased intensity of single-copy fragments in the tumors, relative to several neighboring unaltered NotI fragments, were detected by visual inspection of overlaid autoradiographs and confirmed in many cases by independent profiles of the same DNA samples. For each tumor type, the dot plots display the total number of methylated CpG islands (of 1,184 CpG islands analyzed) observed in each tumor. Under the assumption that the tumors are drawn from a homogeneous distribution, with all tumors having the same frequency of methylation, the loss distributions should be approximately Poisson. The colored curve represents the expected distribution. BRE, breast tumors; CLN, colon tumors; GLI, gliomas; HN, head and neck tumors; LEU, acute myeloid leukemias; PNET, primitive neuroectodermal tumors of childhood; TST, testicular tumors.

Fig. 4. Subsets of CpG islands are preferentially methylated. For each tumor type, the histograms display the number of tumors in which the particular CpG islands were methylated. Most of the 1,184 CpG islands were not methylated in any of the tumors (histogram bar at 0 is not shown), but several CpG islands were methylated in multiple tumors. The black line shows the expected distribution under the null hypothesis that the CpG islands have equal frequencies of methylation. Most of the tumor types show significant preferential methylation.

Detailed Description Of The Invention

In one aspect, the present invention relates to methods for identifying clones within a DNA library that can be used for cancer diagnosis and tumor classification, based on the methylation status of CpG dinucleotides contained within or closely adjacent to the specific clones. Such method employs methylation-sensitive restriction endonucleases (MSREs) and restriction landmark genomic scanning (RLGS) gels to identify new, differentially-methylated CpG islands within malignant cells obtained from patients diagnosed as having cancer. In accordance with the present invention, Applicants have identified 93 clones which can be used to determine whether a tumor biopsy from a patient contains benign or malignant cells.

To carry out such method, tissue (referred to hereinafter as "tumor tissue") which contains a tumor or neoplasm is obtained from a patient known to have a cancer. In some cases, the tumor tissue is obtained from a particular type of solid tumor which has been surgically removed from the patient. In some cases, the tumor tissue is obtained from the hematopoietic system, such as for example, bone marrow or blood, of the patient. The tumor tissue will have been determined to be from either a benign or malignant tumor or neoplasm.

Separately, tissue (referred to hereinafter as "healthy tissue") which does not contain a tumor or neoplasm is obtained from a subject. The healthy tissue, may be obtained by surgically removing normal tissue from the patient or by surgically removing normal tissue from a healthy control subject who does not have cancer. The healthy tissue may also come from the hematopoietic system, such as for example, bone marrow or blood, of a healthy control subject. The healthy tissue will have been determined to be non-tumorigenic or non-neoplastic.

DNA is then isolated from both the tumor tissue and healthy tissue. If the tumor tissue is a solid tissue sample, such procedure may first comprise separating the individual cells contained

within the tissue from each other. For example, if the tissue samples were frozen after surgical removal from a patient, cells may be separated from one another by grinding the frozen tissue with a mortar and pestle. DNA is then isolated from the individual cells using procedures well known to those skilled in the art. Commonly, such DNA isolation procedures comprise lysis of the individual cells using detergents, for example. After cell lysis, proteins are commonly removed from the DNA using various proteases. The DNA is then commonly extracted with phenol, precipitated in alcohol and dissolved in an aqueous solution.

In the procedures which follow, the DNA obtained from the tumor tissue is treated separately from the DNA obtained from healthy tissue (i.e., the two DNAs are not mixed). The DNAs are separately analyzed using a method called restriction landmark genomic scanning (RLGS). The purpose is to analyze both DNAs separately. The two analyses are then compared in order to identify CpG islands that distinguish cancer cells from normal cells.

Both DNA samples are treated with restriction enzymes and the free ends that result from the restriction enzyme cleavage are labeled. However, since the isolated DNA is in linear pieces, there are free ends that exist before the DNA is cleaved with the restriction enzymes. To prevent these ends from being labeled, the ends, preferably, are blocked before restriction enzyme treatment. Such blocking can be done by addition of dideoxynucleotides and sulfur-substituted nucleotides to the free ends before treatment with restriction enzymes. Subsequently, when the DNA is cleaved by restriction enzymes and labeled, only the ends resulting from the restriction enzyme cleavage will be labeled.

After the reaction to block free ends, the DNA samples are cleaved with a first restriction enzyme that can be characterized as an infrequently cleaving, methylation-sensitive restriction enzyme. Examples of suitable first restriction enzymes are NotI, AscI, BssHII and EagI. As used herein the term "infrequently cleaving" refers to a restriction enzyme that is expected to cleave genomic DNA at intervals greater than 10 kilobases. For example, NotI is an infrequently cleaving restriction enzyme. NotI recognizes a nucleotide sequence of 8 base pairs (bp) in the genome (i.e., 5'GCGGCCGC3') and cleaves the DNA at this site. There are an estimated 4000-5000 of such NotI recognition sequences within the human genome. It is estimated that such recognition sequences are spaced at approximately 1 megabase (Mb) intervals within the genome. In contrast, a frequently cleaving restriction enzyme is expected to cleave the human genome at from 5-10 kb intervals. Such an enzyme will have approximately 100-times more

cleavage sites within the human genome than infrequently-cleaving enzymes. Such frequently cleaving enzymes usually recognize a nucleotide sequence of less than 8 bp in the genome and cleave the DNA at that site. However, not all restriction enzymes that have nucleotide recognition sequences of less than 8 bp are frequently cleaving enzymes. BssHII and EagI both have 6 bp recognition sequences but the recognition sequences for these two enzymes are spaced at intervals within the genome that are greater than 10 kb. "Methylation sensitive" as used herein refers to any enzyme that is unable to cleave DNA at its normal restriction site if one or more nucleotides within the recognition sequence is methylated. For example, the restriction enzyme NotI will cleave the 5'GCGGCCGC3' recognition sequence if the sequence does not contain a 5-methylcytosine. However, the NotI enzyme will not cleave this sequence if any of the cytosines have been methylated to become 5-methylcytosine.

Following digestion of the DNA with the first restriction enzyme, the ends of the DNA fragments are labeled. This can be done, for example, by attachment of nucleotides carrying a detectable label, such as a radiolabel, to the ends of the DNA sample. Typically, attachment is accomplished by filling in the recessed DNA ends left by cleavage with the first restriction enzyme such that the ends become blunt (i.e., non-recessed). Such end-filling reaction may employ deoxynucleoside triphosphates having a radiolabeled phosphate at the α phosphate position. Such labeled phosphate is preferably ^{32}P .

The labeled fragments from each sample are then cleaved with a second restriction enzyme. Such second restriction enzyme preferably cleaves human DNA at average intervals of between 5-10 kb. Such enzymes normally have a 6 bp recognition sequence. Preferably, the second restriction enzyme is not methylation sensitive. Examples of suitable second restriction enzymes are PstI, PvuI, EcoRV or BamHI. Cleavage of the DNA fragments with the second restriction enzyme provides a second set of fragments, labeled at the ends left by cleavage with the first enzyme. Many of such second fragments are smaller than the fragments resulting from cleavage with the first restriction enzyme.

The DNA fragments are then separated from one another. Preferably this separation is based on size. Preferably this separation is performed by first-dimension electrophoresis through an agarose tube-shaped gel of approximately 60 cm in length.

After electrophoresis through the tube-shaped gel, the DNA is digested within the gel with a third restriction enzyme. Such third restriction enzymes preferably have recognition

sequences of 4 or 6 bp. Such third restriction enzymes also have the property of being able to cleave DNA which is embedded within agarose. One such enzyme is *HinfI*.

After cleavage by the third restriction enzyme, the DNA is again separated based on size, preferably by electrophoresis through a polyacrylamide gel. Subsequently, the separated DNA fragments are detected based on the labeled ends of the DNA fragments. In those cases where the fragments are radiolabeled, detection is by autoradiography of the two-dimensional gel. Such autoradiography provides a pattern of DNA fragments or "spots." Such pattern is called an RLGS profile.

Each fragment on the RLGS profile obtained from using the DNA from healthy tissues is uniquely identified by its location on the autoradiograph (Y coordinate, X coordinate, fragment number). For each fragment location on the RLGS profile obtained from healthy tissue DNA, the identical location is observed on the RLGS profile obtained from tumor tissue DNA.

In a fragment by fragment comparison of RLGS profiles obtained from tumor tissue DNA with healthy tissue DNA, three different patterns are possible. First, for a given fragment on the healthy tissue RLGS profile, there may be a corresponding fragment at the same location, and of the same intensity, on the tumor tissue RLGS profile. This indicates that the first restriction enzyme cleaved both DNAs at the same sequences (Fig. 1A). This indicates that there were no differences in methylation of the *NotI* nucleotide recognition sequence of that fragment between the tumor tissue DNA and the healthy tissue DNA.

Second, for a given fragment on the healthy tissue RLGS profile, there may be no fragment at the same location on the tumor tissue RLGS profile. Such a pattern indicates that the first restriction enzyme did not cleave the tumor tissue DNA at the recognition sequence required to produce that specific fragment, but did cleave at such sequence within the healthy tissue DNA (Fig. 1A). This indicates that there was methylation within the *NotI* recognition sequence in the tumor tissue DNA but not in the healthy tissue DNA.

Third, for a given fragment on the healthy tissue RLGS profile, there may be a corresponding fragment at the same location on the tumor tissue RLGS profile, but the intensity of the fragment may be of decreased intensity. Such a pattern indicates that the first restriction enzyme cleaved one of two copies (i.e., the genome is diploid) of the tumor tissue DNA at the recognition sequence required to produce that specific fragment (Fig. 1A). In healthy tissue DNA, the first restriction enzyme cleaved both copies of the recognition sequence. This

indicates that there was methylation within one of two NotI recognition sequences in the tumor tissue DNA.

Through comparisons of RLGS profiles obtained from healthy tissue DNA with profiles obtained from a large number of different tumor tissue DNAs, loss of specific fragments in multiple tumors can be associated with a specific type of cancer. Loss of such fragments from RLGS profiles, therefore, can be diagnostic for cancer in a subject. For example, loss of a specific fragment (i.e., methylation of the first restriction enzyme site at the end of said fragment) in a high percentage of tumor tissue DNAs from women known to have breast cancer can be diagnostic for breast cancer in subjects suspected of having the disease. To perform such a diagnostic analysis, DNA isolated from a patient suspected of having breast cancer would be analyzed by RLGS, as described above, to determine whether there was loss of one or more fragments in RLGS profiles that are known to be lost at high frequency in women known to have breast cancer. Similarly, loss of other specific fragments can be diagnostic for other cancers, such as for example, colon cancer, head and neck cancer, lung cancer, testicular cancer, neuroectodermal cancer, gliomas, acute myeloid leukemias, and others.

Loss of a specific fragment in RLGS profiles from multiple tumors can also be diagnostic of several types of cancer, rather than a single type of cancer. For example, loss of a specific fragment can occur in a high percentage of tumor tissue DNAs obtained from individuals with either breast, colon or lung cancer. Loss of such a spot from RLGS profiles using DNA obtained from a patient suspected of having cancer would be diagnostic for either breast, colon or lung cancer in that patient.

Isolated Polynucleotides and Oligonucleotides Diagnostic for Cancer

Individual DNA clones that contain the DNA present in each spot or fragment that makes up an RLGS profile can be obtained. This is done by constructing a DNA library of healthy tissue DNA that has been cleaved with the same first and second enzymes used to perform the RLGS gel analysis. Such DNA library will contain individual clones, each clone comprising DNA that is present in a single spot of the RLGS profile. The totality of clones within the library is representative of the combined DNA spots in the RLGS profile.

Individual clones within the library can be identified that contain the DNA of each spot on the RLGS profile. This can be done by taking DNA from one or a few individual clones of the DNA library and mixing it with healthy tissue DNA, before RLGS analysis is begun. When

this mixture of DNAs is used to produce an RLGS profile, the intensity of the spots that contain the same DNA as the individual clones added to the mixture will be increased. By performing multiple analyses of this type, each spot on an RLGS profile can be matched up with a DNA clone within the library. The result of such an analysis is an ordered human genomic library of restriction fragments containing the same subset of genomic fragments as those displayed on RLGS profiles. In such ordered genomic libraries, an individual library clone corresponding to any spot or fragment in an RLGS profile can be rapidly located.

To design diagnostic CpG polynucleotides and oligonucleotides, the sequence of the DNA within each clone (referred to hereinafter as a “diagnostic clone”) that corresponds to a spot that is absent or exhibits decreased intensity on the RLGS profile of the DNA from malignant tumor tissue is sequenced using standard techniques. Once sequence information is obtained, regions comprising multiple CpG dinucleotides are located. Such regions serve as the target sequence for the CpG polynucleotides and oligonucleotides.

The CpG polynucleotides are from 35 to 3000, preferably from 35 to 1500 nucleotides in length and comprise two or, preferably, more CpG dinucleotides or dinucleotides which are complementary thereto. The CpG diagnostic oligonucleotides are from 15 to 34 nucleotides, preferably from 18 to 25 nucleotides, in length and comprise at least two CpG dinucleotides or dinucleotides which are complementary thereto. The CpG diagnostic polynucleotides and oligonucleotides each comprise a sequence which is substantially complementary to target sequences containing CpG islands that are known to be preferentially methylated in the DNA from one or more types of cancer cells. “Substantially complementary” means that there is enough complementarity between the CpG diagnostic polynucleotides or oligonucleotides and the target sequence so that hybridization occurs between the CpG diagnostic polynucleotides and oligonucleotides under stringent conditions, preferably under highly stringent conditions. Such assays include hybridization assays, such as for example Southern analysis, where the sample DNA is reacted with the CpG diagnostic polynucleotide under stringent hybridization conditions.

The term “stringent conditions, as used herein, is the “stringency” which occurs within a range from about $T_m - 5$ (5 below the melting temperature of the probe) to about 20 C below T_m . “Highly Stringent hybridization conditions” refers to an overnight incubation at 42 degree C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt’s solution, 10% dextran sulfate, and 20 g/ml denatured,

sheared salmon sperm DNA, followed by washing the filters in 0.2x SSC at about 65 degree C. As recognized in the art, stringency conditions can be attained by varying a number of factors such as the length and nature, i.e., DNA or RNA, of the probe; the length and nature of the target sequence, the concentration of the salts and other components, such as formamide, dextran sulfate, and polyethylene glycol, of the hybridization solution. All of these factors may be varied to generate conditions of stringency which are equivalent to the conditions listed above.

Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lower stringency); salt conditions, or temperature. For example, moderately high stringency conditions include an overnight incubation at 37 degree C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2 M NaH₂PO₄; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 ug/ml salmon sperm blocking DNA; followed by washes at 50 degree C with 1XSSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC).

Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

Such assays also include polymerase chain reactions (PCR) where the sample DNA and the diagnostic CpG oligonucleotides are reacted, preferably under conditions which result in the synthesis of a single PCR product. Computer programs, such as for example, the "Primer3" program that can be accessed at "http://genome.wi.mit.edu/cgi-bin/primer/primer_3www.cgi" can be used to determine the size and sequence of the CpG diagnostic oligonucleotides. Optimum conditions are determined empirically.

The CpG diagnostic polynucleotides and oligonucleotides are made using standard techniques. For example, these polynucleotides and oligonucleotides may be made using commercially available synthesizers.

Diagnostic Methods

In another aspect, the present invention relates to methods which use the CpG diagnostic

polynucleotides and oligonucleotides to characterize tissue samples from a subject suspected of having cancer, referred to hereinafter as test sample DNA. To do this, DNA is isolated from the cells of the tissue sample of the patient. Preferably, DNA that serves as a control is also obtained from healthy tissue of the test subject or a control subject as described previously. The

5 diagnostic methods comprise reacting the test sample DNA with the diagnostic CpG polynucleotide or oligonucleotide and assaying the products that are formed as the result of the reaction. In some cases, the sample DNA is digested into smaller fragments prior to reaction with the CpG diagnostic polynucleotides or oligonucleotides. In some cases, a portion of the test sample DNA is first reacted with a chemical compound, such as for example sodium

10 bisulfite, which converts methylated cytosines to a different nucleotide base.

Southern Blot Analysis

One such method for diagnosing cancer in a patient involves cleavage of the test sample DNA with a methylation sensitive enzyme, then Southern blot analysis of said cleaved DNA using a CpG diagnostic polynucleotide or oligonucleotide as a probe. For example, the DNA

15 from the patient and the control, healthy tissue DNA are separately cleaved with a methylation-sensitive restriction endonuclease, such nuclease being the same first restriction enzyme used to identify the diagnostic spot in the RLGS profile that corresponds to the CpG diagnostic polynucleotide or oligonucleotide. After cleavage, the test sample and control DNAs are electrophoretically separated by size in different lanes of the same agarose gel and blotted to a

20 membrane that can be used in hybridization, such as for example, nitrocellulose or nylon. The membrane is then used in a hybridization reaction with a labeled CpG diagnostic polynucleotide or oligonucleotide. The labeled CpG diagnostic polynucleotide or oligonucleotide will hybridize to complementary DNA sequences on the membrane. After hybridization, the location on the membrane where the probe hybridized to the control and patient DNAs is visualized.

25 Such locations will identify DNA fragments or bands within the control and patient DNAs containing the same sequence as the CpG diagnostic polynucleotide or oligonucleotide. Hybridization of the probe to a fragment within the patient DNA that is of higher molecular weight than that of the fragment within the control DNA to which the probe hybridized, indicates that a restriction endonuclease cleavage site flanking the target sequence of the CpG diagnostic

30 polynucleotide or oligonucleotide was not cleaved due to methylation. Such result indicates that the tissue is from a cancer for which the CpG diagnostic polynucleotide or oligonucleotide serves

as a diagnostic tool.

A second method for diagnosing cancer in a patient involves cleavage of patient DNA with a methylation-sensitive restriction endonuclease, such nuclease being the same first restriction enzyme used to identify the diagnostic spot in the RLGS profile that corresponds to the fragment. Such nuclease will cleave the patient DNA at the diagnostic recognition sequence only if the DNA is unmethylated. Using nucleotide information derived from sequencing of the library clone corresponding to the diagnostic spot on the RLGS gel, primers for PCR are selected that span the diagnostic recognition sequence. Using the primers, PCR is performed on the DNA. PCR amplification of the sequences will be successful only if the diagnostic nucleotide sequence in the patient DNA had been methylated and was not cleaved by the enzyme. Successful PCR amplification, therefore, is indicative of cancer in the patient.

Methods Employing a Chemically-Modified DNA Test Sample

Another group of methods for diagnosing cancer in a patient using CpG diagnostic polynucleotides and oligonucleotides are based on treatment of patient DNA with sodium bisulfite which converts all cytosines, but not methylated cytosines, to uracil. The bisulfite converted patient DNA can then be analyzed in a number of different ways. One method of analysis is direct sequencing of the DNA to determine whether the sequence contains cytosine or uracil. Such DNA sequencing requires primers adjacent to the sequenced region to be made. Such primers would be based on DNA sequence information obtained from the diagnostic RLGS spots.

Another method of analyzing bisulfite converted patient DNA is a method called "methylation sensitive PCR" (MSR). In MSR, primers are designed to comprise a sequence which is substantially complementary to the the CpG islands which are known to be preferentially methylated in DNA of cells found in one or more type of tumor tissues. Two sets of PCR primers are made to ^{encompass} ~~encompass~~ this region. One set of primers is designed to be complementary to the sequence that was changed by bisulfite (i.e., cytosines that were originally unmethylated and changed to uracil). As discussed above, these are the modified CpG diagnostic oligonucleotides. A second set of primers is designed to be complementary to the same sequence that was not changed by bisulfite (i.e., cytosines that were methylated and not changed to uracil). As discussed above these are the unmodified CpG diagnostic oligonucleotides, i.e the oligonucleotides which containe at least two CpG dinucleotides or

dinucleotides which are complementary thereto. Two sets of PCR reactions are then run, one reaction with each set of primers, using DNA from the subject as the template. In the case where cytosines within the target sequence of the subject DNA are not methylated, the target sequence will be modified by the chemical reaction and the primers complementary to the modified sequence, i.e., the modified CpG diagnostic oligonucleotides, will produce a PCR reaction product while the primers complementary to the methylated sequence, i.e., the unmodified CpG diagnostic oligonucleotides, will not produce a PCR product. In the case where cytosines within the target sequence of the subject DNA are methylated, the target sequence will not be altered by the reaction with the sodium bisulfite, and the primers complementary to the unaltered sequence, i.e., the unmodified CpG diagnostic oligonucleotides, will produce a PCR reaction product while the modified CpG diagnostic oligonucleotides, which are complementary to the modified target sequence (i.e., unmethylated sequence) will not produce a PCR product

A modification of MSR is bisulfite treatment of patient DNA and PCR amplification of said DNA using primers designed to amplify either methylated or unmethylated sequences. The PCR product is then digested with a restriction enzyme that will cleave or not depending on whether said product contains uracil (rather, thymidine, the complement of uracil; found in PCR product if original patient DNA contained unmethylated cytosine) or cytosine (found in PCR product if original patient DNA contained methylated cytosine).

Another technique referred to as MS-SnuPE, uses bisulfite/PCR followed by primer extension, where incorporation of C (vs. T) denotes methylation.

Methods of Identifying Genes

In another aspect of the invention, the CpG diagnostic polynucleotides and oligonucleotides can be used as probes to identify genes whose expression is increased or decreased in cancerous tissues. To do this, CpG diagnostic polynucleotides are reacted with individual clones of the DNA library. The clones which hybridize with the CpG diagnostic polynucleotide can then be analyzed to determine if they contain an open reading frames that could encode proteins. To determine if the CpG diagnostic polynucleotide hybridizes with the promoter region of a known gene, the open reading frame sequence is analyzed by searching existing DNA databases. For example, GenBank databases can be searched using the BLAST algorithm. If no known genes that correspond to a library clone is found, the sequence can be

searched for open reading frames that could encode a protein. Such searching can be performed using commercially available sequence analysis programs commonly known to those skilled in the art. GCG is an example of one such program.

Sequences from clones of the DNA library that contain either known genes or open reading frames can be used as probes to determine whether genes encoded by the sequences are expressed in tumor tissues as compared to control, healthy tissues. To do this, RNA, preferably messenger RNA (mRNA) is isolated from healthy tissue and from tumor tissue from which it is desired to test expression. Such RNA is examined for the presence of expressed transcripts encoded by the sequences obtained from the library. Examination for the presence of expressed transcripts can be performed using a number of methods. One method is Northern blotting where the isolated RNA is separated by size using gel electrophoresis and then blotted to a hybridization membrane. A fragment, polynucleotide or oligonucleotide from the sequence obtained from a library clone is labeled and then used to probe the hybridization membrane containing the size-separated RNA. Detection of hybridization of the probe to the membrane indicates presence of a transcript encoded by the sequence and indicates expression of the gene encoded by that sequence.

Another method to examine isolated RNA for the presence of expressed transcripts is to use RT-PCR analysis. In such analysis, primers are designed and made that span a region of the gene whose expression is to be tested. The isolated RNA is reverse transcribed into DNA using reverse transcriptase. Such DNA is then amplified with the designed primers using PCR. PCR products are visualized after electrophoresis. The presence of PCR products on the gel indicates that the gene encompassed by the designed primers was expressing RNA transcripts. Such analysis can identify and determine genes whose expression is changed in cancer cells as compared to normal, non-cancerous cells.

The following examples are for purposes of illustration only and are not intended to limit the scope of the invention as defined in the claims which are appended hereto.

Examples

Example 1. Identification of diagnostic markers using NotI and RLGS

A. Isolation and enzymatic processing of genomic DNA

5 Tissue from solid tumors was obtained as surgical tissue samples. Where possible, surrounding non-tumor tissue was taken and used as a control. Where it was not possible to obtain patient-matched normal tissue, normal tissue from multiple patients was used. Tissue samples from patients with acute myelogenous leukemias (AML) consisted of either bone marrow aspirates or peripheral blood. Normal samples were obtained from the same patients who were
10 in remission after chemotherapy.

The surgically removed tissues were quickly frozen in liquid nitrogen and stored at -80°C prior to isolation of DNA. When DNA was ready to be isolated, 2 ml of lysis buffer (10 mM Tris, pH 8.0; 150 mM EDTA, 1% sarkosyl) was added to 100-300 mg of tissue in a 50 ml Falcon tube and frozen in liquid nitrogen. The frozen mixture was then removed from the tube, wrapped
15 in aluminum foil, and quickly broken into pieces with a hammer. The broken pieces of cells were transferred to a chilled mortar and ground to a powder with a chilled pestle. For peripheral blood samples, cells were separated on a sterile Histopaque-1077 (SIGMA) gradient and stored at -80°C before DNA isolation.

Cells were transferred to a 50 ml tube and 15-25 ml of lysis buffer containing 0.1 mg proteinase K per ml of lysis buffer was added and mixed using a glass rod. The mixture was incubated at 55°C for 20 min with gentle mixing every 5 min. The mixture was then placed on ice for 10 min. Subsequently, an equal volume of PCI (phenol:chloroform:isoamylalcohol in a ratio of 50:49:1) was added and the tubes containing the mixture were gently rotated for 30-60 min. The tubes were then centrifuged for 30 min at 2500 rpm and the separated, aqueous phase
25 was transferred to a new 50 ml tube using a wide-bore pipette. The PCI extraction was repeated one time. The collected aqueous phase containing the DNA was transferred to dialysis tubing and dialyzed against 4 L of 10 mM Tris, pH 8 for 2 hr. The dialysis tubing was then transferred into fresh 10 mM Tris and dialyzed overnight at room temperature. One additional dialysis was performed in fresh 10 mM Tris for an additional 2 hr. The DNA was then transferred from the
30 dialysis tubing to 50 ml tubes and RNase A was added to a final concentration of 1 µg/ml. The mixture was incubated at 37°C for 2 hr. Subsequently, 2.5 volumes of 100% ethanol were added

to the DNA and the mixture was gently rotated. The insoluble DNA was transferred to a microfuge tube, centrifuged briefly, and the remaining alcohol removed. The pellet was briefly dried in air. The DNA in the pellet was resuspended to a final concentration of 1 µg/µl. Such isolated DNA had an average size of 200-300 kb.

5 The isolated genomic DNA was blocked at ends where the DNA had been sheared. Blocking was done by addition of dideoxynucleotides and sulfur-substituted nucleotides. In a 1.5 ml tube, 7 µl of genomic DNA solution was added along with 2.5 µl of blocking buffer (1µl 10X buffer 1, 0.1µl 1 M DTT, 0.4 µl each of 10 µM dGTPαS, 10 µM ddATP, 10 µM ddTTP, and 0.2 µl 10 µM dCTPαS; buffer 1 consists of 500 mM Tris, pH 7.4, 100 mM MgCl₂, 1 M NaCl, 10mM DTT) and 0.5 µl DNA polymerase I. The mixture was mixed thoroughly and incubated at 37°C for 20 min. The mixture was then incubated at 65°C for 30 min to inactivate the polymerase. The reaction was then cooled on ice for 2 min. The DNA was digested with NotI by adding to the sample, 8µl of 2.5X buffer 2 (20X buffer 2 is 3 M NaCl, 0.2% Triton X-100, 0.2%BSA) and 2µl (10 U/µl) of NotI. The sample was incubated at 37°C for 2 hr. The DNA was then radioactively labeled. This was done by adding to the sample 0.3 µl 1 M DTT, 1 µl [α -³²P]-GTP, 1 µl [α -³²P]-dCTP and 0.1 1 µl[α -³²P]-GTP Sequenase ver 2.0 (13 U/µl). The mixture was incubated at 37°C for 30 min. The DNA was then digested with EcoRV by adding to the sample 7.6 µl second enzyme digestion buffer (1 µl 1 mM ddGTP, 1 ul 1 mM ddCTP, 4.4 µl ddH₂O, 1.2 µl 100 mM MgCl₂) and 2 µl EcoRV (10 U/µl). The mixture was incubated at 37°C for 1 hr. Then, 7 µl of 6X first-dimension loading dye (0.25% Bromophenol Blue, 0.25% Xylene Cyanol, 15% Ficoll type 400) was added.

B. First dimension gel set-up and electrophoresis

To make the 60 cm long agarose tube-shaped gel, a gel holder was made. To do this, a sharp razor was used to cut one end of PFA-grade teflon tubing (PFA 11 thin wall, natural; American Plastic, Columbus, Ohio) at an angle to make a bevel. The beveled end of the tubing was fed into glass tubes (4 mm inner diameter, 5 mm outer diameter, 60 cm long). Using a hemostat, the beveled end was pulled up through the tapered end of the glass rod until it protruded 2 to 4 cm. The tubing was cut horizontally at the same end, leaving a 2 mm protrusion (this is the top of the gel holder). The opposite end was cut horizontally, leaving a 5 to 6 cm protrusion from the glass tube. The gel holder was inverted and the top protruding end was

pressed firmly against a hot metal surface (metal spatula heated by a Bunsen burner) to fold the edges of the teflon outward onto the rim of the glass support. A rubber stopper with cored center was pulled over the top end of the gel holder until it was just past the taper of the glass rod. A two-way stopcock was attached to a 10 ml syringe and then to the gel holder via 2 to 3 cm of flexible tubing. The stopcock valve was adjusted to the open position.

Then, to a clean 200 ml glass bottle was added, 60 ml 2X Boyer's buffer (20X is 1 M Tris, 360 mM NaCl, 400 mM sodium acetate, 40 mM EDTA) and 0.48 g Seakem GTG agarose (0.8%). The mixture was heated in a microwave oven until the agarose was dissolved. The mixture was then equilibrated to 55°C in a water bath. With the stopcock valve in the open position, the protruding teflon tube was lowered into the molten agarose solution. The gel solution was suctioned into the gel holder until the gel solution reached 1-2 cm from the top of the gel holder. The stopcock valve was then closed. Keeping the gel upright, the gel was suspended from a ring stand. The gel was allowed to solidify for 20 min.

The stopcock valve was then opened and the syringe and connecting tubes were removed from each gel. After adding 2X Boyer's buffer to the bottom of the first dimension gel apparatus (C.B.S. Scientific), the gels were lowered into the first dimension gel apparatus, seating the rubber stopper firmly into the appropriate holes in the top portion of the apparatus. The top chamber was filled with 2X Boyer's buffer.

Between 1.0-1.5 µg of DNA was loaded onto each gel. The sample was electrophoresed at 110 V for 2 hr, and then 230 V for 24 hr.

C. In-gel digest

After the DNA was electrophoresed in the first dimension in the agarose tube gel, the DNA was further digested with an additional restriction endonuclease so it could be electrophoresed in the second dimension. In order to perform this additional endonuclease digestion, the buffer and gel holders were removed from the first dimension apparatus. The gel was extruded into a pan containing 1X buffer K (10X buffer K is 200 mM Tris, pH 7.4, 100 mM MgCl₂, 1 M NaCl) by forcing the gel out through the bottom of the gel holder. This was accomplished using a 1 ml syringe fitted with a pipet tip and filled with buffer K. The tip was firmly inserted into the top of the gel holder and the plunger depressed until the gel began to come out through the bottom of the gel holder. The 1 ml syringe was replaced with a 5 ml

syringe, and the plunger was depressed until the entire gel was expelled. With a razor, a bevel was cut in the low molecular weight end of the gel and a horizontal cut was made at the high molecular weight end so that the gel was approximately 43 cm in length. The gel length was now the same as the width of the second dimension gel.

5 The gel was placed into a separate 50 ml tube containing 40 ml of 1X buffer K. The tube was incubated for 10 min at room temperature. The buffer was poured off and the gel incubated in 1X buffer K for an additional 10 min. The buffer K and gel was poured into a pan containing fresh buffer K. Using a 10 ml syringe attached to restriction digest tubing (PFA grade teflon, 9, thin wall, natural; 2.7 mm inner diameter and approximately 3.3 mm outer diameter; American
10 Plastic, Columbus, Ohio), via a 1 to 2 cm segment of flexible tubing, the gel was suctioned into the digest tubing, low molecular weight (beveled) end first. The gel was suctioned into the digest tubing by placing the end of the tubing in line with the beveled end of the gel and pulling the syringe plunger. The tubing was positioned vertically, with the syringe at the bottom and remaining buffer from the tubing was suctioned into the syringe.

15 In a clean tube, a 1.6 ml mix of 1X *Hinf*I restriction enzyme buffer (50 mM NaCl, 10 mM Tris pH 7.9, 1 mM DTT), 0.1 % BSA, and 750 U of *Hinf*I restriction enzyme was made. The open end of the digest tubing was placed into the tube containing restriction digestion solution. Holding the syringe end up, suction was applied until a small amount of digestion solution appeared in the syringe. The digest tubing was removed and both ends were oriented
20 upward in a U-shape. The syringe was removed and the two ends of the tubing were attached to form a closed circle. This was placed in a moist chamber and incubated at 37°C for 2 hr.

D. Second dimension electrophoresis

25 The digested DNA was now run in the second dimension using a 5% non-denaturing acrylamide gel with a 0.8% agarose spacer. To do this, the second dimension gel apparatus (C.B.S. Scientific) was first assembled. All glass plates were cleaned thoroughly and the non-beveled face of each plate was coated with Gelslick or Sigmacote (only once every 10 uses). The back half of the apparatus was laid horizontally on a table top with the upper buffer chamber hanging over the table edge. The two small clear plastic blocks were inserted at the bottom
30 corners of each apparatus. A glass plate was placed in the apparatus, beveled edge facing upward and near the upper buffer chamber, followed by two spacers, one along each side. Glass

plates and spacers were added in this manner until the fifth plate had been added. After the third plate, flexible Tygon tubing was slid down the side channel of the apparatus, with a bevel cut in the leading end of the tubing. The other end was cut, leaving approximately 10 cm protruding from the apparatus. The Plexiglas "filler" sheet was placed over the fifth glass plate. The front half of the apparatus was positioned by aligning the screw holes of the front and back half. These were secured with the teflon screws. The oblong oval "windows" at the lower, front face were sealed with Plastic tape (Scotch brand). The apparatus was stood upright in the lower buffer chamber.

Using a three-way stopcock, the gel apparatus tubing was attached in series with a 2 L reservoir and a 60 ml syringe was attached to the remaining stopcock outlet. The tubing was attached to the 2 L reservoir through a bottom drain (a 2 L graduated cylinder was used). The reservoir was secured above the gel apparatus to allow for gravity flow. The stopcock valve was adjusted to allow liquid to flow between the 2 L reservoir and the 60 ml syringe. Once the TEMED was added, the acrylamide solution (1X TBE, pH 8.3, 96.9 g acrylamide, 3.3 g bis-acrylamide, 1.3 g ammonium persulfate and 700 μ l TEMED in a total volume of 2 L) was poured into the 2 L reservoir. The syringe plunger was pulled down to the 50 ml mark. The plunger was depressed to push the air out of the upper tubing. Once all air was removed, the valve was adjusted so that all three ports were open. Acrylamide flowed into the apparatus, filling all four gels simultaneously from the bottom upward. The flow was stopped when the level reached 3 mm from the top edge of the glass plates. The solution was allowed to settle for 2 to 3 minutes. After the valve leading to the gel apparatus had been closed, the syringe and reservoir were detached.

The ends of the in-gel digest tubing were separated and the first dimension gel was extruded into a pan containing 1X TBE, pH 8.3. The gel was transferred to a 50 ml tube containing 40 ml 1X TBE, pH 8.3. This was incubated for 10 min at room temperature, replaced with fresh TBE, and incubated for an additional 10 min. The first dimension gel was placed in a horizontal position across the beveled edge of each glass plate. Once all gels were in place, the space between the agarose gel and the top of each polyacrylamide gel was filled with molten 0.8% agarose (equilibrated to 55°C). This connecting agarose was allowed to solidify for 10 to 15 min and then 250 μ l second dimension loading dye (1X TE, pH 8.3, 0.25% Bromophenol Blue, 0.25% Xylene Cyanol) was added along the length of each gel. Then 1X TBE, pH 8.3 was

added to the upper and lower buffer chambers and electrophoresis was carried out at 100 V for 2 hr and then at 150 V for approximately 24 hr.

Buffers were then removed and the apparatus was disassembled. Each gel was lifted from the plates by overlaying with Whatmann paper cut to size for autoradiographic or phosphorimager cassettes. The perimeter of the paper was traced with the edge of a plastic ruler, removing any excess gel. The Whatmann paper and gel were lifted and placed, gel side up, on a second piece of Whatmann paper. This was overlaid with saran wrap and a third piece of Whatmann paper was added to the top and saran wrap was folded over the top of the Whatmann paper. This was placed in a gel drier, in the same orientation, for 1 hr at 80°C while applying a vacuum. The lower and upper Whatman paper was then removed, saran wrap folded under the remaining paper and exposed to X-ray film (BioMax MS).

E. RLGS spots resulting from methylation-sensitive restriction enzymes identify CpG islands

Using this methodology, an RLGS profile of DNA from human cells produces a pattern displaying approximately 2,000 spots. Fig. 1B, for example, shows such an RLGS profile from normal peripheral blood lymphocyte DNA. First-dimension separation of labeled NotI/EcoRV fragments extends from right to left horizontally. Following in-gel digestion with HinfI, the fragments were separated vertically downward into a polyacrylamide gel and autoradiographed. To allow uniform comparisons of RLGS profiles, spots were defined based on their location in the gel by assigning each spot a three-variable designation (Y coordinate, X coordinate, fragment number). This can be more easily seen in the enlarged portion of section 2D of the RLGS profile (Fig. 1C) showing the numbers assigned to each spot.

From a set of 1,567 NotI spots comprising the central portion of the RLGS profile of normal DNA, 392 spots were eliminated from all analyses on the basis of having more than diploid intensity, less than diploid intensity, or a degree of positional overlap with neighboring fragments. In addition, a small fraction of loci in individual tumor profiles was not able to be analyzed due to poor local gel quality. In normal DNA profiles, the less-than-diploid copy-number intensities can result from polymorphism, partial methylation or spots derived from sex chromosomes. Thus, the analyzed spots were of diploid copy number in most samples. Tumor tissue and healthy tissue DNA profiles were compared by visual inspection of overlaid autoradiographs. In those cases in which matched normal tissue was not available, tumor

profiles were compared with profiles matched for tissue type of four to five unrelated individuals. Each CpG island was defined as unmethylated or methylated (a visually apparent decrease in intensity on the RLGS profile, which, through corroboration with Southern-blot data for 26 CpG island loci and more than 100 loss events, corresponded to a 30% or greater level of methylation).

To determine if the NotI restriction sites which produced the RLGS spots, had characteristics of authentic CpG islands, DNA from 210 of the NotI/EcoRV RLGS spots was partially sequenced. This was possible because each spot on the human NotI/EcoRV RLGS profile had previously been assigned to a clone from a NotI/EcoRV genomic plasmid library (see description earlier in the specification). Of the 210 spots, 184 were randomly chosen. Another 26 spots were chosen because they were frequently lost from RLGS profiles from human tumors, suggesting that cytosine nucleotides within the NotI sequence of that spot were methylated in the tumor. From the sequences derived from these clones, the GC content (%GC) was plotted against the CpG ratio for each clone (Fig. 1D; $\text{CpG ratio} = \frac{(\text{number of CpGs})}{(\text{number of guanines})(\text{number of cytosines})(\text{number of nucleotides analyzed})}$). CpG islands have a GC content of greater than 50% and a CpG value of at least 0.6. Fig. 1D shows that, of 210 clones sequenced, 197 (94%) had sequence characteristics consistent with CpG-island DNA.

F. Tumor tissue samples analyzed

DNA used to perform the RLGS analyses was obtained from 98 primary human tumors and, where possible, matched normal samples. These samples were from 8 broad tumor types; breast, colon, gliomas, head and neck, acute myeloid leukemias, primitive neuroectodermal tumors (PNETs) and testicular.

Fourteen breast cancers included 2 adenocarcinomas, 2 lobular carcinomas and 10 ductal carcinomas. The samples were from obtained the Cooperative Human Tissue Network (CHTN). All tumors were from females, 38-89 years of age (average of 54 years). Breast tissue adjacent to the tumor was available for 6 of 14 cases, and 8 tumor profiles were compared with 4 breast samples from the matched sets.

Colon tumors were obtained from Roswell Park Cancer Institute and classified according to the American Joint Committee on Cancer staging manual. The 8 primary tumors included 1 stage I tumor, 2 stage II tumors, 2 stage III tumors and 3 stage IV tumors. Patient ages ranged

from 49 to 77 years (average of 63 years). Normal adjacent colon mucosa samples were obtained for all tumors.

Fourteen gliomas, including 12 World Health Organization (WHO) grade II astrocytomas and 2 WHO grade III anaplastic astrocytomas, from Saitama Medical School, the University of Tokyo, Teikyo University School of Medicine, Komagome Metropolitan Hospital and the University of Washington, Seattle. Patients included 10 females and 4 males with an age range of 7-57 years (average of 34 years). Brain tissue adjacent to the tumor was also obtained for 1 WHO grade II and 1 WHO grade III tumor. Twelve cases were compared with 3 unmatched normal brain samples and with the 2 brain samples from the matched sets.

Fourteen head and neck squamous cell carcinomas were obtained through the CHTN. Tumors were from 11 males and 3 females. Patients were 42-77 years of age (average of 57 years). Tissue adjacent to the tumor was available for 12 of 14 cases, and 2 tumors were compared with 4 samples from the matched sets.

Nineteen acute myelogenous leukemia samples (3 bone marrow aspirates and 14 peripheral blood) from the Cancer and Leukemia Group B Tissue Bank. Samples were classified according to the French-American-British system. Samples were obtained from patients at the time of initial diagnosis with AML and again at complete remission (24-154 days, average 45 days) after induction chemotherapy. Samples were from 14 males and 3 females. Patients were 22-61 years of age (average 40 years). All cases were compared with matched samples (either peripheral blood lymphocytes or bone marrow, but always matched with the origin of the cancer sample) obtained at remission.

Twenty-two PNETs, including 17 medulloblastomas and 5 supratentorial PNETs, through the CHTN, Pediatric Division. Tumors were from 15 males and 7 females. Patients were 2-26 years of age, with peak ages between 3 and 6 years. All tumors were WHO grade IV. Matched peripheral blood lymphocytes were available for 6 of 22 cases, and 18 samples were compared with unmatched normal cerebellum DNA.

Nine testicular tumors included 6 seminomas and three nonseminomas. Samples were obtained from the Norwegian Radium Hospital and from the Helsinki University Central Hospital. Patients were 21-77 years (average of 41 years). Adjacent testicular tissue was available for 7 of 9 cases, and 2 samples were compared with 4 samples of testicular DNA used in the matched sets.

G. Loss of spots from RLGS profiles is due to methylation

In comparing RLGS profiles of DNAs from different tumors with control, healthy tissue DNAs, loss of a fragment or spot from an RLGS profile (Fig. 1A) was frequently detected. Loss of such a spot could be due to either methylation of DNA sequences at the NotI site giving rise to that spot, or to deletion of DNA surrounding that NotI site from the genome of the tumor. The relative contribution of each mechanism was assessed by using clones from the NotI/EcoRV genomic library, specific for lost spots, as probes in Southern blotting studies. In Fig. 2A, a section of an RLGS profile, from normal, healthy tissue was compared with tumor tissue from two gliomas, J7 and J16. This RLGS section contains spot 3C1. In tumor J16, spot 3C1 is absent from the RLGS profile. If there was a deletion of DNA surrounding the NotI site, however, the expected result in the Southern blot would be either no hybridization of the probe to the J16 genomic DNA or hybridization to a band of a size different from those detected in the lane containing normal, healthy tissue DNA digested with NotI plus EcoRV, and tumor tissue DNA digested with EcoRV alone. This result is not seen. These results show, therefore, that DNA corresponding to a missing 3C1 spot in J16 glioma DNA is present in the genome, as shown by the Southern hybridization result.

Likewise, DNA corresponding to specific RLGS spots missing in certain leukemias (Fig. 2B) and neuroectodermal tumors of childhood (Fig. 2C) are found to be present when these DNA are analyzed by Southern blotting. Overall, in 26 tumors where specific spots in RLGS profiles were missing DNA corresponding to the spot, was found to be present in the genome by Southern blotting. These results show that loss of spots on RLGS profiles is due to methylation of the corresponding NotI site and not deletion from the genome of DNA representing that spot. Therefore, methylation is the predominant mechanism underlying loss of spots from RLGS profiles.

H. Heterogeneity in CpG-island methylation across tumors.

To compare the overall pattern of methylated CpG islands among different tumors of the same tumor type, 1,184 spots in each of 98 tumors (and their non-tumorigenic controls) were analyzed by RLGS. The analysis was performed by determining the number of RLGS spots lost, or of decreased intensity, as compared to the controls. Each lost spot or spot of decreased

intensity is equivalent to one methylated CpG island. For each tumor type, the number of methylated CpG islands in each individual tumor, as compared to controls, was plotted (Fig. 3). These data showed that breast, head and neck, and testicular tumors had relatively low levels of methylation, with many such tumors showing no methylation. Colon tumors, gliomas, acute myeloid leukemias and primitive neuroectodermal tumors (PNETs) had a much higher frequency of methylation. Nonparametric comparison (Kruskal-Wallis procedure) of the methylation frequencies of the various tumor types showed significant differences between them ($\chi^2=56.9$, $P<0.0001$).

Within a tumor type, the range of methylated CpG islands in individual tumors was variable. The data (Fig. 3) are not consistent with chance variation between tumors because, in the absence of heterogeneity, the variance of the methylation frequency would not be expected to be greater than the mean¹. A formal test of this overdispersion was performed for each tumor type and the results are shown in Fig. 3 as a superimposition of the expected Poisson distribution on the dot plots. These data showed that aberrant methylation of CpG islands can be quantitatively different in individual tumors within a tumor type and more pronounced overall in particular tumor types.

I. Subsets of CpG islands were preferentially methylated in tumors

Through analysis of the RLGS spots lost in different tumors, it was determined that certain spots on the RLGS gels were lost in multiple tumors. This means that specific CpG dinucleotides were methylated in more than one tumor. This is shown in Fig. 4 where the number of tumors within a specific tumor type that had a particular CpG island methylated are shown.

To test the hypothesis that methylation of these common CpG islands was not random, a standard goodness-of-fit test was used.² This can be seen in the plots of Fig. 4 where the black

¹Heterogeneity of methylation frequencies across samples was assessed within each tumor type by a standard test for evidence that the variance in methylation frequency exceeds the mean. This test is motivated by the Poisson approximation, which applies even if the frequencies of methylation vary across CpG islands. Moreover, a simple result from the binomial distribution shows that the test is conservative, because under homogeneity the population variance cannot exceed the mean.

²Under the null hypothesis of equal methylation frequencies for each CpG island, a goodness-of-fit test (χ^2) was applied to the observed versus expected frequencies of islands exhibiting

line of each plot shows the expected distributions if methylation of specific CpG islands in multiple tumors was random. It can be seen from Fig. 4 that for breast tumors, colon tumors, gliomas, acute myeloid leukemias and childhood PNETs, the actual distributions were significantly different ($P < 0.0001$) from the theoretical distributions indicative of randomness.

5 Similarly, the results for head and neck tumors were significant ($P < 0.025$). The results for testicular tumors ($P = 0.365$) were not significant. However, tumors of this type have a low overall methylation frequency and larger sample sizes are needed. Overall, the data indicate that the patterns of CpG island methylation in tumors is not random.

10 J. Frequencies of aberrant CpG-island methylation of shared and tumor-type-specific targets

Because the data have shown that they are methylated in a nonrandom fashion, CpG islands that are methylated at a high frequency in one or more tumor types can be used for diagnosis of tumors. From analysis of 98 tumors using NotI/EcoRV RLGS analysis, a number of spots that are absent or of decreased intensity, as compared to control healthy tissue DNA, have been found. Table I lists these spots. Each fragment (CpG island) is identified in three ways in the table. First, the location of each CpG island is designated as the distance (in cm) migrated during electrophoresis, from the gel origin, in both the first dimension and the second dimension. Second, each CpG island is given a three-variable designation (Y coordinate, X coordinate, fragment number). The X coordinate indicates horizontal direction on the two-dimensional RLGS profile and is a letter from B-G. The Y coordinate indicates vertical direction and is a number from 1-5. Together, an X and Y designation divide the RLGS profile into 28 sections. Within each section, the spots/fragments are given a number. Such a profile is available at: <http://pandora.med.ohio-state.edu/masterRLGS/>. Third, the partial DNA sequence of individual spots has been determined by sequencing of library clones corresponding to each spot. These sequences are shown in the attached Sequence Listing and have been assigned SEQ ID NOS. from 1 to 82.

The diagnostic NotI/EcoRV spots are of two types (Fig. 1). The first type of spot is absent or of decreased intensity in a single tumor type. For example, the NotI site that is part of the CpG island designated 2.B.53, is methylated only in head and neck tumors. Similarly, the NotI site of CpG island 2.F.2 is methylated only in breast tumors.

methylation in multiple tumors within each tumor type.

The second type of spot is absent or of decreased intensity in more than one type of tumor. For example, the NotI/EcoRV spot designated 2.C.24 is missing in gliomas and AMLs. Similarly, the NotI/EcoRV spot designated 3.B.55 is methylated in breast, colon and PNETs.

5 Table I. Diagnostic CpG islands in tumors.

CpG	1st-D	2nd-D	Type³	Methylated
Island¹	(cm)²	(cm)²		In⁴:
2.B.53	36.85	9.25	t	HN
2.C.24	30.3	5.32	s	Abt/Leu
2.C.29	27.8	5.45	s	Leu/Hn
2.C.35	29.45	6.9	s	Abt/Bre/Cln/Leu/Pbt
2.C.54	32.38	9.42	s	Leu/Hn
2.C.57	30.9	8.5	ND	Tst
2.C.58	31.2	9.2	s	Abt/Leu
2.C.59	30.4	9.35	ND	Hn
2.D.10	27.55	5.3	s	Leu/Pbt
2.D.14	24.25	4.47	t	Leu
2.D.20	26.3	5.3	t	Cln
2.D.25	27.15	6.4	ND	Bre
2.D.27	25.65	5.82	ND	Hn
2.D.34	23.62	6.6	s	Leu/Pbt
2.D.40	23.95	7.25	ND	Pbt
2.D.48	26.1	8.1	ND	Leu
2.D.55	24.2	8.3	s	Cln/Leu
2.D.74	23.95	9.35	s	Abt/Bre/Cln/Leu
2.E.20	20.6	5.95	ND	Pbt
2.E.24	19.35	5.7	s	Abt/Leu
2.E.25	18.27	5.65	t	Bre
2.E.30	20.35	6.4	s	Abt/Bre/Leu
2.E.37	21.42	7.1	ND	Bre
2.E.4	21.1	4.48	s	Leu/Pbt
2.E.40	NA	NA	ND	Tst
2.E.61	19.4	8.08	s	Abt/Pbt
2.E.64	20.5	8.35	s	Abt/Cln
2.F.2	17.27	4.72	t	Bre
2.F.41	NA	NA	t	Tst
2.F.50	15.23	7	s	Abt/Leu
2.F.59	17.49	8	ND	Bre
2.F.70	15.88	13.3	s	Pbt/Tst
2.G.10	10.29	4.49	s	Leu/Tst
2.G.108	7.68	7.44	ND	Bre
3.B.30	35.4	12.55	ND	Tst

3.B.36	34.2	11.8	s	Abt/Cln/Leu/Pbt
3.B.55	NA	NA	s	Bre/Cln/Pbt
3.C.01	31.6	9.7	s	Abt/Cln/Leu
3.C.16	27.9	11.8	t	Pbt
3.C.17	29.2	10.57	t	Cln
3.C.30	31.61	10.37	t	Bre
3.C.35	31.6	11.5	t	Pbt
3.C.64	29.1	14.05	ND	Bre
3.D.21	24.2	10.75	t	Leu
3.D.24	23.2	11.03	s	Abt/Leu
3.D.35	26.1	11.65	s	Abt/Cln/Leu/Pbt
3.D.40	23.4	12.26	s	Abt/Cln/Leu
3.D.44	24.45	12.82	t	Leu
3.D.60	27.2	12.4	s	Abt/Cln/Leu
3.E.04	20.4	14.2	s	Hn/Pbt
3.E.50	20.55	10.7	s	Hn/Tst
3.E.55	18.78	10.55	s	Cln/Leu
3.E.57	18.09	10.9	s	Cln/Hn
3.E.59	18.4	9.72	s	Abt/Tst
3.F.16	16.6	9.75	ND	Leu
3.F.2	16.73	9.35	s	Leu/Tst
3.F.50	16.25	11.6	s	Cln/Leu/Tst
3.F.72	16.9	13.7	t	Leu
3.F.82	13.8	13.12	s	Abt/Cln/Leu
3.G.46	9.88	11.5	ND	Bre
3.G.78	10	12.93	ND	Leu/Pbt
4.B.44	33.7	18.53	s	Cln/Hn
4.B.56	33.2	19.45	s	Bre/Leu
4.C.05	30	14.9	ND	Bre
4.C.25	28.62	17	ND	Bre
4.C.42	NA	NA	ND	Tst
4.C.9	30.3	15.3	ND	Bre
4.D.07	22.9	14.5	s	Leu/Tst
4.D.08	23.5	15	s	Abt/Tst
4.D.12	25	14.85	s	Abt/Leu/Tst
4.D.13	24.95	15.3	s	Abt/Bre
4.D.47	27.6	18.25	s	Abt/Leu/Pbt
4.E.53	19.39	18.43	t	Leu
4.F.15	13.25	15.45	t	Cln
4.F.17	14.1	15.6	s	Abt/Bre/Cln
4.F.22	17.56	16.2	s	Cln/Hn/Leu
4.F.6	14.85	14.59	ND	Bre
4.F.69	12.58	18.86	t	Abt
5.D.9	25.17	23.4	t	Hn

5.E.2	20.58	19.5	t	Bre
5.E.25	18.7	21.3	t	Cln
5.E.4	18.45	19.75	s	Abt/Bre/Leu
¹ Y coordinate, X coordinate, fragment number				
² NA, spots too close to analyze.				
³ T, tumor-type specific target of methylation; s, shared target of methylation; ND, not determined.				
⁴ Types of tumor in which CpG island is methylated: Abt, gliomas; Bre, breast; Cln, colon; Hn, head and neck; Leu, acute myeloid leukemia; Pbt, pediatric brain tumors; Tst, testicular germ cell tumors.				

Example 2. Identification of diagnostic markers for lung cancer using AscI and RLGS

Tissue from lung tumors was obtained as surgical tissue samples. Where possible, surrounding non-tumor tissue from the same patient was obtained and used as a control. DNA was isolated from the tissue as described in Example 1. In preparation for RLGS analysis, the ends of the DNA were blocked as described in Example 1. The DNA was then digested with AscI followed by digestion with EcoRV. The AscI restriction enzyme recognizes the sequence 5'GGCGGCC3' and does not cleave said sequence if cytosines within the sequence are methylated. First dimension gel electrophoresis, in-gel digestion with HinfI, second dimension gel electrophoresis and autoradiography were performed as described in Example 1.

RLGS profiles from lung tumor DNA were compared with RLGS profiles obtained from healthy, non-tumor tissue DNA. Spots which were lost or present at reduced intensity in tumor tissue RLGS profiles as compared to profiles obtained from healthy tissue were noted. Eight spots were lost or altered in the RLGS profiles from multiple lung tumor samples. A compilation of such spots is shown in Table II (lung tumors).

DNA sequence information was obtained from the lung cancer-specific spots. This was done by sequencing individual clones of an AscI/EcoRV library that was made from DNA from healthy tissue. Individual clones of this library that corresponded to spots on the AscI/EcoRV RLGS profile were identified by overloading an RLGS gel with DNA from various groups of library clones, as was described earlier in the specification of this application for the NotI/EcoRV library. After individual clones were matched with spots in the AscI/EcoRV profile, the DNA from the spots that were missing in profiles from the lung tumor DNAs were sequenced. Such sequence information is shown in the attached DNA Sequence Listing.

Table II. Diagnostic CpG islands grouped by tumor type.

Library	Tumor type	Tumor type specific (+), shared (-), or not determined (ND) ¹	CpG island designation`
NotI/EcoRV	Breast	+	2.E.25, 2.F.2, 3.C.30, 5.E.2
		-	3.B.55, 4.B.56, 4.D.13, 4.F.17, 2.D.74, 2.C.35, 2.E.30, 5.E.4
		ND	2.D.25, 2.E.37, 2.F.59, 2.G.108, 3.C.64, 3.G.46, 4.C.05, 4.C.25, 4.C.9, 4.F.6
NotI/EcoRV	Colon	+	2.D.20, 3.C.17, 4.F.15, 5.E.25
		-	3.E.57, 4.B.44, 4.F.22, 2.D.55, 3.E.55, 3.F.50, 3.B.55, 4.F.17, 2.D.74, 2.C.35, 2.E.64, 3.C.01, 3.D.40, 3.D.60, 3.F.82, 3.B.36, 3.D.35
		ND	---
NotI/EcoRV	Glioma	+	4.F.69
		-	4.D.13, 4.F.17, 2.D.74, 2.C.35, 2.E.30, 5.E.4, 2.E.64, 3.C.01, 3.D.40, 3.D.60, 3.F.82, 3.B.36, 3.D.35, 2.C.24, 2.C.58, 2.E.24, 2.F.50, 3.D.24, 4.D.47, 4.D.12, 2.E.61, 3.E.59, 4.D.08
		ND	---
NotI/EcoRV	Head & neck	+	2.B.53, 5.D.9
		-	2.C.29, 2.C.54, 3.E.04, 3.E.50, 3.E.57, 4.B.44, 4.F.22
		ND	2.C.59, 2.D.27
NotI/EcoRV	Acute myelogenous Leukemia	+	2.D.14, 3.D.21, 3.D.44, 3.F.72, 4.E.53, 2.C.29, 2.C.54

		-	2.D.10, 2.D.34, 2.E.4, 2.G.10, 3.F.2, 4.D.07, 4.F.22, 2.D.55, 3.E.55, 3.F.50, 2.E.64, 3.C.01, 3.D.40, 3.D.60, 3.F.82, 3.B.36, 3.D.35, 3.C.01, 3.D.40, 3.D.60, 3.F.82, 3.B.36, 3.D.35, 2.C.24, 2.C.58, 2.E.24, 2.F.50, 3.D.24, 4.D.47, 4.D.12
		ND	2.D.48, 3.F.16, 3.G.78, 4.B.56
NotI/EcoRV	Pediatric neuroectodermal tumor of childhood	+	3.C.16, 3.C.35, 3.E.04
		-	2.D.10, 2.D.34, 2.E.4, 3.B.55, 2.C.35, 3.B.36, 3.D.35, 4.D.47, 2.E.61
		ND	2.D.40, 2.E.20, 3.G.78
NotI/EcoRV	Testicular	+	2.F.41
		-	2.G.10, 3.F.2, 4.D.07, 3.E.50, 3.F.50, 4.D.12, 3.E.59, 4.D.08
		ND	2.C.57, 2.E.40, 3.B.30, 4.C.42
AscI/EcoRV	Lung	+	
		-	
		ND	A.2.F.45, A.2.F.50, A.2.F.67, A.3.F.38, A.4.D.30, A.4.D.36, A.4.E.32, A.5.E.28 ²

¹ND, not determined. Indicates that the designated CpG island was methylated in the indicated tumor type but its methylation in other tumor types was not determined.

²The "A" preceding the X, Y, number designation for the CpG islands indicates that these islands are from the AscI/EcoRV RLGS profile.

Example 3. Design of primers for cancer diagnosis

Primers are designed for diagnosis of cancer using methylation-specific PCR (MSR). The primers are designed to amplify regions of the human genome whose sequences are contained within the library clones disclosed in this application. Two sets of primers are needed for each library clone whose DNA sequence is to be used for diagnosis of cancer. Each primer

set is designed to amplify the same region of the genome, said region beginning at the end of a library clone containing the methylation-sensitive restriction enzyme recognition site (i.e., the NotI site for the library described in Example 1; the AscI site for the library described in Example 2) and ending at a region contained within the clone up to 200 nucleotides from the methylation-sensitive restriction enzyme recognition site.

The first set of primers is designed to amplify template genome DNA whose cytosine residues are not methylated and, after bisulfite treatment, the cytosines of said genome DNA are converted to uracil. The second set of primers is designed to amplify template genome DNA which is methylated on cytosines that comprise CpG dinucleotides. Such methylated cytosines are unaffected by bisulfite treatment. Therefore, by using two sets of primers, one set that will amplify only unmethylated DNA and another set that will amplify only methylated DNA, methylation state of the template DNA can be determined. Such methylation state can be diagnostic for cancer.

The primers used for MSR are designed to be from 15 to 34 nucleotides in length and contain within their sequence either CpG dinucleotides or dinucleotides complementary to CpG dinucleotides that have been treated with bisulfite. It is preferred that the 3' ends of primers used to amplify unmethylated DNA are CpA dinucleotides. It is preferred that the 3' ends of primers used to amplify methylated DNA are CpG dinucleotides.

For each library clone to be used diagnostically, the first set of primers are designed to amplify genome DNA that is not methylated. After treatment of such genome DNA with bisulfite, all such unmethylated cytosines are converted to uracil. PCR primers that will use such DNA as a template and amplify it, will have adenine residues which are complimentary to these uracils.

For the first set of primers, the 5' end of one of the primers begins at the end of the library clone containing the methylation-sensitive restriction enzyme recognition site. The sequence of this primer is identical in sequence to the strand of the template which has its 5' end as part of the methylation-sensitive restriction enzyme site, except that guanine residues are replaced with adenine residues. The adenines allow the primer to hybridize with the template strand in which cytosines have been converted to uracils by bisulfite. This primer extends to a length of between 15 and 32 total nucleotides. Preferably, the 3' end of said primer ends with a

CpA dinucleotide, the adenine of said dinucleotide hybridizing to a uracil which, before bisulfite treatment, had been a cytosine that comprised a CpG dinucleotide.

The diagram below shows implementation of these rules to select a primer that can be used to amplify clone 2.B.53 of the NotI/EcoRV library (Table I and attached sequence listing).

- 5 In the diagram, I shows the end of the 2.B.53 clone containing the methylation-sensitive NotI site (NotI recognition sequence is shown in bold letters). CpG dinucleotides are shaded. To amplify a region of this clone rightward of the NotI site, the first primer is identical to the top strand of the duplex shown in I. However, since bisulfite treatment of the DNA in I converts cytosines to uracils, guanines within the PCR primer must be replaced with adenines. II shows the sequence of the bottom strand of I after bisulfite treatment converts cytosines to uracils. A primer complementary to the bisulfite-treated bottom strand has the sequence shown in III.

I

15 5' **GCGCGCGCGG**TTAGCTTCTCCTGT**CCGAA**CGCAGGG-----
3' **CGCGCGCGCG**CAATCGAAGAGGACAG**CGCTT**CGGTCCC-----

II

20 3' UGUUGGUGUUAATUGAAGAGGAUAGGUTTGUGTUUU-----

III

25 5' ACAACCACAATTAAGTTCTCCTATCCAAACA 3'

III shows the entire sequence of one of the two primers used to amplify unmethylated genome DNA corresponding to library clone 2.B.53. This primer encompasses 5 CpG dinucleotides, as shown by the shading in I above. Encompassment of 2 or more such CpG dinucleotides is preferred so that this primer will not hybridize to a bisulfite-treated template which contains methylated cytosines. The 3' end of the primer shown in III ends in a CpA dinucleotide. This is also preferred in order to provide maximal discrimination of the primer between methylated and unmethylated template DNA in MSR. The primer shown in III has a length of 31 nucleotides.

The second primer is designed to work with the first primer in PCR amplification such that a fragment of less than about 200 base pairs is amplified. Therefore, this primer is made to a sequence rightward of the sequence shown in I. The sequence of this primer is complementary in sequence to the strand of the template which has its 5' end as part of the methylation-sensitive

restriction enzyme site, except that guanine residues are replaced with adenine residues. This primer is preferably between 15 and 32 nucleotides in length. This primer is also designed to preferably encompass 2 or more CpG dinucleotides. Preferably, the 3' end of said primer ends with a CpA dinucleotide.

5 The diagram below shows implementation of these rules to select a primer that can be used to amplify unmethylated genome DNA corresponding to clone 2.B.53 of the NotI/EcoRV library. IV shows a region of the 2.B.53 clone about 70 nucleotides rightward of the sequence in I of the earlier diagram. The CpG dinucleotides within the sequence are shaded. To amplify a region leftward of this region, this second primer must be complementary to the top strand of the duplex shown in IV. However, bisulfite treatment of the DNA in IV converts cytosines to uracils. A primer complementary to this bisulfite-treated top strand has the sequence shown in VI.

IV

15 5' -----GGAGT **CGCG**GT **CGCG**GGAGGCTG **CGC** **CGCG**CAC **CGA**-----3'
 3' -----CCTCA **CGCG**CA **CGCG**CCTCCGAC **CGC** **CGCG**GTGGCT-----5'

V

20 5' -----GGAGTUGUGGTUGUGGGAGGUTGUGUUGUGUAUUGA-----3'

VI

3' ACACCAACACCCTCCAACACAACACATAACT 5'

25 VI shows the entire sequence of the second primer used to amplify unmethylated genome DNA corresponding to library clone 2.B.53. This primer encompasses 8 CpG dinucleotides, as shown by the shading in IV. Encompassment of 2 or more such CpG dinucleotides is preferred. The 3' end of the primer shown in VI ends in a CpA dinucleotide. This is also preferred. The primer shown in VI has a length of 31 nucleotides. Together, the first and second primers amplify a PCR fragment of 128 base pairs in length.

30 The above primers amplify genome DNA that does not contain 5-methylcytosine. The above primers will not amplify genome DNA containing 5-methylcytosines because 5-methylcytosines are not converted to uracils by bisulfite treatment. The two primers already described (III and VI), therefore, will not be complementary to bisulfite-treated genome DNA which is methylated.

5

15

20

19

25

30

A second primer is designed to work with the primer shown in IX to amplify methylated genome template DNA. Design of such a primer is shown below. In the diagram, X shows the same region of clone 2.B.53 (approximately 70 nucleotides rightward of the sequences shown in VII) that is shown in IV. Treatment of the DNA in X with bisulfite produces a top strand with the sequence shown in XI. In XI, only unmethylated cytosines are converted to uracil by bisulfite.

X

5' -----GGAGT**CGCG**GT**CGCG**GGAGGCTG**CGCG**CAC**CGA**-----3'
 3' -----CCTCAG**CGCG**CAG**CGCG**CCTCCGAC**CGCG**GTGG**CT**-----5'

XI

5' -----GGAGTCGCGGTCGCGGGAGGUTGCGUCGCGUAUCGA-----3'

XII

3' GCGCCAGCGCCCTCCAACGCAGCGCATAGCT 5'

A primer complementary to the bisulfite-treated top strand (XI) has the sequence shown in XII. Said primer will prime PCR amplification of sequences leftward of those shown in X. The primer shown in XII encompasses 8 CpG dinucleotides. Encompassment of 2 or more such CpG dinucleotides is preferred. The 3' end of the primer shown in XII ends in a CpG dinucleotide. This is also preferred. The primer shown in XII has a length of 31 nucleotides. Together, the first (IX) and second primers (XII) of the second set amplify a PCR fragment of 128 base pairs in length.

Example 4. Use of oligonucleotides to diagnose cancer

The library clones, and DNA sequences within, can be used to detect DNA methylation in a genome at the specific sequences identified by the sequences within the clone. Such detection can be diagnostic for cancer. Various methods can be used for such diagnosis.

A. Diagnosis of cancer using methylation-sensitive restriction enzymes followed by Southern blot

Cleavage or lack of cleavage by a methylation-sensitive restriction enzyme at a specific restriction enzyme recognition site can be detected by a probe for the specific recognition site,

using Southern blotting. Genomic DNAs were isolated (as described in Example 1) from tumor tissue from a patient with acute myelogenous leukemia (AML). Cells from the same patient after chemotherapy and remission of the disease served as a source of control, healthy tissue DNA. The AML and control DNAs were designated as 26T and 26N, respectively. The DNAs were digested with NotI and EcoRV for 4 hours and then electrophoresed through a 0.8% agarose gel. DNA within the gel was depurinated by soaking the gel in 0.2 N HCl for 10 min. The gel was equilibrated in transfer solution (0.5 N NaOH, 1 M NaCl) for 10 min. and then blotted to Zeta Bind-GT nylon membranes (Bio-Rad). Blots were crosslinked with UV light, baked in a vacuum oven and then prehybridized for 1 hour at 65°C in a solution of 7% SDS, 500 mM sodium phosphate buffer (pH 7.2) and 1 mM EDTA. The blot was hybridized overnight at 65°C in prehybridization solution with 10 ng of α -³²P-labeled probe at a specific activity of 10⁸-10⁹ dpm/μg. The DNA probe used was the 2.C.40 clone from the NotI/EcoRV 2.C.40 library. The purified NotI/EcoRV fragment (50 ng) was labeled with [α -³²P]dCTP by random priming using the Prime-It II random-prime labeling kit (Stratagene). The blot was washed with two quick rinses at 65°C in wash solution 1 (100 mM sodium phosphate buffer, pH 7.2, 0.1% SDS), followed by one 30 min. wash at 65°C in wash solution 1. The blot was next washed for 30 min. at 65°C in wash solution 2 (40 mM sodium phosphate buffer, pH 7.2, 0.1% SDS). Bands were visualized by autoradiography using Kodak X-OMAT AR film.

Fig. 2B shows the data. The first 2 lanes of the autoradiograph are relevant. The first lane, labeled 26N is the normal, healthy tissue DNA cleaved with both NotI and EcoRV. The 26N lane shows a band near the bottom of the autoradiograph labeled "NotI/EcoRV." This is fragment resulting when the NotI site present in the 2.C.40 clone is unmethylated. The adjacent lane, labeled "26T," is the tumor tissue DNA cleaved with both NotI and EcoRV. It is seen that this band, labeled "EcoRV," does not migrate as fast as did the 26N band. The reason is that the NotI site present in the 2.C.40 clone is methylated and the NotI enzyme was unable to cleave at this site.

B. Diagnosis of cancer using methylation-specific PCR (MSR)

MSR is a technique whereby DNA is amplified by PCR dependent upon the methylation state of the DNA. In this example, the specific areas of the genome whose methylation status is to be determined are the regions at the ends of the CpG islands that are demarcated by the

methylation-sensitive restriction enzyme recognition sequence. In the case of the NotI/EcoRV RLGS profiles, this is the NotI site. In the case of the AscI/EcoRV RLGS profiles, this is the AscI site, at the end of each clone.

For the purposes of this example, the methylation status of genomic sequences corresponding to the NotI site of clone 2.B.53 of the NotI/EcoRV library is examined. Genomic DNA is first isolated from normal tissue and from tumor tissue, as described in Example 1. This DNA is then treated with bisulfite. This is done by taking 1 µg of genomic DNA in a volume of 50 µl and denaturing said DNA in a final concentration of 0.2 M NaOH. Thirty microliters of 10 mM hydroquinone and 520 µl of 3 M sodium bisulfite, at pH 5.0, are added, mixed and incubated under mineral oil at 50°C for 16 hours. The modified DNA is then purified using the Wizard DNA purification resin (Promega) and eluted into 50 µl of water. Modification is completed by NaOH (final concentration, 0.3 M) treatment for 5 min. at room temperature, followed by ethanol precipitation. DNA is resuspended in water.

Each genomic DNA is then used in two PCR reactions. One PCR reaction will amplify DNA that is not methylated and has, therefore, been modified by bisulfite. The second PCR reaction will amplify DNA that is methylated. Separate primers are used for each reaction. To determine the methylation status of the NotI site in the genomic DNA which corresponds to the 2.B.53 clone, the two sets of primers described in Example 3 are used. Each PCR reaction contains 1X PCR buffer (16.6 mM ammonium sulfate, 67 mM Tris, pH 8.8, 6.7 mM MgCl₂, 10 mM 2-mercaptoethanol), dNTPs (each at 1.25 mM), primers (300 ng each per reaction), and 50 ng bisulfite-modified DNA in a final volume of 50 µl. Separate control reactions are run which contain DNA that has not been modified by bisulfite. Reactions are hot-started at 95°C for 5 min. before the addition of 1.25 units of *Taq* polymerase. Amplification is carried out for 35 cycles (30 sec at 95°C, 30 sec at the annealing temperature, and 30 sec at 72°C), followed by a final 4 min. extension at 72°C. Each PCR reaction is directly loaded onto nondenaturing 6-8% polyacrylamide gels and electrophoresed. Gels are stained with ethidium bromide and visualized under UV illumination.

If input genomic DNA is not methylated at cytosines within CpG dinucleotides at the NotI site corresponding to the end of the 2.B.53 CpG island clone, the PCR reaction using the primers specific for nonmethylated DNA (primers III and VI in Example 3) will produce an amplification product of 128 base pairs in length. Using the same input genomic DNA, the PCR reaction using

the primers specific for methylated DNA (primers IX and XII in Example 3) will not produce an amplification product.

If input genomic DNA is methylated at cytosines within CpG dinucleotides at the NotI site corresponding to the end of the 2.B.53 CpG island clone, the PCR reaction using the nonmethylation-specific primers will not produce an amplification product. Using the same input genomic DNA, the PCR reaction using the methylation-specific primers will produce an amplification product of 128 base pairs in length.

Example 5. Detection of gene expression

The library clones (Tables I and II) and DNA sequences (attached sequence listing) are useful for determining whether genes encoded within said clones are being transcribed in tumor tissue or cultured cells. To determine transcription, RNA was isolated from five different human glioma cell lines (U87MG, U178, T98G, U251 and LN235) using Trizol (Gibco BRL). Such RNA isolation reagent is known to those skilled in the art. RNAs were quantified using a spectrophotometer and then treated with amplification grade Dnase I (Gibco). The RNA (2 µg) was reverse transcribed by incubation with oligo-dT and random primers in a 20 µl reaction, heated to 70°C for 10 min. and placed on ice. A mix containing 1X reaction buffer (Gibco), DTT (10 mM), dNTPs (0.5 mM each), and RNasin (80 U, Promega) was added to each sample. The samples were divided into two tubes, each containing 19 µl, and incubated at 37°C for 2 min. M-MLV reverse transcriptase (RT, 200 U) was added to one of the two tubes and each was incubated at 37°C for 1 h. DEPC-treated water (30 µl) was added to each sample and heated in boiling water for 5 min.

PCR amplification of the reverse transcribed RNA was then performed. In this study, transcripts encoded by sequences within the 2.C.24 library clone (Table I) were looked for. A computer search using the BLAST program had identified an open reading frame within the sequence of this library clone. PCR primers were made to this region. Primer 1 was 5' TGGTGCTGAAGTCGGTGAA 3'. Primer 2 was 5' GGGCCATCTTCACCATCTG 3'.

These primers (10 pmol of each) were used in 10 µl PCR reactions which contained 1.5 µl of the reverse transcription reaction, 1X reaction buffer, Taq polymerase (0.5 U, Boehringer),

and dNTPs (250 μ M each). For each gene, separate amplification reactions were carried out using RT-positive and RT-negative reactions as template. Amplification was not detected from the RT-negative reactions. The PCR reactions were carried out by heating the samples to 94°C for 5 min and then amplifying for 35 cycles, each cycle consisting of 94°C for 30 sec., a 30 sec. annealing step at 56°C, and 72°C for 45 sec. The reactions were then incubated at 72°C for 7 min and cooled to 4. The sample was then electrophoresed through an agarose gel containing ethidium bromide and PCR products were visualized using an Eagle Eye gel documentation system (Stratagene). The correct identity of the PCR products was confirmed by nucleotide sequencing of both strands.

10 The data showed that no transcripts encoded by this region of the 2.C.24 clone were found in any of the 5 glioma cell lines. Such expressed transcripts are present in RNA obtained from human fetal brain and adult brain.

15 In addition to examination of cell lines, tumor tissue obtained from patient samples can be similarly tested for the presence of transcripts by one skilled in the art. Other techniques to detect transcripts can also be used. Such techniques include, for example, Northern blot hybridization, RNase protection and primer extension assays.

CLAIMS

What is claimed is:

1. A method of identifying CpG islands which are preferentially methylated in malignant cells contained within a tumor or neoplasm, comprising:

- 5 a) digesting genomic DNA obtained from the malignant cells with an infrequently-cutting, methylation-sensitive, restriction enzyme to provide a set of malignant cell restriction fragments;
- b) digesting genomic DNA obtained from non-malignant, control cells with an infrequently-cutting, methylation-sensitive, restriction enzyme to provide a set of control cell
- 10 restriction fragments;
- c) attaching a detectable label to the ends of the malignant cell restriction fragments and the control restriction fragments;
- d) digesting the labeled malignant cell and control cell restriction fragments with a second restriction enzyme;
- 15 e) separating the labeled malignant cell restriction fragments and the labeled control cell restriction fragments, wherein the malignant cell restriction fragments and the control cell restriction fragments are separated by electrophoresis on two different gels;
- f) digesting the restriction fragments in each of said gels with a third restriction enzyme;
- 20 g) electrophoresing the restriction fragments in each of said gels in a direction perpendicular to the first direction to provide a first pattern of detectable malignant cell restriction fragments and a second pattern of detectable control cell restriction fragments; and
- h) comparing the first pattern to the second pattern to identify diagnostic control cell restriction fragments in said second pattern which are absent or exhibit a decreased intensity in
- 25 the first pattern, wherein said diagnostic control cell restriction fragments comprise a CpG island that is unmethylated in the DNA of the control cells and methylated in the DNA of the malignant cells.
2. The method of claim 1 further comprising the step of determining the sequence of at least a portion of a diagnostic control cell restriction fragment, wherein said portion is located at or
- 30 near an end of the fragment .
3. The method of claim 1 further comprising the step of obtaining a clone from a DNA

library which comprises a diagnostic control cell restriction fragment.

4. A method of preparing a polynucleotide or oligonucleotide for characterizing tissue obtained from a subject suspected of having cancer, comprising:

synthesizing a polynucleotide or oligonucleotide which comprises a sequence which is identical to or substantially complementary to a target sequence on one of the strands of a diagnostic control fragment identified according to the method of claim 2, wherein said target sequence comprises at least two CpG dinucleotides, wherein said oligonucleotide is from 15 to 34 nucleotides in length, and wherein said polynucleotide is from 35 to 2000 nucleotides in length.

5. The method of claim 4 wherein said target sequence is located at or near the control restriction fragment end which was cleaved by the methylation-sensitive, restriction enzyme.

6. The method of claim 4 wherein the target sequence is located from about 100 nucleotides to about 500 nucleotides downstream of the control restriction fragment end that was cleaved by the methylation-sensitive, restriction enzyme.

7. The method of claim 4 wherein the control restriction fragment comprises a sequence selected from the group consisting of SEQ. ID. NO.:1, SEQ. ID. NO.:2, SEQ. ID. NO.:3, SEQ. ID. NO.:4, SEQ. ID. NO.:5, SEQ. ID. NO.:6, SEQ. ID. NO.:7, SEQ. ID. NO.:8, SEQ. ID. NO.:9, SEQ. ID. NO.:10, SEQ. ID. NO.:11, SEQ. ID. NO.:12, SEQ. ID. NO.:13, SEQ. ID. NO.:14, SEQ. ID. NO.:15, SEQ. ID. NO.:16, SEQ. ID. NO.:17, SEQ. ID. NO.:18, SEQ. ID. NO.:19, SEQ. ID. NO.:20, SEQ. ID. NO.:21, SEQ. ID. NO.:22, SEQ. ID. NO.:23, SEQ. ID. NO.:24, SEQ. ID. NO.:25, SEQ. ID. NO.:26, SEQ. ID. NO.:27, SEQ. ID. NO.:28, SEQ. ID. NO.:29, SEQ. ID. NO.:30, SEQ. ID. NO.:31, SEQ. ID. NO.:32, SEQ. ID. NO.:33, SEQ. ID. NO.:34, SEQ. ID. NO.:35, SEQ. ID. NO.:36, SEQ. ID. NO.:37, SEQ. ID. NO.:38, SEQ. ID. NO.:39, SEQ. ID. NO.:40, SEQ. ID. NO.:41, SEQ. ID. NO.:42, SEQ. ID. NO.:43, SEQ. ID. NO.:44, SEQ. ID. NO.:45, SEQ. ID. NO.:46, SEQ. ID. NO.:47, SEQ. ID. NO.:48, SEQ. ID. NO.:49, SEQ. ID. NO.:50, SEQ. ID. NO.:51, SEQ. ID. NO.:52, SEQ. ID. NO.:53, SEQ. ID. NO.:54, SEQ. ID. NO.:55, SEQ. ID. NO.:56, SEQ. ID. NO.:57, SEQ. ID. NO.:58, SEQ. ID. NO.:59, SEQ. ID. NO.:60, SEQ. ID. NO.:61, SEQ. ID. NO.:62, , SEQ. ID. NO.:63, SEQ. ID. NO.:64, SEQ. ID. NO.:65, SEQ. ID. NO.:66, SEQ. ID. NO.:67, SEQ. ID. NO.:68, SEQ. ID. NO.:69, SEQ. ID. NO.:70, SEQ. ID. NO.:71, SEQ. ID. NO.:72, SEQ. ID. NO.:73, SEQ. ID. NO.:74, SEQ. ID. NO.:75, SEQ. ID. NO.:76, SEQ. ID. NO.:77, SEQ. ID. NO.:78, SEQ. ID. NO.:79,

SEQ. ID. NO.:80, SEQ. ID. NO.:81, SEQ. ID. NO.:82, SEQ ID NO: 83, SEQ ID NO. 84, SEQ ID. NO. 85, SEQ. ID. NO. 86, SEQ. ID. NO. 87, SEQ. ID. NO. 88, SEQ. ID. NO. 89, SEQ. ID. NO. 90, SEQ. ID. NO. 91, SEQ ID. NO. 92, and SEQ. ID. NO. 93.

8. An isolated polynucleotide or oligonucleotide for characterizing cells that are obtained from a subject suspected of having a cancer which is associated with methylation of one or a plurality of CpG islands in the genomic DNA of malignant cells, wherein said polynucleotide or oligonucleotide comprises a sequence which is identical to or complementary to a target sequence on one of the strands of a diagnostic control fragment identified according to the method of claim 2, wherein said target sequence comprises at least two CpG dinucleotides, wherein said oligonucleotide is from 15 to 34 nucleotides in length; and wherein said polynucleotide is from 35 to 3000 nucleotides in length.

9. The isolated polynucleotide or oligonucleotide of claim 8 wherein said target sequence is located at or near the control restriction fragment end which was cleaved by the methylation sensitive restriction enzyme.

10. The isolated polynucleotide or oligonucleotide of claim 8 wherein the target sequence is located from about 100 nucleotides to about 500 nucleotides downstream of the control restriction fragment end that was cleaved by the methylation-sensitive, restriction enzyme.

11. An isolated polynucleotide or oligonucleotide for characterizing cells which are obtained from a subject suspected of having a cancer which is associated with methylation of one or a plurality of CpG islands in the genomic DNA of malignant cells, wherein said polynucleotide or oligonucleotide comprises a sequence which is identical to or complementary to a modified target sequence on one of the strands of a diagnostic control fragment identified according to the method of claim 2, wherein said modified target sequence is derived from a target sequence that has been modified by treatment with sodium bisulfite, wherein said modified target sequence lacks cytosines and comprises at least two UpG dinucleotides, wherein said oligonucleotide is from 15 to 34 nucleotides in length; and wherein said polynucleotide is from 35 to 3000 nucleotides in length.

12. The isolated polynucleotide or oligonucleotide of claim 11 wherein the modified target sequence is derived from a target sequence located at or near the control restriction fragment end that was cleaved by the methylation sensitive restriction enzyme.

13. The isolated polynucleotide or oligonucleotide of claim 11 wherein the modified target

sequence is derived from a target sequence that is located from about 100 nucleotides to about 500 nucleotides downstream of the control restriction fragment end that was cleaved by the methylation-sensitive restriction enzyme.

- 5 14. An isolated polynucleotide for characterizing cells which are obtained from a subject suspected of having a cancer selected from the group consisting of glioma, acute myeloid leukemia, primitive neuroectodermal tumors of childhood, breast cancer, colon cancer, head and neck cancer, testicular cancer and lung cancer; wherein said polynucleotide is from 35 to 3000 nucleotides in length and comprises at least two CpG dinucleotides, and wherein said
- 10 polynucleotide comprise a sequence which is identical to or complementary to a target sequence located within a sequence selected from the group consisting of SEQ. ID. NO.:1, SEQ. ID. NO.:2, SEQ. ID. NO.:3, SEQ. ID. NO.:4, SEQ. ID. NO.:5, SEQ. ID. NO.:6, SEQ. ID. NO.:7, SEQ. ID. NO.:8, SEQ. ID. NO.:9, SEQ. ID. NO.:10, SEQ. ID. NO.:11, SEQ. ID. NO.:12, SEQ. ID. NO.:13, SEQ. ID. NO.:14, SEQ. ID. NO.:15, SEQ. ID. NO.:16, SEQ. ID. NO.:17, SEQ. ID. NO.:18, SEQ. ID. NO.:19, SEQ. ID. NO.:20, SEQ. ID. NO.:21, SEQ. ID. NO.:22, SEQ. ID. NO.:23, SEQ. ID. NO.:24, SEQ. ID. NO.:25, SEQ. ID. NO.:26, SEQ. ID. NO.:27, SEQ. ID. NO.:28, SEQ. ID. NO.:29, SEQ. ID. NO.:30, SEQ. ID. NO.:31, SEQ. ID. NO.:32, SEQ. ID. NO.:33, SEQ. ID. NO.:34, SEQ. ID. NO.:35, SEQ. ID. NO.:36, SEQ. ID. NO.:37, SEQ. ID. NO.:38, SEQ. ID. NO.:39, SEQ. ID. NO.:40, SEQ. ID. NO.:41, SEQ. ID. NO.:42, SEQ. ID. NO.:43, SEQ. ID. NO.:44, SEQ. ID. NO.:45, SEQ. ID. NO.:46, SEQ. ID. NO.:47, SEQ. ID. NO.:48, SEQ. ID. NO.:49, SEQ. ID. NO.:50, SEQ. ID. NO.:51, SEQ. ID. NO.:52, SEQ. ID. NO.:53, SEQ. ID. NO.:54, SEQ. ID. NO.:55, SEQ. ID. NO.:56, SEQ. ID. NO.:57, SEQ. ID. NO.:58, SEQ. ID. NO.:59, SEQ. ID. NO.:60, SEQ. ID. NO.:61, SEQ. ID. NO.:62, , SEQ. ID. NO.:63, SEQ. ID. NO.:64, SEQ. ID. NO.:65, SEQ. ID. NO.:66, SEQ. ID. NO.:67, SEQ. ID. NO.:68, SEQ. ID. NO.:69, SEQ. ID. NO.:70, SEQ. ID. NO.:71, SEQ. ID. NO.:72, SEQ. ID. NO.:73, SEQ. ID. NO.:74, SEQ. ID. NO.:75, SEQ. ID. NO.:76, SEQ. ID. NO.:77, SEQ. ID. NO.:78, SEQ. ID. NO.:79, SEQ. ID. NO.:80, SEQ. ID. NO.:81, SEQ. ID. NO.:82, SEQ ID NO: 83, SEQ ID NO. 84, SEQ ID. NO. 85, SEQ. ID. NO. 86, SEQ. ID. NO. 87, SEQ. ID. NO. 88, SEQ. ID. NO. 89, SEQ. ID. NO. 90, SEQ. ID. NO. 91, SEQ ID. NO. 92, and SEQ. ID. NO. 93.
- 25 15. The isolated polynucleotide of claim 14 wherein said subject is suspected of having glioma and said polynucleotide comprises a sequence which is identical to or complementary to
- 30

a target sequence located within SEQ. ID.NO. 78, SEQ ID NO:71, SEQ ID NO:75, SEQ ID NO:18, SEQ ID NO:4, SEQ ID NO:22, SEQ ID NO:82, SEQ ID NO:27, SEQ ID NO:38, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:59, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:2, SEQ ID NO:7, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:45, SEQ ID NO:72, SEQ ID NO:70, SEQ ID NO:26, SEQ ID NO:54, or SEQ ID NO:69.

16. The isolated polynucleotide of claim 14 wherein said subject is suspected of having acute myeloid leukemia and said polynucleotide comprises a sequence which is identical to or complementary to a target sequence located within SEQ ID NO:10, SEQ ID NO:44, SEQ ID NO:48, SEQ ID NO:58, SEQ ID NO:73, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:9, SEQ ID NO:14, SEQ ID NO:24, SEQ ID NO:33, SEQ ID NO:56, SEQ ID NO:68, SEQ ID NO:76, SEQ ID NO:17, SEQ ID NO:52, SEQ ID NO:57, SEQ ID NO:26, SEQ ID NO:38, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:59, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:38, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:59, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:2, SEQ ID NO:7, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:45, SEQ ID NO:72, SEQ ID NO:16, SEQ ID NO:55, SEQ ID NO:61, SEQ ID NO:63, or SEQ ID NO:70.

17. The isolated polynucleotide of claim 14 wherein said subject is suspected of having a primitive neuroectodermal tumor of childhood and said polynucleotide comprises a sequence which is identical to or complementary to a target sequence located within SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:50, SEQ ID NO:9, SEQ ID NO:14, SEQ ID NO:24, SEQ ID NO:37, SEQ ID NO:4, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:72, SEQ ID NO:26, SEQ ID NO:15, SEQ ID NO:19, or SEQ ID NO:61.

18. The isolated polynucleotide of claim 14 wherein said subject is suspected of having breast cancer and said polynucleotide comprises a sequence identical or complementary to a target sequence which is located within SEQ ID NO:21, SEQ ID NO:28, SEQ ID NO:41, SEQ ID NO:80, SEQ ID NO:37, SEQ ID NO:63, SEQ ID NO:71, SEQ ID NO:75, SEQ ID NO:18, SEQ ID NO:4, SEQ ID NO:22, SEQ ID NO:82, SEQ ID NO:12, SEQ ID NO:23, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:43, SEQ ID NO:60, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:77.

19. The isolated polynucleotide of claim 14 wherein said subject is suspected of having colon cancer and said polynucleotide comprises a sequence which is identical to or complementary to a target sequence located within SEQ ID NO:11, SEQ ID NO:40, SEQ ID NO:74, SEQ ID NO:81,

SEQ ID NO:53, SEQ ID NO:62, SEQ ID NO:76, SEQ ID NO:17, SEQ ID NO:52, SEQ ID NO:57, SEQ ID NO:37, SEQ ID NO:75, SEQ ID NO:18, SEQ ID NO:4, SEQ ID NO:27, SEQ ID NO:38, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:59, SEQ ID NO:36, or SEQ ID NO:46.

20. The isolated polynucleotide of claim 14 wherein said subject is suspected of head and neck cancer and said polynucleotide comprises a sequence which is identical to or complementary to a target sequence located within in SEQ ID NO:1, SEQ ID NO. 79, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:62, SEQ ID NO:76, SEQ ID NO:8, or SEQ ID NO:13.

21. The isolated polynucleotide of claim 14 wherein said subject is suspected of testicular cancer and said polynucleotide comprises a sequence which is identical to or complementary to a target sequence located within in SEQ ID NO. 29, SEQ ID NO:33, SEQ ID NO:56, SEQ ID NO:68, SEQ ID NO:51, SEQ ID NO:57, SEQ ID NO:70, SEQ ID NO:54, or SEQ ID NO:69

22. The isolated polynucleotide of claim 14 wherein said subject is suspected of having a lung cancer and said polynucleotide comprises a sequence which is identical to or complementary to a target sequence located within SEQ ID NO: 83, SEQ ID NO. 84, SEQ ID. NO. 85, SEQ. ID. NO. 86, SEQ. ID. NO. 87, SEQ. ID. NO. 88, SEQ. ID. NO. 89, SEQ. ID. NO. 90, SEQ. ID. NO. 91, SEQ ID. NO. 92, and SEQ. ID. NO. 93.

23. An isolated CpG diagnostic oligonucleotide for characterizing cells which are obtained from a subject suspected of having a cancer selected from the group consisting of glioma, acute myeloid leukemia, primitive neuroectodermal tumors of childhood, breast cancer, colon cancer, head and neck cancer, testicular cancer and lung cancer; wherein said oligonucleotide is from 15 to 34 nucleotides in length and comprises at least two CpG dinucleotides, and wherein said oligonucleotide comprise a sequence which is identical to a target sequence located within a region extending from nucleotide 1 through nucleotide 99 of a sequence selected from the group consisting of SEQ. ID. NO.:1, SEQ. ID. NO.:2, SEQ. ID. NO.:3, SEQ. ID. NO.:4, SEQ. ID. NO.:5, SEQ. ID. NO.:6, SEQ. ID. NO.:7, SEQ. ID. NO.:8, SEQ. ID. NO.:9, SEQ. ID. NO.:10, SEQ. ID. NO.:11, SEQ. ID. NO.: 12, SEQ. ID. NO.:13, SEQ. ID. NO.:14, SEQ. ID. NO.:15, SEQ. ID. NO.: 16, SEQ. ID. NO.:17, SEQ. ID. NO.:18, SEQ. ID. NO.: 19, SEQ. ID. NO.:20, SEQ. ID. NO.:21, SEQ. ID. NO.:22, SEQ. ID. NO.:23, SEQ. ID. NO.:24, SEQ. ID. NO.:25, SEQ. ID. NO.:26, SEQ. ID. NO.:27, SEQ. ID. NO.:28, SEQ. ID. NO.:29, SEQ. ID. NO.:30,

SEQ. ID. NO.:31, SEQ. ID. NO.:32, SEQ. ID. NO.:33, SEQ. ID. NO.:34, SEQ. ID. NO.:35,
 SEQ. ID. NO.:36, SEQ. ID. NO.:37, SEQ. ID. NO.:38, SEQ. ID. NO.:39, SEQ. ID. NO.:40,
 SEQ. ID. NO.:41, SEQ. ID. NO.:42, SEQ. ID. NO.:43, SEQ. ID. NO.:44, SEQ. ID. NO.:45,
 SEQ. ID. NO.:46, SEQ. ID. NO.:47, SEQ. ID. NO.:48, SEQ. ID. NO.:49, SEQ. ID. NO.:50,
 5 SEQ. ID. NO.:51, SEQ. ID. NO.:52, SEQ. ID. NO.:53, SEQ. ID. NO.:54, SEQ. ID. NO.:55,
 SEQ. ID. NO.:56, SEQ. ID. NO.:57, SEQ. ID. NO.:58, SEQ. ID. NO.:59, SEQ. ID. NO.:60,
 SEQ. ID. NO.:61, SEQ. ID. NO.:62, , SEQ. ID. NO.:63, SEQ. ID. NO.:64, SEQ. ID. NO.:65,
 SEQ. ID. NO.:66, SEQ. ID. NO.:67, SEQ. ID. NO.:68, SEQ. ID. NO.:69, SEQ. ID. NO.:70,
 SEQ. ID. NO.:71, SEQ. ID. NO.:72, SEQ. ID. NO.:73, SEQ. ID. NO.:74, SEQ. ID. NO.:75,
 10 SEQ. ID. NO.:76, SEQ. ID. NO.:77, SEQ. ID. NO.:78, SEQ. ID. NO.:79, SEQ. ID. NO.:80,
 SEQ. ID. NO.:81, SEQ. ID. NO.:82; SEQ ID NO: 83, SEQ ID NO. 84, SEQ ID. NO. 85, SEQ.
 ID. NO. 86, SEQ. ID. NO. 87, SEQ. ID. NO. 88, SEQ. ID. NO. 89, SEQ. ID. NO. 90, SEQ. ID.
 NO. 91, SEQ ID. NO. 92, and SEQ. ID. NO. 93; or a sequence which is the reverse complement
 of a target sequence located in a region extending from about nucleotide 100 through nucleotide
 15 500 in said SEQ ID NO.

24. The isolated oligonucleotide of claim 23 wherein said subject is suspected of having
 glioma and said oligonucleotide comprises a sequence which is identical to a target sequence
 located within a region extending from nucleotide 1 through nucleotide 99 in SEQ ID 78, SEQ
 ID NO:71, SEQ ID NO:75, SEQ ID NO:18, SEQ ID NO:4, SEQ ID NO:22, SEQ ID NO:82,
 20 SEQ ID NO:27, SEQ ID NO:38, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:59, SEQ ID
 NO:36, SEQ ID NO:46, SEQ ID NO:2, SEQ ID NO:7, SEQ ID NO:20, SEQ ID NO:30, SEQ ID
 NO:45, SEQ ID NO:72, SEQ ID NO:70, SEQ ID NO:26, SEQ ID NO:54, or SEQ ID NO:69, or
 said oligonucleotide comprises a sequence which is the reverse complement of a target sequence
 located in a region extending from about nucleotide 100 through nucleotide 500 in said SEQ ID
 25 NO.

25. The isolated oligonucleotide of claim 23 wherein the subject is suspected of having acute
 myeloid leukemia; and said oligonucleotide comprises a sequence which is identical to a target
 sequence located within a region extending from nucleotide 1 through nucleotide 99 SEQ ID
 NO:10, SEQ ID NO:44, SEQ ID NO:48, SEQ ID NO:58, SEQ ID NO:73, SEQ ID NO:3, SEQ
 30 ID NO:5, SEQ ID NO:9, SEQ ID NO:14, SEQ ID NO:24, SEQ ID NO:33, SEQ ID NO:56, SEQ
 ID NO:68, SEQ ID NO:76, SEQ ID NO:17, SEQ ID NO:52, SEQ ID NO:57, SEQ ID NO:26,

SEQ ID NO:38, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:59, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:38, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:59, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:2, SEQ ID NO:7, SEQ ID NO:20, SEQ ID NO:30, SEQ ID NO:45, SEQ ID NO:72, SEQ ID NO:16, SEQ ID NO:55, SEQ ID NO:61, SEQ ID NO:63, or SEQ ID NO:70, or said oligonucleotide comprises a sequence which is the reverse complement of a target sequence located in a region extending from about nucleotide 100 through nucleotide 500 in said SEQ ID NO.

26. The isolated oligonucleotide of claim 23 wherein said subject is suspected of having primitive neuroectodermal tumors of childhood; and said oligonucleotide comprises a sequence which is identical to a target sequence located within a region extending from nucleotide 1 through nucleotide 99 of SEQ ID NO:39, SEQ ID NO:42, SEQ ID NO:50, SEQ ID NO:9, SEQ ID NO:14, SEQ ID NO:24, SEQ ID NO:37, SEQ ID NO:4, SEQ ID NO:36, SEQ ID NO:46, SEQ ID NO:72, SEQ ID NO:26, SEQ ID NO:15, SEQ ID NO:19, or SEQ ID NO:61, or said nucleotide comprises a sequence which is the reverse complement of a target sequence located in a region extending from about nucleotide 100 through nucleotide 500 in said SEQ ID NO.

27. The isolated oligonucleotide of claim 23 wherein said subject is suspected of having breast cancer; and said oligonucleotide comprises a sequence which is identical to a target sequence located within a region extending from nucleotide 1 through nucleotide 99 of SEQ ID NO:21, SEQ ID NO:28, SEQ ID NO:41, SEQ ID NO:80, SEQ ID NO:37, SEQ ID NO:63, SEQ ID NO:71, SEQ ID NO:75, SEQ ID NO:18, SEQ ID NO:4, SEQ ID NO:22, SEQ ID NO:82, SEQ ID NO:12, SEQ ID NO:23, SEQ ID NO:31, SEQ ID NO:34, SEQ ID NO:43, SEQ ID NO:60, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:67, SEQ ID NO:77, or said oligonucleotide comprises a sequence which is the reverse complement of a target sequence located in a region extending from about nucleotide 100 through nucleotide 500 in said SEQ ID NO.

28. The isolated oligonucleotide of claim 23 wherein said subject is suspected of having colon cancer; and said oligonucleotide comprises a sequence which is identical to a target sequence located within a region extending from nucleotide 1 through nucleotide 99 of SEQ ID NO:11, SEQ ID NO:40, SEQ ID NO:74, SEQ ID NO:81, SEQ ID NO:53, SEQ ID NO:62, SEQ ID NO:76, SEQ ID NO:17, SEQ ID NO:52, SEQ ID NO:57, SEQ ID NO:37, SEQ ID NO:75, SEQ ID NO:18, SEQ ID NO:4, SEQ ID NO:27, SEQ ID NO:38, SEQ ID NO:47, SEQ ID

NO:49, SEQ ID NO:59, SEQ ID NO:36, SEQ ID NO:46 or said oligonucleotide comprises a sequence which is the reverse complement of a target sequence located in a region extending from about nucleotide 100 through nucleotide 500 in said SEQ ID NO.

29. The isolated oligonucleotide of claim 23 wherein said subject is suspected of having head and neck cancer; and said oligonucleotide comprises a sequence which is identical to a target sequence located within a region extending from nucleotide 1 through nucleotide 99 in SEQ ID NO:1, SEQ ID NO. 79, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:53, SEQ ID NO:62, SEQ ID NO:76, SEQ ID NO:8, or SEQ ID NO:13., or said nucleotide a sequence which is the reverse complement of a target sequence located in a region extending from about nucleotide 100 through nucleotide 500 in said SEQ ID NO.

30. The isolated oligonucleotide of claim 23 wherein said subject is suspected of having testicular cancer; and said oligonucleotide comprises a sequence which is identical to a target sequence located within a region extending from nucleotide 1 through nucleotide 99 in SEQ ID NO. 29, SEQ ID NO:33, SEQ ID NO:56, SEQ ID NO:68, SEQ ID NO:51, SEQ ID NO:57, SEQ ID NO:70, SEQ ID NO:54, or SEQ ID NO:69, or said oligonucleotide comprises a sequence which is the reverse complement of a target sequence located in a region extending from about nucleotide 100 through nucleotide 500 in said SEQ ID NO.

31. The isolated oligonucleotide of claim 23 wherein said subject is suspected of having lung cancer; and said oligonucleotide comprises a sequence which is identical to a target sequence located within a region extending from nucleotide 1 through nucleotide 99 of SEQ ID NO SEQ ID NO: 83, SEQ ID NO. 84, SEQ ID. NO. 85, SEQ. ID. NO. 86, SEQ. ID. NO. 87, SEQ. ID. NO. 88, SEQ. ID. NO. 89, SEQ. ID. NO. 90, SEQ. ID. NO. 91, SEQ ID. NO. 92, and SEQ. ID. NO. 93, or said oligonucleotide comprises a sequence which is the reverse complement of a target sequence located in a region extending from about nucleotide 100 through nucleotide 500 in said SEQ ID NO.

32. A method for determining whether cells obtained from a subject suspected of having a cancer are malignant or non-malignant, comprising:

a) digesting DNA which has been isolated from the cells with a methylation-sensitive restriction enzyme to provide a set of restriction fragments;

b) hybridizing said restriction fragments with a CpG diagnostic polynucleotide comprising a sequence which is identical to or complementary to a target sequence on one of the

strands of a diagnostic control fragment identified according to the method of claim 2, wherein said method employed said methylation-sensitive restriction enzyme, wherein said target sequence comprises at least two CpG dinucleotides, wherein said polynucleotide is from 35 to 3000 nucleotides in length, and wherein said reaction is conducted under stringent hybridization conditions; and

c) assaying the reaction products of step b to determine the size or the sequence of the restriction fragment to which the CpG diagnostic polynucleotide has hybridized.

33. The method of claim 32 wherein said subject is suspected of having a cancer selected from the group consisting of glioma, acute myeloid leukemia, primitive neuroectodermal tumors of childhood, breast cancer, colon cancer, head and neck cancer, and testicular cancer;

wherein said DNA sample is digested with NotI; and

wherein said polynucleotide comprises a sequence which is identical to or complementary to a target sequence located in SEQ. ID. NO.:1, SEQ. ID. NO.:2, SEQ. ID. NO.:3, SEQ. ID. NO.:4, SEQ. ID. NO.:5, SEQ. ID. NO.:6, SEQ. ID. NO.:7, SEQ. ID. NO.:8, SEQ. ID. NO.:9, SEQ. ID. NO.:10, SEQ. ID. NO.:11, SEQ. ID. NO.:12, SEQ. ID. NO.:13, SEQ. ID. NO.:14, SEQ. ID. NO.:15, SEQ. ID. NO.:16, SEQ. ID. NO.:17, SEQ. ID. NO.:18, SEQ. ID. NO.:19, SEQ. ID. NO.:20, SEQ. ID. NO.:21, SEQ. ID. NO.:22, SEQ. ID. NO.:23, SEQ. ID. NO.:24, SEQ. ID. NO.:25, SEQ. ID. NO.:26, SEQ. ID. NO.:27, SEQ. ID. NO.:28, SEQ. ID. NO.:29, SEQ. ID. NO.:30, SEQ. ID. NO.:31, SEQ. ID. NO.:32, SEQ. ID. NO.:33, SEQ. ID. NO.:34, SEQ. ID. NO.:35, SEQ. ID. NO.:36, SEQ. ID. NO.:37, SEQ. ID. NO.:38, SEQ. ID. NO.:39, SEQ. ID. NO.:40, SEQ. ID. NO.:41, SEQ. ID. NO.:42, SEQ. ID. NO.:43, SEQ. ID. NO.:44, SEQ. ID. NO.:45, SEQ. ID. NO.:46, SEQ. ID. NO.:47, SEQ. ID. NO.:48, SEQ. ID. NO.:49, SEQ. ID. NO.:50, SEQ. ID. NO.:51, SEQ. ID. NO.:52, SEQ. ID. NO.:53, SEQ. ID. NO.:54, SEQ. ID. NO.:55, SEQ. ID. NO.:56, SEQ. ID. NO.:57, SEQ. ID. NO.:58, SEQ. ID. NO.:59, SEQ. ID. NO.:60, SEQ. ID. NO.:61, SEQ. ID. NO.:62, , SEQ. ID. NO.:63, SEQ. ID. NO.:64, SEQ. ID. NO.:65, SEQ. ID. NO.:66, SEQ. ID. NO.:67, SEQ. ID. NO.:68, SEQ. ID. NO.:69, SEQ. ID. NO.:70, SEQ. ID. NO.:71, SEQ. ID. NO.:72, SEQ. ID. NO.:73, SEQ. ID. NO.:74, SEQ. ID. NO.:75, SEQ. ID. NO.:76, SEQ. ID. NO.:77, SEQ. ID. NO.:78, SEQ. ID. NO.:79, SEQ. ID. NO.:80, SEQ. ID. NO.:81, and SEQ. ID. NO.:82.

34. The method of claim 32 wherein the subject is suspected of having lung cancer; wherein the DNA is digested with AscI;

and wherein said polynucleotide comprises a sequence which is identical to or complementary to a target sequence located in , SEQ ID NO: 83, SEQ ID NO. 84, SEQ ID. NO. 85, SEQ. ID. NO. 86, SEQ. ID. NO. 87, SEQ. ID. NO. 88, SEQ. ID. NO. 89, SEQ. ID. NO. 90, SEQ. ID. NO. 91, SEQ ID. NO. 92, and SEQ. ID. NO. 93.

5 35. A method of determining whether cells contained within a tissue sample obtained from a subject suspected of having cancer are malignant, comprising:

a) treating DNA isolated from the tissue sample with a compound which converts non-methylated cytosines to a different nucleotide base;

10 b) reacting a portion of the treated DNA with a CpG diagnostic oligonucleotide which is complementary to a target sequence which comprises CpG islands that are preferentially methylated in malignant cells of subjects known to have said cancer;

c) reacting a portion of the treated DNA with a modified CpG diagnostic oligonucleotide which is complementary to a modified target sequence in which the cytosines in said target sequence are replaced with the different nucleotide base; and

15 d) assaying the reaction products of step b and step c to determine whether the treated DNA has hybridized with the CpG diagnostic oligonucleotide or the modified CpG diagnostic oligonucleotide; wherein hybridization of the treated DNA with the CpG diagnostic oligonucleotide as opposed to the modified CpG diagnostic oligonucleotide indicates that the DNA has been obtained from malignant cells.

20 36. The method of claim 35 wherein the chemical compound is sodium bisulfite and the non-methylated cytosines are converted to uracil.

37. The method of claim 35 wherein the assay is a polymerase chain reaction,

wherein a portion of the treated DNA is reacted with a first primer set which comprises two diagnostic CpG oligonucleotides; and

25 wherein a portion of the treated DNA is reacted with a second primer set which comprises two modified diagnostic CpG oligonucleotides.

38. The method of claim 35 wherein the subject is suspected of having a cancer selected from the group consisting of glioma, acute myeloid leukemia, primitive neuroectodermal tumors of childhood, breast cancer, colon cancer, head and neck cancer, testicular cancer and lung cancer;

30 and

wherein the CpG diagnostic oligonucleotide comprises a sequence which is identical to a target

sequence located between nucleotide 1 and 100 in a sequence selected from the group consisting said polynucleotide comprises a sequence which is identical to or complementary to a target sequence located in SEQ. ID. NO.:1, SEQ. ID. NO.:2, SEQ. ID. NO.:3, SEQ. ID. NO.:4, SEQ. ID. NO.:5, SEQ. ID. NO.:6, SEQ. ID. NO.:7, SEQ. ID. NO.:8, SEQ. ID. NO.:9, SEQ. ID. NO.:10, SEQ. ID. NO.:11, SEQ. ID. NO.:12, SEQ. ID. NO.:13, SEQ. ID. NO.:14, SEQ. ID. NO.:15, SEQ. ID. NO.:16, SEQ. ID. NO.:17, SEQ. ID. NO.:18, SEQ. ID. NO.:19, SEQ. ID. NO.:20, SEQ. ID. NO.:21, SEQ. ID. NO.:22, SEQ. ID. NO.:23, SEQ. ID. NO.:24, SEQ. ID. NO.:25, SEQ. ID. NO.:26, SEQ. ID. NO.:27, SEQ. ID. NO.:28, SEQ. ID. NO.:29, SEQ. ID. NO.:30, SEQ. ID. NO.:31, SEQ. ID. NO.:32, SEQ. ID. NO.:33, SEQ. ID. NO.:34, SEQ. ID. NO.:35, SEQ. ID. NO.:36, SEQ. ID. NO.:37, SEQ. ID. NO.:38, SEQ. ID. NO.:39, SEQ. ID. NO.:40, SEQ. ID. NO.:41, SEQ. ID. NO.:42, SEQ. ID. NO.:43, SEQ. ID. NO.:44, SEQ. ID. NO.:45, SEQ. ID. NO.:46, SEQ. ID. NO.:47, SEQ. ID. NO.:48, SEQ. ID. NO.:49, SEQ. ID. NO.:50, SEQ. ID. NO.:51, SEQ. ID. NO.:52, SEQ. ID. NO.:53, SEQ. ID. NO.:54, SEQ. ID. NO.:55, SEQ. ID. NO.:56, SEQ. ID. NO.:57, SEQ. ID. NO.:58, SEQ. ID. NO.:59, SEQ. ID. NO.:60, SEQ. ID. NO.:61, SEQ. ID. NO.:62, , SEQ. ID. NO.:63, SEQ. ID. NO.:64, SEQ. ID. NO.:65, SEQ. ID. NO.:66, SEQ. ID. NO.:67, SEQ. ID. NO.:68, SEQ. ID. NO.:69, SEQ. ID. NO.:70, SEQ. ID. NO.:71, SEQ. ID. NO.:72, SEQ. ID. NO.:73, SEQ. ID. NO.:74, SEQ. ID. NO.:75, SEQ. ID. NO.:76, SEQ. ID. NO.:77, SEQ. ID. NO.:78, SEQ. ID. NO.:79, SEQ. ID. NO.:80, SEQ. ID. NO.:81, SEQ. ID. NO.:82, SEQ ID NO: 83, SEQ ID NO. 84, SEQ ID. NO. 85, SEQ. ID. NO. 86, SEQ. ID. NO. 87, SEQ. ID. NO. 88, SEQ. ID. NO. 89, SEQ. ID. NO. 90, SEQ. ID. NO. 91, SEQ ID. NO. 92, and SEQ. ID. NO. 93; or a sequence which is the reverse complement of a target sequence located in a region extending from about nucleotide 100 through nucleotide 500 in said SEQ ID NO.

39. An isolated polynucleotide for characterizing cells which are obtained from a subject suspected of having a cancer selected from the group consisting of glioma, acute myeloid leukemia, primitive neuroectodermal tumors of childhood, breast cancer, colon cancer, head and neck cancer, testicular cancer and lung cancer; wherein said polynucleotide is from 35 to 1000 nucleotides in length and comprises at least two CpG dinucleotides, and wherein said polynucleotide comprise a sequence which is identical to or complementary to a target sequence which originates between nucleotide 1 and nucleotide 15 of SEQ. ID. NO.:1, SEQ. ID. NO.:2, SEQ. ID. NO.:3, SEQ. ID. NO.:5, SEQ. ID. NO.:6, SEQ. ID. NO.:7, SEQ. ID. NO.:8, SEQ. ID.

NO.:9, SEQ. ID. NO.:10, SEQ. ID. NO.:11, SEQ. ID. NO.: 12, SEQ. ID. NO.:13, SEQ. ID.
 NO.:14, SEQ. ID. NO.:17, SEQ. ID. NO.:18, SEQ. ID. NO.: 19, SEQ. ID. NO.:20, SEQ. ID.
 NO.:21, SEQ. ID. NO.:22, SEQ. ID. NO.:23, SEQ. ID. NO.:24, SEQ. ID. NO.:25, SEQ. ID.
 NO.:27, SEQ. ID. NO.:28, SEQ. ID. NO.:29, SEQ. ID. NO.:30, SEQ. ID. NO.:31, SEQ. ID.
 5 NO.:32, SEQ. ID. NO.:33, SEQ. ID. NO.:34, SEQ. ID. NO.:35, SEQ. ID. NO.:36, SEQ. ID.
 NO.:37, SEQ. ID. NO.:38, SEQ. ID. NO.:40, SEQ. ID. NO.:41, SEQ. ID. NO.:42, SEQ. ID.
 NO.:43, SEQ. ID. NO.:44, SEQ. ID. NO.:45, SEQ. ID. NO.:47, SEQ. ID. NO.:48, SEQ. ID.
 NO.:49, SEQ. ID. NO.:50, SEQ. ID. NO.:51, SEQ. ID. NO.:52, SEQ. ID. NO.:53, SEQ. ID.
 NO.:54, SEQ. ID. NO.:55, SEQ. ID. NO.:56, SEQ. ID. NO.:58, SEQ. ID. NO.:60, SEQ. ID.
 10 NO.:62, , SEQ. ID. NO.:63, SEQ. ID. NO.:64, SEQ. ID. NO.:65, SEQ. ID. NO.:66, SEQ. ID.
 NO.:67, SEQ. ID. NO.:69, SEQ. ID. NO.:70, SEQ. ID. NO.:72, SEQ. ID. NO.:73, , SEQ. ID.
 NO.:75, SEQ. ID. NO.:76, SEQ. ID. NO.:77, SEQ. ID. NO.:78, SEQ. ID. NO.:79, SEQ. ID.
 NO.:80, SEQ. ID. NO.:81, SEQ. ID. NO.:82, SEQ ID NO: 83, SEQ. ID. NO. 86, SEQ. ID. NO.
 87, SEQ. ID. NO. 88, SEQ. ID. NO. 89, SEQ. ID. NO. 90, SEQ. ID. NO. 91, SEQ ID. NO. 92,
 15 and SEQ. ID. NO. 93.

SEQUENCE LISTING

<110> Ohio State Research Foundtaion
Plass, Christoph

<120> Detection of Methylated CpG Rich Sequences Diagnostic for
Malignant Cells

<130> 22727/04075

<160> 90

<170> PatentIn version 3.0

<210> 1

<211> 677

<212> DNA

<213> Homo sapiens 2.B.53

<220>

<221> n

<222> (1)..(677)

<223> a or g or c or t

<400> 1
gcggccgcggg ttagcttctc ctgtccgaac gcagggtttc actggggcgc cgctacggtt 60
cctatggcaa cgcggctcct cgacgcagcc caggagtcgc ggtcgcggga ggctgcgccg 120
cgcaccgagc tcttcctgt ggccgccga gccgccagcc tcttcctgct catgcttttc 180
ctcatcttca tctcggctg agtgggctct ggacctctcc accagcctct gccccagaac 240
tgttaactgc gggggggaaa aaaggaattt gtcgtcgcaa cgcgcgttcc gatggagccg 300
cacgccacaa aggaagactc atgctgcacc ccgcggggca gatgcggcga cactggacat 360
cgctgcacag ctgggtctgc ccgtttccag agctgcttag cgccgacgcc cataaatgag 420
gaggactccc tgtgtattaa aagggggatc cgcagggttt aatttgataa ggattatagc 480
cttcataaag gcatttttaa caaaaagatg taggtggcat ggtaatcgag tattatttac 540
gcattctctc gcacacgcac tcatacctga aaacgttntg gcaggcacia aatgattttt 600
ttgtgtataa aagaatgtgt gtaactcgtg gatgggtggg ttcagcagga caagatagtg 660
acattagata aattaca 677

<210> 2

<211> 380

<212> DNA

<213> Homo sapiens 2.C.24

<220>

<221> n

<222> (1)..(380)

<223> a or g or c or t

<400> 2
 gcggccgcct tgaaggcgct ggacgggatg gtgctgaagt cggatgaagga gccccggcag 60
 gtgagctcgc ggccccgccag cccgctgccc acgcagtagt ggaagaggcc gaagtagcca 120
 ggcttggggg tgctcacgct gtcgcccacc cagtagggct ggatgaagac caccacgttg 180
 atgatggcga agcagatggg gaagatggcc cacagcacgc cgatggcccg cgagttccgc 240
 atgtantgct cgtggtagag cttggaacct cctgcgaggg cagcatgggtg cccggangcg 300
 gggccggcgg cggctgtngc tggcngnggc cgtcggcccg ggacngacgc ctggctgccg 360
 ggcgggaact ggggactcac 380

<210> 3
 <211> 566
 <212> DNA
 <213> Homo sapiens 2.C.29

<400> 3
 gcggccgcgg cgctagtgac tacttctctc tactccttct cctcctgctc cggcctcctg 60
 gcgccctgct ccaggtctct cggcgccctg ctccaggctc tccggcgccc tccagccagg 120
 caccggccga accgggtagt gccgcaaggt gtaattactg ctttgaaact ttaaaggcat 180
 ttggaaagaa actacgggtt atgcttactt tttttgtttt tgattattat tttgtaggag 240
 acacaaagtt taaaaataga aagcaaaaag tgtgacacat ttaaagagtt aaaggaaata 300
 aacgtttcca atttacctta taacatgatt ttcatacact ggatttggtt aaaacagact 360
 gactacatgg ataacttttc taggaattgt tcttaactct gatagctggc tcaactgatg 420
 taggcattaa aataacgtca tattaccatc tttcctccac gaattgatga tatttgacta 480
 tagctttgtc agggttatgt ccaactattg tataatatgt gtcagtttcc tattgctacc 540
 gtaacaaatt accccaaatt tactgg 566

<210> 4
 <211> 1297
 <212> DNA
 <213> Homo sapiens 2.C.35

<220>
 <221> n
 <222> (1)..(1297)
 <223> a or g or c or t

<400> 4
 gttcacttct cgctgcgccc cgggttctgt agaagcgcaa gaatggggct gattattccg 60
 gtgccacat gccgccccca cagccccca ccccgctccg gcgcaagact tcccttggcc 120
 aaaagaggcg tttaattagt tctggggccc cggagagcca gcgtggccga caaagcccgg 180

ctccccaggt aacccgggtt ccctgcggac ccgggagggg gcgcgcgggg ccggagcacc 240
ggccttgggc tgcgcgctcc ctccggcgac actgccgtcc ccctggcctc cggcccggtc 300
ccccgcaggc caaaggctca tctgccgggc ttgggtggcc cgggccagcg ccgcctgcgg 360
tccccgagtg cggctggctc taaggccggc gccctctccc cggctttcag tgctcagagc 420
caggccagcg ggaaagaagg cagcatggtc cgcaaaagac aggtggcagt ggcagtcttg 480
catgatactt gtccttcttc cctgttcccc attttgggga aacactggaa acacttttct 540
ctttatgcgc attcgcgtct cagcaccgag tgctccaagc cctgcgcgca gcgccgggct 600
tggaaggcgg cgaatggctg cctagccgcc gccctacta gtgacactcg gccgccagcc 660
cccgccagg atgtgcacat ctgctggcag cactggcccc ggtggcagtc accgggccac 720
ccactccaca ggtacaaccg cacccaatcc aacctggaac tcggagggct gtgcgcgccg 780
agctgggatc gcgccccaac gagccgggcc tttggctgcg ccaggggcca ggccgagtca 840
tcccccgct cgcgtcgccg cgaggcggga caccgtgtaa tacctttgcc gtgggctggg 900
cgtcggccgc gggccggaga gcgggtgtcc cacctcgct catcatttga tttccgccag 960
cgtctgagga cggcgcaccc aattcgttcc actcgctgcg ctctgtgaac cagcggcggg 1020
cagggcgggg gaggcgggc tggggnaggg cagggtggtc ccaatcccc gcccccggcc 1080
cgccggcctc gcggagcaca agtgttggga ttccacagg caggcgtgct ctgcggctgg 1140
aggcccgagc gccaggggc caggagacgt ggccgacaca gaggggtttg taggcacggt 1200
gacctcgtg ctctgtctct gaaaggcct gaaaggagcg gtttatggtg cattaccagt 1260
caagggtca ggtaccagcg cctgtgtcgg gaacccg 1297

<210> 5
<211> 651
<212> DNA
<213> Homo sapiens 2.C.54

<400> 5
gcggccgccc cgggggacgc tcagatctcg cgagaagagg gcgagcgcgc tgccccctg 60
gtgggcgggg cgaagcccgg gagagggtgg gcgccaccgg aggggaggag gggaacaggg 120
aactgaagga agtgggaggg gccggcgggg cggggaagcg gaaagggggc gtggctgagg 180
gcgggaggat taagctgcct ttttgaaagt ggagcgccag gtcccgggtt ctgggtggag 240
gtggttgctg attggtggag ctccgagcgg cggttgggag ggtcctggtc acatggtggg 300
gagtgggagg ggggaagtcc ggagagcggg agcgggatgg tagtgggctg ggccccactg 360
ggctgggaca gcaggaggat agtcttgagg aggagcgtgg ggtgctagat gtgtaactac 420
gtcccgaaact ggttcctgtg tttttctagg gcatgtggac tagggatggg tacttgagta 480

gaagcctgca acttgaagag tttgtgcagg agttagctgc agtgtcggaa attagtgtcc 540
tgtatgtctca acaaggtatt cggactgggt gtgcacacca cagctctcag gactggaagg 600
tggaatttta atctacgaag ttcccttaaa ctgcataagc ttcgggacct c 651

<210> 6
<211> 710
<212> DNA
<213> Homo sapiens 2.C.57

<220>
<221> n
<222> (1)..(710)
<223> a or g or c or t

<400> 6
gcggccgcac ggagttgaag aactaacc agctaagcca catacagacc ctcacggccg 60
cctggtctac acaggccgcc acagctacac aggtcaggc ctcagcctgg tcacaatggt 120
cacaccacac ctctcgggtc ccacagtttt gcgggagcgg tgacacacac ccgctcccaa 180
ctgaccacgc ccacacacgc tggcttcagc cgcacacgca cacagtagcc acgccccctt 240
atgctccagc cttgccagca cccgccctcg ccacgctggt cacgcccaca cacacacaca 300
cacacacaca cacacgcacg caggcctggg gcacgcccct cccccacacg caggcgtgcg 360
gcacgccttc ccatacacac acacacgcgc gcgggcctgg ggcacgcccct ccacacacat 420
gcaggggtag ggcacgcccc cacacacaca cagccgggcc tggggcacgc tcgcgcgcac 480
atgcacacac atacacacgc acaggcctgg ggcaggcccc acccccacac acgcaggcct 540
ggggcacgcc ccccccacaca tgcaggcctg gagcatgcgc aactcgcag gccttgggca 600
cacgcgcaca cactcatgca cagacacgca cgcacacatc gagccccgcc cncggaagca 660
catgagaggc acttgctttc actgactgan ggcanggctt tgggcccgcn 710

<210> 7
<211> 1204
<212> DNA
<213> Homo sapiens 2.C.58

<400> 7
gcggccgctc ctctttattc tactctcacc cgaggcccgc gcccgctccg gggagcggct 60
ctgccaggaa aacggccccga ccagtgcgcg gcgcctgggc tgcgtccgag cccaccttct 120
tccctcgtcg tcgtctccca gactaaatcc cggaaaggga aagcgggatg tttgcgcca 180
ccgcgctgta gctggtcctg acacttgcaa aatggctcagt ggctcctgct cggccaggct 240
gagtgtgtgc gtgtgtgtga gcaagggagc gaggggtgtgc ggtgtgcagg ggggtgcgctg 300

tgtgtgcgcg	cgtctccggg	aaggtctcgc	ggcgggtgga	gccgggactg	acagcccggg	360
cggagcgag	gcagctccac	acgctaaacc	tctcgctct	cccctcacc	ccacccttc	420
cactcccctc	tccttcccc	accctccccg	gccccttcca	agctctctga	ttggccaatg	480
ggacaaaagt	ttctgtggag	acggctgggc	gctgacgtca	cgggcagaat	tgtcccat	540
agggatcccc	ggggcagtgc	gcatgctgca	ggctgcaggt	tagaggcaga	aggaggtagc	600
agcggggccc	gcggcagcca	ggtggcagaa	aggagcacgc	agcatccagg	tggggggacg	660
actccagcag	ggtttccatg	gagattcctc	tgggtctagc	ctaaaaacag	cagatcagct	720
gacaccatta	gctcaggacc	taattactgc	ttattggagc	aacaaatgag	ggaaagggcc	780
agctgcaaag	gaagagtttt	tatcccccca	cccattccc	ccatctcctt	tctcccctc	840
tctccatccc	tcttgagtcc	cgggtgaatt	ctcattaact	tgcaagattc	ctgcaacaac	900
agctcccctt	ctccagaggc	caccccgact	gcttttattc	ttttatttcc	ttcttttgta	960
ttaaaaagaa	atgctaaaat	aatcagttg	ttgagtcctt	gaatttttgt	tcaatacgta	1020
ttagaccata	gagctcagag	aagacactgt	ccaatgaagt	cacaagtga	tctaatacaa	1080
gggactcagg	ggaaaaatat	cactttcaat	ttattgaggt	gaatcttttag	atatttcaca	1140
ttaaaaaaat	cttaatatct	taaatacata	aatatttgaa	acacgcaatt	ggacagaaga	1200
tatc						1204

<210> 8
 <211> 687
 <212> DNA
 <213> Homo sapiens 2.C.59

<220>
 <221> n
 <222> (1)..(687)
 <223> a or g or c or t

<400>	8	
gcggccgcac	aagcgcacac	gcacacgtcc agggcggagg aacactacta gtaacacccg 60
cctccttcta	gcctccctat	cccaaagtta tgggtccgat tttgtccgcg gcaggggctc 120
caggggcaca	ctcataaatt	cggtgcggag gaacacaact agcagcacca ccccccgcc 180
actgccagaa	ccaaagtgac	ggtgcccaga cccctccgca agcgcaaggc cgacttccat 240
aagtaattag	ccagagcacc	gtcccgttcc tgtcagcacc gagccccagc caggacaccg 300
gtattcccag	caccatacaa	gaactacttt ttogatgaag caacccaaaa gctgcgagcg 360
gttcccggtg	aggccgcccc	ctcacctggc cggcgcagac aagctccgtg cgtcaagaca 420
taacagcgta	agtgtacgac	gttgcgagc gacgcggggg ccttcgggaa atgtagtcta 480

caactggaaa cgggccggat cgtgtctgcg caggcccagc agctaagatc ggggccggcg 540
 ctccagaaca gaacgatccc tgaggctccc ttgctcgaac tgtgggactt accctactat 600
 ggtccgagcc taccctatct cattatactc aagtaacgcc ccagaaattn cagagaatct 660
 acacaaagag gttgagtctt gccgtgg 687

<210> 9
 <211> 1520
 <212> DNA
 <213> Homo sapiens 2.D.10

<400> 9
 gcggccgcga ggacagctcg gacgggggag agaaaaggagg tttccagtaa aaataataac 60
 gccagagaga aaaccgtaac tcgcgtgaca cagacagaaa tttccagtaa taatcatcag 120
 gtgatagaga aggaaggctt ccaaaatgaa gaacaagtga aataaagggtt ttagtcatga 180
 attacagcac gtgcgatgga tgagtgggta tttctcatca taaatggtaa ctcgggagat 240
 agagaaacgt gtccagccct aaactacaac agggtttggg ttgaaagaga ggtgctgtca 300
 taaagcggaa ctcaggggat ggggaagacg gcctccgtcc caaatgacaa ctcaatgaca 360
 gagaacaaaa gatccaaact aaagtgatgg agaaaaaggg tttccaacca ccacacaaat 420
 gaagagaaag actgatacaca taatgaagta ttcagtcatt aatacatgat aaacccgggtg 480
 atagagaaag aggccttagtc acaaattact cagataatgg agaaaaaagc cttattcatg 540
 tatcactcag gtagatacat caaggcaggt ttcttgccat aaaggataac acagctaaaa 600
 gagaaataaa ggttttttagta ataagtgaca attcatataa cagagaaaga aggccttctgg 660
 ccataaggat aactcatgta ataaagaaaa gtttttagtca taaataatag agaaagaaag 720
 gtttccgata gaaaatggta gagatagaaa ggttctaggt aacaaacggt aactgaagtg 780
 atagagcaag gtcacaaata ataactcagg taatagagaa agatttctag tcataaataa 840
 tacatctgct acagaaataa gggttttgat tcataaagtt atgtcataag tgataagtgg 900
 tagaaaagga aaggtttttag ttataaatta tgattcaagg gatagaaaaa caaagggtttc 960
 aagttataaa tatcatttca atgggtcaaga aaggttttca gtcataaatg aaaactgggt 1020
 gaagttttcc agtcacaggt tataactcag gcaatggaca gagaaggaaa gatttttctg 1080
 atcaatcaac tcagggtggag aaggaaagggt ttttcaataa gaaataactc agttgagtga 1140
 aagaaggctt gaggtcatga atgataatta ggtgatagag aaagaaatgt tccagtcata 1200
 aggggttaaat cagatgctag agaaagaaag gtttttagtc ataaataaaa ctgagctgct 1260
 agaaagaata gggctaccag tcataattga taactcaggt gagagaaaga ttgctgggtca 1320
 taaattgtaa cccaggtgac agaaaagaag gtgtcactca cacatgataa ttcgggttat 1380

gaggaaggtt tccagccaca gtggtaactc aggtgctagg gaaagaaggt ttgggcaata 1440
atgacaactc aggtaatata gaaaaacgat tacagtcata aatgacagag aaggaaaggc 1500
ttttattcat aaaggatatc 1520

<210> 10
<211> 575
<212> DNA
<213> Homo sapiens 2.D.14

<400> 10
gcggccgcgg ctgtggctcc tcttggccgc gcagctgaca ggtaaggcgg cggcgcgcgg 60
gctaccaag ggtctgcgct cccggggcct gagcggggag gtgataagt gctgtcctgg 120
ccctggctct ggcaggggtgc agcgtcgagc ccgcggtggc ggggcgcccg ggaggcagct 180
tggcaggcac ggtccctaag ggtggaaata aaataccccc atatcgatt accccggggg 240
accggagagc ccctggactg aggccacctc ccctcaaaag cctggacgca ggagaagggg 300
aggcagtga aaggggagcg agtgaggga ggaagagag ggtcgctgga ggtcaccagg 360
ggaaggaaac aggtccctgc ccaggggtccc cgcaggatgt gctcggagga aggttggcca 420
ggccatgggt cctgtggaca catttttatt acttccgggg aagtgtttgt agtacaatca 480
gacaaacatc gggcgttctc agttctcgga gggctagggc agggatgatcc ctctggctcc 540
cgttctccct gatgtcgctg gtgttgggtg tcatg 575

<210> 11
<211> 741
<212> DNA
<213> Homo sapiens 2.D.20

<220>
<221> n
<222> (1)..(741)
<223> a or g or c or t

<400> 11
gcggccgcgt cgctcgctgag tacaccagct gcctcatcta tctggagccc ggcctccatc 60
tcgccaggct cagcgcgccg gtccgtgtcg gtgccggagc cattggccgc gcctagcaac 120
acctcgtgta tgcagcgctc cgtagctgca ggcgcgcgca ccgcagcagc ctcttatccc 180
atgtcctacg gccagggcgg cagctacggc caaggctacc ctacgccctc ctcttcctac 240
tttggcggcg tggactgcag ctcatacctc gcgcccatgc actcacatca ccaccgcac 300
cagctgagcc ccatggcacc ctctccatg gcgggccacc atcatcacca cccacatgcg 360
caccaccgtg tgagccagtc ctacggccac caccaccacc atcaccacca ccaccaccaa 420
ggctacgggt gctctggggt tgccttcaac tctgccgact gcttggatta caaggagcct 480

ggcgccgctg ctgcttcctc cgcttgaaa ctcaacttca actccccga ctgtctggac	540
tataaggacc aagcctcatg gcggttccag gtcttgtgag cccaggaatg aaagaggaga	600
agaaacgcaa ctacctgcg cctccgtggt cccgatcctg ttgctgctgc tgcaccgccc	660
gcctttgcct cgtcttctcc aaaaactgat tntcaccccc caaaagatgt ccattgcctg	720
cactgccgcc cncatttttg t	741

<210> 12
 <211> 458
 <212> DNA
 <213> Homo sapiens 2.D.25

<220>
 <221> n
 <222> (1)..(458)
 <223> a or g or c or t

<400> 12	
gcggccgcc gtagcagagc ccagcacatt gcgggtgccc agttcatctt cgtgggggta	60
aacctgcggg aagagaggga aagggccctt agtttccatg gagatcgggt gccagggggc	120
ggaggggtca aggctggaga gcagaggac cccatctttt tgtgggatca ggggtgcccc	180
agcatcttgg agggccactg aggcctgggg gggcgcggtt taacttctag catcaggagc	240
ttaggcctgg gggaggcgct gggaagtggc aggtggggca ggagggttct gcacctgaag	300
gttgtgcacc tggattgggg gtgtagaagc ggngcaggag cgccgcggtg ggggcgtcca	360
ggcccgggcg gnggagcaag cctgggggag ggagctctgc acgcgttgct gggatgtggg	420
gggcgngggg aggcggcatg gggggagggg cgttgtgt	458

<210> 13
 <211> 615
 <212> DNA
 <213> Homo sapiens 2.D.27

<400> 13	
gcggccgccc ggcgtccgc tctggggggc cgggaccgaa gcgctcacgg cccggggacg	60
cggggttggg ccaggctgcg gcctgtggcg cgtgcaggcc tgaaggaggc gagatgccga	120
tgccgccacc gctgggtccg tggaccaggc cccttgggtc agcctcccct cccgcagccg	180
ccgtcttggg ggtgttcgca gccccgggct cccccggccc gcccgccggg gagggtggag	240
gcgatggcg cccgcctccg gctcttacgg agagcgcgcc tccccctcaa ctccggcggc	300
ggtgagccgg ggtgcgatgc gcggccgagg cctcgcccgg accgcccgtc cccatcgct	360
ccctgggcca gggagggggc gttggccgga gatggcggag gggcgtagcc gccccgctg	420

cccgccgtcc ccagccctca ggcctgggg aagccccctgc tgtggcagtg ctcgggcgct	480
atccggagga agaggagcag ttcctctttc ttggctgcgg cagggctgct tgggccggaa	540
aactaacttg tgtcggcgcc cagccgcccc ggcgggctg ccggctagct caggccgacg	600
ccgaggggag cggcg	615

<210> 14
 <211> 669
 <212> DNA
 <213> Homo sapiens 2.D.34

<400> 14	
gcggccgcgc ggggcagcgc gaggaactgt tgatttgcct gcgccttggg cccttgcgtc	60
tctcccaggc ggcggtctcc gctttctctca aaggccgtgt cgggtttgtt gtttgggtgtg	120
ggtgccggga aagggcgctt ctccccagtg aggtggggaa cttgggtgat gggaccacgg	180
aggcgccggt tcgtgcccgg tggggacggg tgaggcaggg gagagtgaga ttttattctc	240
ccccaaggaa ggagtgtccc cttctcctta ttttgagggc tattcaagct tattgaaacc	300
agaaagcggg gtttcttgtc aatctctcag ccccttcttc caaccaagaa caattgtcga	360
tgagtttcca tcacaggcgc ttgtgagaga accggtaaac ccagtacagc aaaatccaag	420
cccttggttt ccacatgcat tttgctagca gtttttggca ttgaccctcg cctcccgtg	480
tttccactcg acatcattta gcgtttgagg ttttttccc tcctcaaat tgcaaatgag	540
aaaaaaagag gaaaccagga aaagggggtg gggggtagca tttaaattgg atgtgagttt	600
ctgttgagaa ttctagcgaa gtcccctgta cactgaagcg ccgagagatt tttccgtttg	660
tgtatcttc	669

<210> 15
 <211> 998
 <212> DNA
 <213> Homo sapiens 2.D.40

<220>
 <221> n
 <222> (1)..(998)
 <223> a or g or c or t

<400> 15	
gatatccatt ataatactat ttgacctcaa agtgaatttt attgttccac acaagcaaca	60
gattacacca atttcacaac tcccagaatc caaacctaca aagacccttc ccaccaagca	120
ctttacaaaa aacgggcttc atctccatct tcctttcttt cacagttgaa aaactgcct	180
tcctaattaa gccaaccaac ttcttacctc aataaaatcc ttgtttttca gtagcatgta	240
cagtatttcc agtgatgaac agtgaactgt ctttcgtctc acacagtaac ctccgtgaag	300

aagatccacc ttgttcttta ctgtatattc ctggcatgct aactgcatcc tcagacaatt 360
ttaagtgact gaaaactcag gcaaagaaag gcaagagggc aaatagaagg gcacaggaga 420
caacgctttt caaatttttc tcaactgcgac ctacagaaac aactgtaga acacctccta 480
gtacactcac acgtgtgtgt acacctgaag tgtcaagaaa caatacccta agtgaacac 540
cctctgatat tttctatttc aagtggccgt gatctactaa actgatttcc aactcacaa 600
taggattcag tttgaaaaac actgcaataa atcaaaccctt acagttgcat tccacaagct 660
actaatgaac tcttgaaaat ccagcataca gcagagacgc tgaccaacta caagatccaa 720
accccccagg tgggcagtgt ccttctgttc agcagtggca gttccccacc accaccagcc 780
ctgagagtta attatctccc aaactcccag agtttcccaa gtagcctgag gtgtctgtca 840
tatgcccttt taacctcttt ataaattcag tcccgccgt ctcttacggg ggcaaagtcc 900
atztatcgtc ggctgtggaa agcaatacnt tctttttgtc cccttcagga acccagaatt 960
aatgaccagg ttggtgcccg gtgtgccttt atgatcta 998

<210> 16
<211> 797
<212> DNA
<213> Homo sapiens 2.D.48

<220>
<221> n
<222> (1)..(797)
<223> a or g or c or t

<400> 16
gcccctctga gttacgggga gccctgcaga caccagccc ctggggatcc tctccccgac 60
ctgcccttcc cctccgacac ttgccagtac tccccggcct ggtattcctt tcgagacccc 120
ctcacctatt ccaggtgtc ctccactgag gcgaagctct atgaagtagc ccaatttcaa 180
tataattcac gttgtgtaaa agaactttga agacggacta catcgtgcaa ggacaccgtc 240
acccgaaaac cattgggtgga acgttaaaac aaacaaaaaa caaacggca aaaccttttt 300
gaaggcaatt ttgacattta tgaatttaca gttattattc ggtttgtccc tgaaatgtca 360
cttctgaaaa tttgcatagt tttcattatc actaaaataa tctagtaaat attcccgaat 420
gaatgcattc aagaatattc actaaattat tttagtata aggaaaaagt ggaaatagct 480
gacagtcata aatttataaa taaaatgatg gttaaataaa atgatgaaca ttcataataa 540
ggaatactct atattcagac gagatctgtg tgctcacagg caaacaggtc taagcttact 600
ttaaataaaa aaggataaat tgcaaaaaga atagtttgtg taatatgatt ccacatttgt 660
aaaaatggag aaagaaatng taagcanatg tctgcaagca atcagatatg attagtgact 720

taatttcatg gatagttata taggaaatat atgtatatatt tatatgcaca tagatatgga 780
ggaatataact ttcactg 797

<210> 17
<211> 1024
<212> DNA
<213> Homo sapiens 2.D.55

<220>
<221> n
<222> (1)..(1024)
<223> a or g or c or t

<400> 17
gcggccgcgg cgctgcacgg gcgtgacgtc atggcgccgc ggagccgcgt cctccccgcc 60
ccgcccccg cgggggtcac ccacccgctg ccggggctga cagagaccct ggcccgcggt 120
ctgcagcctc ctcagtcgtg cgtgcgttca ttccgctcat agcttctgtc actcagcaag 180
cgctcaacac agacgcatga gataccctgg ctggaaggcc ctgaaaggta gtcgtccatt 240
caacacgtgc ttagcgcgct gctgatctgt gccaggcact gggccagggc cccgacacgc 300
gtcagggtag aagcaagcag aagcctggcc ctggtggagc ttacattggt aaataaccaa 360
gataatttca ggtaaattatt aggtcctatt aaaaatatgc gtcttcgcca ggcgcggtgg 420
atcacgcctg taatctcagc actttgagag gtcgagcacg ggcggtctc ctgaggtcaa 480
gagttcgaga ccaacctgng taaatggtga aaccgcatct ctacaaacat acaaaaaaaaa 540
aaattagcag tgagctgtga gcttgacca ctgcactcca gtctgggcaa caggacgaga 600
tcttctaaca acaacaaaaa aaaagtatgg gccacctagt ccagccaaaa aaacaaagtg 660
cttttttttt gctttttttt tttttttttt tttttgagat ggagtctcgc tgtgtcgccc 720
aggctggagt gcaggggcgc gatctcagct cactggaagc tccacctncc gggtttacgc 780
cattctcctg gctcagcctc ccgagtagct gggactacag gcacatgcca ccatgcctgg 840
ctaattnttt gatttttttg ttgggtgttt agtagagacg ggttcacgt gtagccagat 900
ggctaactct gactgtgatc tgcacttgcc tccagtgtgg atacagggga ccacttgag 960
caaagctcta ttctgtagg aggggtgttg tgaatcagac ccaatttgga aatcaaattc 1020
tagt 1024

<210> 18
<211> 1854
<212> DNA
<213> Homo sapiens 2.D.74

<220>

<221> n
 <222> (1)..(1854)
 <223> a or g or c or t

<400> 18
 gcggccgctg cagaccctgc tccaggcgcc gtagccttgc aggaagagca gacaaagaca 60
 ggagagagggc aaagcgccgc ttgccagag atgcagtcgg ctcaagtcaat agaggggaaat 120
 cgcctccaaa ccagggctgg gaatgagga ggaggggcca ggcggtctgg gactagaaaa 180
 agcagcaggg aattaacgtg acagtcagag ccagccagt gcctcgccgg cgctgctctc 240
 tcgcctcgcg gttgcggngt ccggaatgga gagaggaggc gggggctgag ccgttggtctg 300
 ccggagacca gctgaggtag gagtattaac tccctctgct gctctcgct gccttctctg 360
 caccctcta cacagctcta ctgcagcag gctatggccc cattctttct cctatttttc 420
 taactactga gatcagagct gaattaagct ggtgaaagga gcaaacgtg caagggattg 480
 attgccctcc ttgggggaaa agcggaggct taaaatcaat tcgacaaatg agtggttact 540
 gggtgctgag tactgtgctc cgctattgtg agggagggtt atgaataagg taccctctc 600
 ccgccccagg gtccgttgct agatctcaga atcagtttcc cctgcagttc tggaagccca 660
 aagtttcggg gttgagttgt ggtccctgat ccggtctc aaccaatcta gctttctaaa 720
 tcagaagaag gtggaattca attttcttt ctcttctctg ggatgacttt aacctgcagc 780
 cgaaatggag tctataggcc ccttaaaaaa gcgcgcgcac gccagtgtgt gtgtgtgcga 840
 gcgcgctcgc gtgcgcgcgt gtgttttaag agtaagtcaa attaatggtt ttagtgatgt 900
 tcttatttca tgattttaat tatttaccat atctgcagta gacaccagtt tggggcagag 960
 gaaccgcct ctccagactc taaaaatacc acctttttt cttaaagcttt tttccgctac 1020
 ccagtcctc tgactcgagg cagaaatctt tcccctctct ttgccctctc agaattttat 1080
 ttgccaatca cttgcggaac ttatatatt atagatttat ctcttctc acatatgagt 1140
 attccctgtg ctttttgttt gtttgttctc actgcaacat ccagcagtggt tttgtatcta 1200
 atgggtactc aaggaaagct tatccagttg aaggtcattt tctcttctg tatgagctaa 1260
 atctcagtggt ctctagaatt aaagagactc cagggatgga acttttgatt taggggtgtg 1320
 tgaagggacc cacacatata gttagactca cagcccctt actggaaagg taataaagta 1380
 ttaattcat tttggtctct agacaatcaa ccttctccca ctgaccaccc acctctgttt 1440
 cctgaattcc caaaagcaaa agaaaaccaa actgctaagc aactgcctag agcaagacat 1500
 gtatgttcag ctgccaacac ctagagcaaa ccattccaa gtggagaatg accaaaaaat 1560
 cttgattatt tcttgacctg tgtcaagtat gttgaaagcc tgccaaagtt tctcatttc 1620
 tattgaagca ctctattctt ggatgcattt tagaacagtt tgaacagtggt tacattgctc 1680

agaggtgaag aaaattgctt tgtagtttaa ggatatttaa gatttgtttg tttgtttggt 1740
 tgttttctgt cccaccttct acaaattgca cgatagatac ctcagatcag gaatgctgca 1800
 tgaaaaagta tgtccataat gcaggagatt agactaaatg actcttaaga tatc 1854

<210> 19
 <211> 674
 <212> DNA
 <213> Homo sapiens 2.E.20

<400> 19
 gcggccgcct tcccttccca ttcactggct gcctcctttg tgaactaatg actgtaatta 60
 ttacctccca gagctctttt gttatctcca accccaagcc ccggagaggg ggaatgggct 120
 ctttagtgaa atgaaagtca ttacaaagca aattaccgtc tagggagggg cagccttcag 180
 gaaagacaaa tcagatctcc atctgcatct gaagtagggg gtgtttaaat aaaaaatgta 240
 aatatcacca ttagatccaa agtactccag agctgtggga tttaatggag tttaaacggc 300
 agcacttgaa gccattgctt taccaaaaag aaaaaaaaaat cagttaaatt cagggtgtttt 360
 aatcgcgtttc ttctttgggg gttttgtgtg atttaaagc ttgcttttaa gaacctttat 420
 gttttcaacc actcatccat agtagaaaag ttctgcaacc ctagactgct ggcttgaagg 480
 aaaacctttg caggatttga tatggatttc aacaaagaac cagcctctgc gaggctggag 540
 agagctgcgg agctgccatg cctgaagtgc agatggctga accacaagtc ttaggtttc 600
 cggagttggt attgtggtga cctagagtgt cagagccagg agagcaagaa agaggagcca 660
 aactgagccc tgag 674

<210> 20
 <211> 676
 <212> DNA
 <213> Homo sapiens 2.E.24

<220>
 <221> n
 <222> (1)..(676)
 <223> a or g or c or t

<400> 20
 gcggccgcag acgcgccagg cccgccaggg cgccgcacgc cgggcgcgcc acgatgtcca 60
 cgaagcccac gatggacagc aggaaggcgg cgtcggtgtc gggcacgccc gcgtccttgg 120
 cgtagttcac cagcaggatg gcggggacga agagcccagc cgccatcagg aacttggtga 180
 cggcgtacac ggcgaaggcg cggtcggtgc aactgccaa gtccagcagg cgccggcggg 240
 gccggaccct gggggatgcc tcgcgcagct gcagccccgc accgtcagcc tccgcctcgc 300

ccggagcgtc cccggcgcggtc tgcgcggcgc tgteccctgcg cggtcgcggg cccggcccgg 360
gcggcgccct catgacagcc ccgcaggcgc agcagtgcag caggagccc cgcagcagca 420
ggaagccgcc gcgccagccg aagcgtcca gcagctgctg gccgagcggc gacagcgcg 480
acaggaacac ggngctgccc gccgncgnca gcccggtggc cagaggccgc cgcgcgtcga 540
agtacagccc cagcatgatg agcgacggct ggaagttgag ggccaggccc aggcctgcgg 600
gcgaggcggt gctgtgccgg ggtccccgga gagccccctc ttgggccccca caggagggag 660
gggccaggcc ccgga 676

<210> 21
<211> 455
<212> DNA
<213> Homo sapiens 2.E.25

<400> 21
gcggccgcgg ctggggggcg ggaggggggc gcaggacccc aagtgggggt cccggagcca 60
gaggcaagtg tcctgggggt ctggggggcg cgtgccggcc ggcccgctgc cctggcctag 120
gctggtccgg gggctagcgc gccgggggct gcggccgatg ggccggggcg gggcccgcg 180
gggtggcgag cccggggggc acgggggtcg ggggtgccc agggggcgcg gccggggcg 240
gggtggccagg gatgggggtc actgggggca aaggggatcc agtggggggg tcccgatgga 300
ggcgtgcagg gccagggcg cccgaggcgt gcgggggtcg ggtgcccag actggtggcg 360
tcagacaggc gtgggtcggt gggggcctgg gtcgcggctt gactgagggc ccggccgggg 420
ctgtggggcg tcaggagagc gtggggtgtt atggg 455

<210> 22
<211> 156
<212> DNA
<213> Homo sapiens 2.E.30

<400> 22
gcggccgcgc ttcgacgacg acgacgactc cttgcaggag gccgccgtag tggccgccgc 60
cagcctctcg gccgcagccg ccagcctctc tgtggctgct gcttcggggc gcgcggggac 120
tggtgggggc ggcgctgggg gtggctgtgt ggccgg 156

<210> 23
<211> 978
<212> DNA
<213> Homo sapiens 2.E.37

<220>
<221> n
<222> (1) .. (978)
<223> a or g or c or t

<400> 23
 gcggccgcta cagtgcgtca acaggcgctg taatccgagc gcataaacga ggggtccggg 60
 ggtggggggc cggggcggcc gtggcagtgg cccggggctg gcagcccgct ttgaaaatct 120
 ggcgaagtcg gggagcctgc gtttgctttg gcagctgcga aggcgcacag gtgcacgggg 180
 gcgggggggct ggctggcggc gccaccaccg accgtcactg acagagcctc gccatggggc 240
 cccaaattcg ttcacttgcg aattgcgtaa gcggccctcc ggtaccaaac ctctgggaat 300
 tacgcgggct tgtgcctgtg gccaccttgc taggccccac cgctccagcc tgaactccca 360
 ccgctccctg ccttgcgctt gatgttccag caacttcgaa ctgtttttat ctctgtaaa 420
 ccaagccgct tctctccttg acgtggcctt tcctgcctgg cttgccctcc cgccttcttt 480
 tgccttttaa gaccgggcag ctatcccacc ccgccagtat atgccccctc tctgggctcc 540
 ttggcttctt gtttatacct acgtgactgt gcttactttt ttgcacatgg tttttcttat 600
 ccttctgtaa gtttcttgaa ggtaggagcc atgtcttacc ctgccaagca cattgtctgg 660
 cacgtagtag ctgttcagta gaggaagtgg tccctttccc taaagggctt tncgtctcac 720
 tggagagaaa ggctagcctg gtaccagga ctgccgagat caagtgatgg cagtacgtgc 780
 gattcgatgg tgccgaaagt gacctagaga ggcagctgng agtgctctgg tgctcgcgga 840
 tagagctttg gcgatattgt catttacaat gaggactgta ctctgagacg tggaccttct 900
 aacagaccat tataaccttt gctctggagg agtgagcnag caacggactc tgacancatg 960
 ttttgacaat ggggtattg 978

<210> 24
 <211> 321
 <212> DNA
 <213> Homo sapiens 2.E.4

<400> 24
 gcggccgcac cggtcgggc tctgccaagg gaccggcct gcccgaatgc cgccggcggg 60
 cggtgcccg tgcacctgc acctgactgc gaggcgcggg aaatgaccgg gtctgtcagc 120
 ctcccatcgc ggcttcgctc tacaggtact acctgtgctc tgtccagcct cagccactgg 180
 acgatccttc ccgtagccgt aggaaggggc ggcgcttctt tggaggggat attagaggcc 240
 cgaattcgcc cgggaagcgg cgggagggcg ggggtgccgg gaaggaggga ggggagaagg 300
 agtgagggaa gtgggtgtat g 321

<210> 25
 <211> 1023
 <212> DNA
 <213> Homo sapiens 2.E.40

<220>
 <221> n
 <222> (1)..(1023)
 <223> a or g or c or t

<400> 25
 gcggccgcgg gctgggggag agcgcacacc ccgcgcgcgt ggagttcact gccgggagcc 60
 ggcattgggccc tgggggaggg gtgcacaggg cccggagggt gcgtgggtgt ggggtgcgcc 120
 cggaggagag cgaggctgcc agagtgcgtg tgccgactga gccagtgtga gtgtgcaggg 180
 gctggcggag agactgggag cgagtgtgtg tgcattctaac cgggaggttg tgagtttgtg 240
 tgcgcgcacg cccgcagaga agttgtgagc ctgtgtgtgc acctaacaca gaggttctaa 300
 gtgtgtgcac ttgtatgtgt gtgtgcacac gcggacagag tgattgtaag gatattgtgtg 360
 cacctcacag agaggttgtg agattgtaag ggtttgcgca cctaaccggag atgttgtgag 420
 tgctttttttt cctgacaggg tgtgagtttg tgtgtgtgtg attagagggt tgtatggacc 480
 tgactgaggg gttgtggaat gtgtgtgcgt gagcatgagc ctggagaggt tctatgcctg 540
 ttcactcctg acagagtttg tgagtgtgta tgattgtgtg actacaccac ccaactggcg 600
 gattgaatgt gttgtataca tctactngga gggcgtgtgt gtgtgtaaat tgtatacaat 660
 gaggtgtgtg gcatcagtg acctaaccac gaacctgtgt gtacagatgt gtgtgccttt 720
 ctgtgtatca gacatgaggg catgtgtctg nggtgtgtta gttggttgtg caagtgtgtg 780
 agtctggggg ggagagaggg agttcggagc cttcccgtt tctccttctn cactctntgc 840
 ttgtctcggc caccagcatg ttggaggact acaaggctgc ccttcaggcc ctttagaccc 900
 gcttaaggca cttgtgatcc tatatgccag atgccctccc aaagtgccag gctaccacat 960
 ggcttggtgt attgattggc attgaccacc catttgttct ttgcttcctg ggcggtcat 1020
 aaa 1023

<210> 26
 <211> 964
 <212> DNA
 <213> Homo sapiens 2.E.61

<400> 26
 agccacatgt gtacccatct tctcctctg tggaaggcgg aaggaaacag atgccctcca 60
 aatatggaca gctgaaatga tgaagtgtgt aagccctggc ccagaccctc agagagatgt 120
 actcaaccac ctccccaccc ttggacaagc aaaaaaccag agaaaacaaa ggccagcaac 180
 tgtggctcag cccgcataaa tttcttctgg aactggcct gtctatttga atatctgtaa 240
 tgtttgggtg agtcaggggt gaggtgtca gcctttggct gctgcatctc cagacaccaa 300
 tcatggggtt cttttctttt ttttaatttt tttttttttt ttggaaccgg attccaaggg 360

gccaatttaa gttaacttcg gcttccaagg ttcaaggcaa ttcttctggc ttaaccttcc 420
aaagtggctg ggaataccag gattgcacma cmatgccsgg ytaatttkgw attttwagka 480
raracarggt tttccatgt kggtwaggyt ggyctmaaac tytsgacctm aggwgatcca 540
cccgytsgg cctccmaaag tgctggratt acagsswtga sccaccgkgc csggcccatc 600
atgggtcttac taatgggtat tttcccctta acatgtcatt tgagcccctg cctgctcatc 660
agtaaactgg gctaattaat aataccctcc tgtagggctg ttgtaagaat aaaatggact 720
atgtgagaaa agggcttaac aacagggtat agtgacagag gactcggtaa ctgctttttt 780
gtgcttatta agagagaata ctacagcaac ctatgggaag atttgagtc acgaaaacct 840
gttctccgtc cttggagcca cagctggact acatttccca gccttccttg cagctgggca 900
tggtcacatg actgtgctcc agccaatgga atgtgaatgc aagtgatatc aagcttatcg 960
atac 964

<210> 27
<211> 748
<212> DNA
<213> Homo sapiens 2.E.64

<400> 27
gcggccgctc cgttgactgc agggccccgg cggctcttct ccgctgttcc gaggccgttg 60
agggctgatg tgctccatcc tcccacttgt gggttggcaa gccatccagc cgactacaaa 120
cccacgtttg tgagttacct gctggctgtg acgcttccgt caaatctgag taacagtttc 180
ctcatctcta agatgggtaa catagtatct acctcacagg atcgtgtggg cagtacatgc 240
atagaaagga tttaacacgc agtgtactca gctagtttta ttatttatcc gtaatgatca 300
tttgttcttt tcccctaact gtgcctcaca agcatgaaac agaatccacc aacatttag 360
gtctgggtag tggttggatg gaaacccatc gcgggttaac gcttccaaca ccagtccctt 420
gacactctcc cgccgaggag gctgatttgt aaacttgctg agaagagaat acccagcaga 480
tctttcaggt ttcaaatcca cgttctttac aagttgtgtt aattgtttgt atatgctttc 540
gatatagagt ctctaggaag taatactagt acatgtttta aaattcaaat actgccaaac 600
agtgagatgt aagtctccct cctaacttct gtttcccaaa tcccatgtcg tttcttctga 660
tgcaatagac attgtatgtg tgtgtgtcta gatagataca tatgtgtatc tctcggcttt 720
ttttttttct tttaaagagt aaaccaag 748

<210> 28
<211> 250
<212> DNA
<213> Homo sapiens 2.F.2

<400> 28
gcggcccgccg ggggaagggcc ctggaagagc aggaccaggc agagcgggcg ctgggggtctg 60
cgctggagct tgcgctgagg ccgggggtctg gccaggagcc gcagttgcag ccgctgctgc 120
cgcaggggtct gaggatgagg ctggagccgc agcgggaacc ggagccgcag ccggtgctgg 180
cgttggcgct ggaactgagg ctggggccgc cgccgggact ggggttggcg tggccggagg 240
agcacttact 250

<210> 29
<211> 657
<212> DNA
<213> Homo sapiens 2.F.41

<400> 29
gcggcccgct acggacagcc agtgcattag gcagggctcc cctacgcgcc cggagagcgc 60
ggaccgctgc ctggggccgg cgccgcctcc tgccgcctgc cgccgctcgc ggagcccagag 120
ccccagcccg agccgccgcc taccacaggc cggggcgctc agcagccggc ggcctgtcca 180
tgtggggcta gccctcgcgc ctggcctgca tcaggaccag caacatggag gcggccgttt 240
gcgaccccgga cacgcgagga ccagggcggt gcggagcccc gcgaggacgc gacgccccatg 300
gacgcctgtc tgcggaaact gggcttgtat tggaaactgg tcgacaagga cgggtcgtgc 360
ctgtttctgg cccgggcgga gcaggtattg cactctcagt ttcgccatgt ggaagtcaga 420
atggcctgta ttactcgtc tcgagagaac agagagaaac ttgaagcgat tatagaacga 480
ccatttgaag gaattttaaa gcgcttcgga aattcacagg aatgggtatg acaaattggaa 540
aaaagagccc tttctcttat gtacaggaaa gattttattc ctaaactgga gccaaagggt 600
ctttctcaca agtaactgaa aatattttcc tgaaaggggt tactggtggt tttaaat 657

<210> 30
<211> 318
<212> DNA
<213> Homo sapiens 2.F.50

<220>
<221> n
<222> (1)..(318)
<223> a or g or c or t

<400> 30
gcggcccgcg agcgattgca tgcaggggcc gcgtaccgng aagtgcagaa gctgatgcac 60
cacgagtggc tgggcgcggg cgcnggccac cccgtgggcc tagcgacccc ccagtggcta 120
cccacgggag gaggcggcgg cggcgattgg gccggcgggc cgcacctaga acacggcaag 180
gcaggcgng gcggcaccgg ccgagccgac gacggcgggc gcngcggagg tttccacgcg 240

cgccctggtgc accagnnggn ntgcccacgc ggtcgcagna tgggcgagg gcaatnncaa 300
aacancactt gggcccng 318

<210> 31
<211> 525
<212> DNA
<213> Homo sapiens 2.F.59

<400> 31
gcggccgcct cccgccagga aggggtggcg gcccggaagg ccagagatgc cccagtgtct 60
cccgcgccgc tacgcaccta gctgcccgcg ggtcccacat ggctgcggcc ggaggggtccg 120
caccaggacc gccgccgcct ggggaagcgc ttccctgtgg gcagggcgcg gcgggcagtg 180
cggaagcccc aaagctaccg gagccccggg cagggcgcg gcgatgcaga ggcggcgttc 240
gggggcccc agctgcctgc ggctcggcta cccagccgcg atcagagggg gcgggggacg 300
caggaacccc ggcgtccggg cgggtgtcag ccgcagacct attccaagtt tccacgtagt 360
tgcgagagcc caaaaactgt cacgtgcacg tcgctgctga gtgggaggag gtgtttgtca 420
tcgcgttcaa aagggcggtt tcgggtgtct ccgtcatgca agcaaagtgt atggctctcg 480
gccgcctttg aataaacgag tgcttcgaac cctttaccag gaggg 525

<210> 32
<211> 1032
<212> DNA
<213> Homo sapiens 2.F.70

<220>
<221> n
<222> (1)..(1032)
<223> a or g or c or t

<400> 32
gcggccgcgg cgggggggct gagaagggcc tgggtgcctg tcgcccggga gccgaggttt 60
cccggcctcc cccgaccccg ggcgccaaga gcagtcggtc ccccgccct cccgccggca 120
aaggggcccct ggggcccagg cgcgcggccc ctgctggtgg gcagggcgcc caggccagcg 180
ccggcggcta gagaaggcct ccagtcagg cctcatggaa gggcctgcct cgagcggccc 240
ctcaacgccc cgcagtggtg cactggaagg gacctaaaaa cccacctggc tttctccttt 300
ccccttcccc acgttccca gggcccaatg cccgcattct agtttcgctt tccggcaggg 360
tcaggggtga gagggaggaa ttctcaggtg tcacctctc acccgctgg agggcgaggc 420
tagaaagacg tcggggcact ctggagggga ggaagaggtg tgcctagaat tctctctctt 480
aaacgctcgc gttatcacgg aggagacttt ataaacactt taaacacaac accaaccatt 540

ttatcagcaa aagcgagggg aggggggcgt acagtaaagtg ctgagagatg ttcgagaagc	600
cccaagacgt tccctgcgga aggagaacgg aagaaagaaa ttacgggcg g aaaaagagta	660
aatattagct ccacacctaa ccaacttncnc agccccaac taggagagaa tctgctaaga	720
ttcgctttat atttatatag tctatgtgat gttaacaata ggggttgcaa atattgcatg	780
ggggcattct tagagtaaaa aattggatc tacctgaaat tcaaaaattt aactgggcat	840
cctgtatttt tattggctaa tcctgcaatt ctaactaaaa aacancttgt gaagaaatca	900
tatagaagga agctaattgc tgatgaatac agtattggga actgttatgg aactggctgg	960
aaagaaatga ttctctacga tactttgagc catgtaggtg agagagatga tgagcactgg	1020
atgtctacta tt	1032

<210> 33
 <211> 708
 <212> DNA
 <213> Homo sapiens 2.G.10

<400> 33	
gcggccgcgc ccaggcgccc cttcccctgt ggggcaaccc aagccgggga cgcgtgaacc	60
acctccgtag ccgccccgcc agcaccacca gccgtgcgcc cctgcaccac gcagctgccc	120
tgcccatgga gcccagaggg acagcaggcc cggccccag caccaccggc ctgccgggag	180
gttcgggaaa ctggcgtcgc agcggagagg gcatcaggcc aacgcctccc ccgaggctca	240
gctgcgggct cccaggcgta ggcaccacg gcccttacgc tgaccgtagc ttggacgccg	300
ctgccgcggg ggtccaatgc cggtcatgcc catcccgcgg gggttgtgct ccttccatgg	360
tccacacacc acctgcctgc atgcggctctg tgggcccgtg ggcgcctccc acctggcccc	420
caccaagtac aacagcttcg aggtgtgcat caagacgcgc tggctgtagg gcttcatcca	480
cttctgctc tacttcagct gcagcctgtc actggggcac gctggccgcc ttcttctgcc	540
tgcagtactt gggcgtagc gtctctctgt gcttccaaca caagctgtgg gtgctgctgc	600
tgctgcttgg cccgctggcg cggtgaaatt tcgctgttga acgagctgct catctacagc	660
atccacgtca acatgcttgt tgtatggggg cctgggctgg atgcctaa	708

<210> 34
 <211> 569
 <212> DNA
 <213> Homo sapiens 2.G.108

<400> 34	
gcggccgcac acgtgtccag gcgtcacgtc cgcgcgcgcc cccggggctt gcgtcagcgg	60
ctgttccaga agcgggtggg ccagggtctt gcgcaccgct ggggttcggg gccggggacg	120
ccgccgggag gagggcaccg cgcgggggtcc gacgcggagg cgtgctcgga acgccggggg	180

ctgcggagtg catcagcgcg gtccagccct ccgcctgccg ggcgccgagc gtctccgccg	240
cccggacctg ggctgggcg cgtggcggtt cctcggagct cgctgccgc ggggcgcgca	300
ccgccttgac ccgggcgcc ccgcggcagg caggcgcccg cagttccatg gttggttcgg	360
agcgcgatga gccgcccgtc ctccaccggc ccagcgcta ataaaccctg cagcaagcag	420
ccgcgccgc agccccagca cactccgtcc ccggtgcgc ccccgggcgc cgccaccatc	480
tcggctgcgg gccccggctc gtccgcggtg cccgcgcgg cggcggtgat ctggggcccc	540
ggcgggcgcg gcgggccggc ccggtgtcc	569

<210> 35
 <211> 916
 <212> DNA
 <213> Homo sapiens 3.B.30

<400> 35	
gcggccgcgc tgagctcact ccgggccctg cggaagaat tcgtaccgtt cctgttgaac	60
ttcctgaggg agcagagcag ccgcgtcctc ccgcaggggc ccccgacccc cgccaagacc	120
ccgggcgctt cggcagcctt gccagggagg ccgggaggcc cgccgcgggg tagccgcggg	180
gcgcgcagcc agcttttccc tccgaccgag gccctgagca ccgctgccga ggccccctctg	240
gcccgcgcgc ggggcaggag gcggggcccg gggccggccc gcgagcgtgg aggcgcgggc	300
ctggaggagg gggtcagcgg ggagagcctg cccggagccg ggggccggag gcttaggggc	360
tctggcagcc ctagccgcc cagcctcacg ctgtctgatc cgccaaacct cagcaacctg	420
gaggagtccc ctcccgtagg ctcggttccc cccggcccta cagggtgaga ctcagctctc	480
atgcaggaga tgggtaccac gaaggctctg gggagtcagt cattcgagct cggcgctccg	540
cagtggagcg ccaggatggg tagaaggctg ggggtgatgg tgagggtttt tgtggggttt	600
cttcgcagcg gccatgctct gcccgtggg ccgtcatttt gtcgtttcgt tttctctata	660
atgtaataac taactaggca aaaagtgtta aaattaataa ctactaaata tccgatgtca	720
ttacaacatt tataatatat aacaatatta aaacatataa ttaataataa aaaaaacctt	780
attttaatat ttttcttttt gttaatttat atcaccttat ataccatttt tctcaatacc	840
attcgataca atcataaatt tattttattgt atattgtcaa aataaaatat tcctctatat	900
aaaaataact ctcta	916

<210> 36
 <211> 998
 <212> DNA
 <213> Homo sapiens 3.B.36

<400> 36

gcggccgcag catggctttc ggccactact cggagcactg gaaggtgcag cggcgcgcag 60
cccacagcat gatgcgcaac ttcttcacgc gccagccgcg cagccgcaa gtcctcgagg 120
gccacgtgct gagcgaggcg cgcgagctgg tggcgctgct ggtgcgcggc agcgcggacg 180
gcgccttcct cgacccgagg ccgctgaccg tcgtggccgt ggccaacgtc atgagtgccg 240
tgtgttttcgg ctgccgctac agccacgacg accccgagtt ccgtgagctg ctacgccaca 300
acgaagagtt cgggcgcacg gtgggcgcgg gcagcctggt ggacgtgatg ccctggctgc 360
agtacttccc caaccgggtg cgcaccgttt tccgcgaatt cgagcagctc aaccgcaact 420
tcagcaactt catcctggac aagttcttga ggcactgcga aagccttcgg cccggggccg 480
ccccccgcga catgatggac gcctttatcc tctctgcgga aaagaaggcg gccggggact 540
cgcacgggtg tggcgcgcgg ctggatttgg agaacgtacc ggccactatc actgacatct 600
tcggcgccag ccaggacacc ctgtccaccg cgctgcagtg gctgctccct ctctttcacc 660
aggtaaagcg ctctgggagg cgtgggccag gtcttttctc ctctgaaaar ggcggagtag 720
agacagaata tgctgagttt gcaagcaggg ccccsggttt ggggtttcgc tccaggtccc 780
caccctcaa aaccaagaat cgcgtcggtg arggractca cagtgagggc tgcgacacgc 840
gcacgcgccc caccagcggtg tgcgccgaac ccctccggtc yyctatctkg yytctatcgt 900
cccctcmcyt gcttkcgagt gagaacacat ttgcaaagac ccctccaccc cccggaaaaa 960
caagagtttt taaatgcttg gagatgagcc ctgatatc 998

<210> 37
<211> 514
<212> DNA
<213> Homo sapiens 3.B.55

<400> 37
gcggccgcgg cgctgttggg ccagcagggc agcaccgagc ccgacttggg gccgcagtac 60
tgcgggggac tgcggggcgcc ccagcccgcg gggtcggcgt agtagccgag cgggcggcca 120
gtgcagcctg cagcctgcag cggcagcgcc ttcacgcccg ccgcccgcgt agagagcagc 180
gtggccgcgt tgcgcgcgaa gtccgtggcc gtgtcatagg ccgaggccgc gaagtccagc 240
cggttggttg cggcggtcac aaaccagcgt tgcggcgagg gcgcgcccgg gtcctcgggc 300
tgctgcggcg acagcagccc gttggtgtgc ggcacgctgc ggtccgtacc cggcccgggg 360
cccgcgcccg cgcccgggtg gaagcggggc ttggcgtagt tgctcacgaa ctggtcctgc 420
aggaaagagc cggccatggc gtagcggggc cggggcacga tctgcgagcg cggcgagtcg 480
ttgggcgagg gggtcaggcg gtccatgtca cagc 514

<210> 38

<211> 608
<212> DNA
<213> Homo sapiens 3.C.01

<400> 38
gcggccgcgg cgcagcggag gggctgcggg cccggaaccc aggccgggtca gcgtgtaagc 60
gccccagccg gccggggctcc gtgggggggtc agctccctga cccctacagc gcggtagcgc 120
ctctccgaga gctccgggac cagcggcccg gccgccccca aagccagcct ccctctccct 180
tccccgcacc gggatcccag accagggagg gggcgcacgt ccgacggctg aggaatagca 240
gggcgcgagc cggcccggca ggtgcccacg gtcgccctct gggaccccgg tggcgcgctc 300
tgtcctccgc gccacgctca gccaccaccc cggctgtttg ggacccggca cccagccgag 360
cgcgccgccc cctcggggac ccgctgggag gggctgagcg aggcttggag tgcgggagaa 420
gggacgtggg gcgaacccgg ggcgctgcgc cacctcggct gtctccagcg gagaccggcg 480
ccctcgcccc ccgtctccgt tcattgtgct gtattcatcc agcagatttt gaaacaattc 540
tcgtgtaaaa aggcatttta ctccgcgcgt cttccttaca gccatttagt tgggagtttg 600
cggtgggc 608

<210> 39
<211> 1025
<212> DNA
<213> Homo sapiens 3.C.16

<400> 39
gatatcctcg ctggggcgccg ggggctgcag ctgcctctgc tgctgctgct ggtagaagtt 60
ctcctcctcg tcgcagtaga aatmcgsctg caccgagtcg tagtcgaggt catagtccct 120
gttggtgaag ctaacgttga ggggcatcgt cgcgggaggc tgctggagcg gggcacacaa 180
agcgggaggc agtcttgagt taaaggggtc ttggtgcgra aacctggcgc agcgcgcagt 240
gcgcgccaca gtcccgaaac tctccccttg cagagctatc ccctaaagcg gctgggtggt 300
cttggtgggg gaataaaggg agcacccttt caccctctt ggacagtccc ctgctatctc 360
ggagacgcac ttagtgaacc agcggcttgg tgcccgccga gccccgctc ccccgggagc 420
ccggagcgca aagcccggga gtcggccccg cagcggcaga ggaatcgaat tcggccctgg 480
cgcccttaag aagccgcggg aggtggcggt gaggaataca atttgccaaa atccaaggca 540
caaagttttg cgccacctga aggagaaggc gagaggcgcc tgggcgctag cggctgcgtg 600
aaccgcgctc cgcgccgggg cccctccgct gcggctgttc ccactcgcg cctagccgct 660
ctcctacccc cgccggcacc gcagccccct ccaaccttcc ytytcaccg sccccgtccc 720
cacccccagt accgcccccg tccaacaact cttttgccag cttttcttct ttctctcgcc 780
ggctggagtg gcgagctcag ccgcgggctt taacaccctt ccataaatac arggggggtg 840

tcaaataata ataggggcac ctcccttcgc actcaatacg gagatgcaac tgcgccagag	900
accccgtgc gatacctccc ccggagccac cccaccaagg gtagcagctg ttctggaacc	960
gcccagagcc ccgctcctcg cagttcctyc gcctctcggg cgcgaggaca cccgagggcg	1020
gccgc	1025

<210> 40
 <211> 1010
 <212> DNA
 <213> Homo sapiens 3.C.17

<220>
 <221> n
 <222> (1)..(1010)
 <223> a or g or c or t

<400> 40	
gcggccgcgg accgacttcc ttcgccggcc accggaggga gggggcgccc ctaccccggg	60
agggggctgg gcgagccggg agacgggtcaa gttggggctg ggggagcgcg ggcgtccgc	120
actctggggc acgcggggac gagcccgcc gcattgtctg cgcggcctcg gaacaagcac	180
ggccggcggt ggcaccggcg ggcgcgggga ggagttgccg tcccctttcg ccgccgcgc	240
ccaccgcgtt ctttgtgtgt ctctcgccgc cctccagccg cttcgccgct cgcctgacag	300
ctgatgggct caccgcgcg ggtcccgct cctctcgcc gcagccggcg gagcccgcc	360
cggcaggagg aggaggggag aagaggagcg ttgacagatg ctgtcttga gcgggacccg	420
ccgggggaaa agtctggact gcctcggcga gaagcggccg gtaggcaacc ggccccagcc	480
tcgcattcgc ctcaaagacc ccaattggct aggagccctt ccctccgcag cggctcgcgc	540
agctccgctc ttgcgccccg cgcccggctc agcggacgga ctagcgcgcc cggtaagaa	600
tcctggggaa ccgctccgc cccctggctc cagcgccctc caatggatgt cggcgtagac	660
aggggctggt ccgcccacac aggtgtcggg aagcccagcc agtccccggg gagtgtagcc	720
aatagaaggc gacttcggca cacaccgcc ctgatccact aggacaaacc gctcgagccg	780
gggtggtgga ccgataccta ggcagatcag ccagtcgcc aaactgtgng caagtagatc	840
tgagacggtc cgtgttaatg actatatcta agagntggat gggaacgggg cgcccaattt	900
tcctngtat acgcttttgg caagttgggt tgaaaactga caacctgagc tggttaatgag	960
gcttctttaa ctgtttatgc tatacgcccta gtggctcaga caacgttttt	1010

<210> 41
 <211> 413
 <212> DNA
 <213> Homo sapiens 3.C.30

<400> 41
 gcggccgccc taaagcgcg atgcgcggcg tggccacgcc ccttcagtgc ttgtgacgca 60
 ggcgccctgg gctttttggg cgcgaaaaag aagcagtcct gggttgtacc cggcgcagct 120
 gggagcggct gcttcctccg gggtcgtatc tccgcccggc atggggctgc tggacctttg 180
 cgaggaagtg ttcggcaccg ccgaccttta ccgggtgctg ggcgtgacgac gcgaggcctc 240
 cgacggcgag gtccgacgag gctaccacaa ggtgtccctg caggtacacc cggaccgggt 300
 gggtgagggc gacaaggagg acgccacccg ccgcttccag gtatgcaggg acccgccccg 360
 aagacgaccg gctgcgcggg cctcccccta gacttttggc taccgggccc cgc 413

<210> 42
 <211> 927
 <212> DNA
 <213> Homo sapiens 3.C.35

<220>
 <221> n
 <222> (1)..(927)
 <223> a or g or c or t

<400> 42
 gcggccgccc ctccctgcct gaccgcttgc tccccgcccg cccgcccgcc gggttgtcgg 60
 cgcggggcca ctggcgggtc gtgatgagca ctcgctcgcg ccccgacag cacacgcgaa 120
 acccgccccg gcccgccgcg ccgccccgcc tctcgactc ccggagctcg cccaccggcc 180
 gcgctggctc acactctccc tcacagcacg ccggccgagg gaggaagggg gcggtccggg 240
 ctcccagggc gtggggaggg ctgtttattt tggggggagg aggggcgcga ggcaggaacg 300
 agctgactgg cggggatcct ccgaccgcc actgtggcag caccgggaag gcggggagag 360
 agaaagaggg agggaggag ggaccgggat gtagaactcc agcccgcgcg ggaggctacg 420
 gcgagggggg cgggtggcgg ccgcgggggg ggcggtgcca ggccccctcg gcaatctccg 480
 tagtctcctc gctggctgcc cgaggaggc cgggaagcga tcggggaagc tcgggaatct 540
 ccggcacggg cctgggattg tctggaggc acagcgcggc tggagtgcgg ggcancgcgg 600
 ggggggcggg gtctgtctcc tttctgggcg gggccgtatc ctgaagcagg cggggcttga 660
 gagaccgaa agccacggag tggctcctgc ttgcggtact agttggacag agtaaagtcc 720
 tggagttacc tcgcctgagc accctggttt cccgagaggg aatgggcact ctgtgagagg 780
 caagctatct gctgtcttcc cctccgcaga agaaaaaagg ctcaattgga aggtggagga 840
 tgaagccacc ctctatggtc accccaatct gagagcttta ctttatataa ctacattcta 900
 aggagtagta aaatacccg ggtggaa 927

<210> 43
 <211> 1365
 <212> DNA
 <213> Homo sapiens 3.C.64

<400> 43
 gcggccgcaa ggaccggctg agarmtgkkg gscsctgtgc tgggggcs g arggagrcgg 60
 ccytraggac tgcscscccc ccacaccggg gcccgggcgg gacacacgcc caacgggacc 120
 cctgagcccc caggctgggg accggcaggg gctccgggga ggctggtag gccaggacgg 180
 agccgccycc acgcgtagcc gtgaagcggg aggtacgcgg ccccttgagg ctgccccgac 240
 tgcagccgag ggcgcgacct gtggtgcaa ccgcctgacc ctgcttgccc gccgcgcct 300
 gcgggtctcc agcaggtccc accccacgcg cccgcggggc ccgctccaga ggctcctcca 360
 aggccgctgc agaggcgcgg ccaggctccc atttctgcgc atccctggcg ctacagacacg 420
 gcctgagccg ggtaccggc gactcccttc ggctccacc gcctcctggg gagggaccgc 480
 gcgctgctcc cacgcgggccc cgggggtctc cgcagccctg gcctgggtgc gtccgtcggg 540
 ctgctcgggt cggagcacc cccgccccgc cgcaccacca gcgcctytc ggagcgtca 600
 cccgcccccc gactcgtgtt gttgttgctt gggttttttc tctaattctc cggagttact 660
 cttttgttgc caattgtttc tatgcccga ggccacgctg taaatgagat gttacatctg 720
 caccgagcta agtaaacact ttataaatga ataaataagt gaataaataa cgaaatcgtc 780
 atctcggggc ggcccggctg ccagggtccc ggccgcgggc ctgcgggggt ctgtgtggtc 840
 ccgggccctg ccctggggtc ggggaggcgc cgggaggggc cgtttcccag ccgtgtccct 900
 accctgaccc catcttcctt cctctcccaa atcatcctc agactctggg cgtttggctc 960
 ccagatgtcg tgtgggattc gtggcttcca cccaccgctt ctcaaacaaa aacgggttgt 1020
 caccgcggct cttaacctg ggcgagccac ggagcgtttc ttcccgggat cgggatcggg 1080
 ccgcggctcg aaccggcacc tgcagaagga agaccggcc ctgtaggccg ccgccgcccc 1140
 aggaccggac tgggtggcctc tccacgtcgt gtccggaccc gacwcatcgc ctccaacgcs 1200
 aacaaacgga agcagcggag cctccgcctc cmasscykcg cyctgyscgs yswgmcmggc 1260
 gcattsragt gcwcsakkym sgcycaatym mgagagckct gracktkca aytatcwccg 1320
 actarsrrsr rcawwtkmww argsactcay tgagtaactg atatc 1365

<210> 44
 <211> 608
 <212> DNA
 <213> Homo sapiens 3.D.21

<400> 44
 gcggccgcac tgccctggcc gccacgctcc gcgcctgcgc cgcgcacctc aggggccccg 60

cgagagggcg gggaggtgac gaggtgaggt gggggcaggg agcgggctgc gcgaacgcac 120
 cgcacacgcg gcctgggagg gaccaccggc ccgcagcccc gggggaggcc cagcggccccg 180
 cgccccctgc cggaggcctt gcgccgcgc agtctccctc tgggccggga agagcccctc 240
 ccgagccccg agggcgatcc caccctctag gattactcca cgccaggcgg ccagcgaatt 300
 tatcccgccc gcctccaccg ccccttcaag ccctggggaa ctgggagaaa cgtggcgcg 360
 agcggcacct ttccacgct gtcctcaag ggaaaggacg cgagtggctt tgcccagggt 420
 aggcaaggca gatggcatct cagaccccg agtgtgccag ccgctgttg gggacagaga 480
 ggccgaggac ctctgcacgg ttttactgag gccacaccag agaaccacct agggctagga 540
 tgctgccctc agggcaagag ggtgaaacct gaagactgcg agtcgttggt gagtttcacc 600
 cgattcct 608

<210> 45
 <211> 1947
 <212> DNA
 <213> Homo sapiens 3.D.24

<220>
 <221> n
 <222> (1)..(1947)
 <223> a or g or c or t

<400> 45
 gatatcatct attttaaaag acatatgtaa aacccaaccc ttaagaaagg attcctatca 60
 ctgttcccca caggcatcct cctcagtctt acacctttcc accccccaaa acaaatcatt 120
 cagcatatct atttcatact gtaatatagg aaatagctat tttttagact ttttatatta 180
 ttagcactga tcatacaaac atggaataga aattccttat gttttatctg gatttaagggt 240
 gatacataat ggaatatatt tctatcaagc cgtacacatt agagataatg aaatcacttg 300
 tgttctagtt taaacattat gggaatttca gaactgcaac ataacaaata atcctcggat 360
 gaaaactaaa tctctcctct ggtcaggcat ctatgtgcat cagwgatgag aagacagggga 420
 ctgtggaagg gaaaacagcg agtcaggaag gactgtggcc acgtccattc cctggtcctt 480
 caagtaatta aatcctgacc tcctctaccc cagtctgtcc tggggaatgg ccaacactgg 540
 cctttcacia ctgtgtgtta ctagaaatgc aacagaaacc cagctgaatc ccctcctctg 600
 cccttctcaa aggaaagatc tgtcccagga ccatttggtc caacattttc aattatgaga 660
 actgggaaga taaagttatt ttacattta taaagaaaca catatttatt cacmctcatt 720
 wcaagraagg tcaagaatct atmcaaanac caagaggaat ttttaaaatc ccataatwcc 780
 accatcaaaa gagccacact tagcatgttg gtccacaggc ttcttttagca ccctcttyy 840

```

ttggtgatg cacaaaatgc acaatcacat tctgtctaca ttttataatt tgcctgtttg 900
ttgattamca ctatatattg aacaattttt aagacctgca acatatgttg acaacattac 960
ttccaaacaa tgtatttaca aataaatgca cacacacact atctgtctta tataaacgt 1020
gtcttacttt ctaattctcc actcttgaag atttaggttt ttccaacttt ttcttaatat 1080
attcaccagg agtcagcaac ttttttccat aaaaggccaa agagtaggcc gggcgagtg 1140
gctcacgcct gtaatcccag cattttggga ggccaaggcg ggcagatcac gaggtcagga 1200
gatccagacc atgctggcta acacggtgaa accctgtctc tactaaaaac aaaaaaatt 1260
agctgggtgt ggtgagtgtg gcggcgga cctgtagtcc cagctactcg ggaggctgag 1320
gcaggagaat ggcgtgaacc cgggaggcag aggttgagct gagccaagat cgcaccactg 1380
cactccagcc tgggcgacag agcaagactc tgtcacaaaa gmaaagaaaa aaaaaaggcc 1440
aaagagtaga tattttaaac tctgcaggcc ataggtttct gttgcaacac tcaactctgc 1500
tgttgcaggg aaagaagcca tacacaattt gtaaataaat gggcatgact gtgttcttcc 1560
cgacatgggt tgccagcccc tgatgtataa cactacagag gatgctgtta gaatgaaawt 1620
tctttacata tctctgatga tctccttagg actaattact agacatgaca tcatggtagc 1680
tgtgggtcaa agggcatgca tgctctggga tgtacattcc cagattgtct atcatgagcc 1740
tttctcatgt caaaatgttt tgtgaccacc agaaaggctg gttctgcttt tawtaccat 1800
ggawtgagga atagaaatga catggcatgg cccttccccca cagcaccacg gcttctcttc 1860
ctcagcacgg cgacaggggc ttccccttgg ccgccgcgcg ccgccaagct ccgccgcgcg 1920
cgccaagct ccgccgcgcc cgcggcc 1947

```

```

<210> 46
<211> 1637
<212> DNA
<213> Homo sapiens 3.D.35

```

```

<220>
<221> n
<222> (1)..(1637)
<223> a or g or c or t

```

```

<400> 46
gatatcttct gataaagaac caatctgcct gggagtttca aatctgaaaa agcaaatcat 60
agtttactgg agtaaactgc tgtttaaaaa taaaagagaa aggaaaaaaaa aaagaatgtt 120
tcctagttcc agaactgaca actagagcct aaataaatac ctggacaagg gtaaatatga 180
cctcaaattt ataaccgccc tgaacgcaga acatcaaccg cgacagctgt ggcacagcg 240
gcgacagtaa ttttctccct ggcattcaac cagagggcag ttggactgtg caccgactgc 300

```

actagtgggtg ggtagccaaa gctagcctcc aaagtgaacc acggtctggg gcctgggtccc	360
gtttgaccga aaatgctatc cagaacmccc wycgagactg caggcccttc ttcttgattg	420
agctagaggt gagtgaagac agggctctggg gtagggaggg gcgtccacgc cagcttgccc	480
attacctgcc ggccttggtg atgatcatct cagtgcctat ctcatgaaag cgcttccaga	540
gctcggtccc ctgcagatcc acccgcgggg cctgcggcga gggcagaggg gtcccgggccc	600
gggccagggga gncgcgccgg agacccttg ggggaagcct cccggtgacg ccagagggga	660
agctccytgc tggaagccgt cctcacagcc gcctggacag caaaggacag agaanaggra	720
actggtgagg gaaaacagag gggaagcmag ccgcggagac ggsccacact ggtggctgag	780
aagargaaaa tgaccgggag aaaaggggaa gctttggtgc catcaggtcc tcctaaagaa	840
caagccagtc gatagacacc cacattctgc ctgtcgaagg ggcgcattca gagctccagt	900
gtggcctgct tgggtcccaa gtcccaagcc cggrakaggc gygcggsmag cgtccacmcc	960
accccgctgk gcctccgcag gkcsarggcm cmasmaraaa aggcttcacg ccgnccgccc	1020
gggtctggga cgcttgcccg acggagtcag agragctccc sggcmagag tccacagtgc	1080
aaactycgac gcaacctgcg cttgaarcg caagcagcaa aagcgcccs g cactctgktc	1140
ccaagagcyt gggcctcctt aagccataag cgtytgcggc gcctcgcttt kggccttctt	1200
ttgggcccgg cgggaggmatt cttctagaar gctcttyaga acmccgcttt ygycaaactm	1260
ycggnccccc tgcgcttcca rcccarcaga agaaaagtgt gaaaagcaag cccgcgggtcg	1320
ccgtcggcct tggcagagaa atcaagagga gaagggaagg gaaccgctca actacccttc	1380
gggaaaccaa gtttccaaat atgccgccct cttcctggtt tgcacaaaacg gtttagggca	1440
ttcgttccgg tttcaggggtg gggatgccg tcgctcccct cctccccgcc ctgtgctttt	1500
aaaagttagg aaacaaaaaa gagcacccat tggctggaac cccaagggag gcagatgcag	1560
gaagcacaga gctgcaccgc taggcgcagc aaacagccgc ggccgaaggc gcgggtcgcc	1620
gagtgggcgg cggccgc	1637

<210> 47
 <211> 900
 <212> DNA
 <213> Homo sapiens 3.D.40

<220>
 <221> n
 <222> (1)..(900)
 <223> a or g or c or t

<400> 47	
gcggccgcgg cccggaccag ccgtccccc cgcgccagc tactacggcg cggcgcgacc	60

gcgggctccg gccccagccc aggcacgtgc gcccaggccg cggggaggcg ccggcgccctc 120
 ccggaacgcg ctcttggcct gcgagtgtg cccgctcagt ctccgggtgg gaagtgcgct 180
 cgccccggac cgaggggaaa gcccaacatc cccgggatgg aacagagagg cggccacccg 240
 tgagtgggcg tgaccattg gttcccttgc gcagcatctg tggagaatta ggctttcccc 300
 tcctctcttg ccagccgttg ttcctaact tgtctttttt aaggaggaa agcaggagaa 360
 ctcatgacac tttgtatcac aggaaatcaa gttggtggag agagggtttg ctgacctctc 420
 ccgtcccttc tcagggtccc taggagaatt tttgaagaag taatcggcag caaggagatg 480
 ggggcaatag agagtctcag actcgcaggg acccatgttc gtccccagcg cactactttt 540
 caaacgttta tccctcagag ctgtttcttc acctccaca caactctccc gggttcgatg 600
 acatatata tcccaccagt tcatcttggg acaggccaaa aggttaattca aaaagcgaaa 660
 cgaatctcat nttctgacct gtgccctcgg taaagtcccc angtttccac cccaagtaca 720
 cttggaagcc agggccctnc acacangctg ancaccacct tncacaaact gaaaacaaag 780
 anaatccctt ggtttcaaag ttagaatagg gatacngcgt gagtggggtg aattgcnatt 840
 gggtaagga aaaaaaaaaa gtaaatnaat taanttttnt tgacctctcg cgctgcccac 900

<210> 48
 <211> 1511
 <212> DNA
 <213> Homo sapiens 3.D.44

<400> 48
 cgggcgcggc gagccccact ttctcccggc aggaaggggg gagggccgaga gcatttctctg 60
 ttgtgcagct gagccctgcg gagacgtcat tgcattcatg ctccctcggg tgcagcgga 120
 cggggggcca aagttcaagc cgcgtccagg gcaggcagcg cgcggcggcg cggcggcgcg 180
 gggcggcgcg ccagggtctc cctctccgcg tggcgctccc ggcgctccg tccccggcgg 240
 gcccagcgct gctaccggag gccagccctg gggctccgcg gggaagagct gctcttctctc 300
 ccggaggaaa ccgagctcgc aagcccagcg ctcccagccg cagactgcag agctccagta 360
 aggtgaaagt aggcaagaag gccccctgag acgtttctaa aagcatattc tatatgtttt 420
 cattatgaaa acaccactg cactcctttt atttattagg accttaagtt atcctatctc 480
 aactaatact ttaacaatc agaatctctt aagaatcttt caatcttata cttatccact 540
 ttaatagcca acaaaacctt tagccagagt gttttaaaat ggaaattacc tgttcatggt 600
 tcttaaagat ttttaaagtc tccttctaaa tttccagcct tccatttagt ttcaagccat 660
 aaaccagatt ataacaatgt gtaattgtag agaagctgtg gcttacgggt aataacgatt 720
 aaaaataagg ccataaggta ttttatgatc attttgaaat aaaaaattga aatagtttaa 780

tttcagcttg tgcagtttga gacagatcgt caactacaaa acaaattgta gattctgttc	840
tcatggtgaa caaacattac agatgtttta ctgtgtcaac atctctaaca tttgaactaa	900
gcaatgtttc acatcagaac atgaattaaa acaatgtaaa ctatggacct ggggtgacca	960
tgatgtgtcg atgtaggttc ttggattata acaaattgtac cactctagcg caagacttcg	1020
atagtggagg aggctgtgtg tatgtgggga caggaagtac atgggaaatc tctgtacctt	1080
ccgctggatt ttgctgagaa gctaaaacta ccctaaaaat ataaactcta tttttaaaca	1140
tatgtttagg gttttatgag tatcctgata cttaaaatgt gcattgcatt gtaacctatg	1200
aattgacaag aaattaatct taagaattgg cacagaaatc atctcgatgt tttcatgaag	1260
ttcatcctcg gttctactgc ttcttgataa acaagtttca tgtttagaag gttactgaaa	1320
tttttttata tggtaaaggc acatcaaaga ctttaccatt taatatatat tagttgtcct	1380
atccagtcac gtactattta aggcaatatt aaaggtaact tagatttccc cacttacagt	1440
gatgcaaagc cttcaataa tattctgttg tcttatttcc taaacatctg aataatacaa	1500
ctttatcaca t	1511

<210> 49
 <211> 835
 <212> DNA
 <213> Homo sapiens 3.D.60

<220>
 <221> n
 <222> (1)..(835)
 <223> a or g or c or t

<400> 49	
gcggccgccc cgagaccggc gtccgcagcg gctgcgcac tcgggcctgc agcggggcgc	60
ttggcggggc ggggccgggg gagagcctgt ttgcgcagta ccccgaggagg gcggaaggcc	120
gccgaggtaa gagccgggac tcggccaggt gggagtgggc accttggggc gggcctgcag	180
ggcgggtcccc gagcgtcccc gggtaggggt ggctccctgg ggacgatgcc cagggccccg	240
gccgcgctcc ggtcgcgccc caccgaggc gcagcgcggc cttggggcgc tgctggcctc	300
gccgcggggg tgggagcggc cgcggcctgg agcagctccg ggcggggccc aggcctctggg	360
gccagggcca gctgcgcgca ggggtgagtg agcagcccc gggccctcaa gtgagccct	420
gtccgctccc caccttgcac ttctcctctc cgcagtgggc gtggcgcccc tttgtgtat	480
agggggcgcc ccaaattgaa gaaggctggg ggggagaacg cataaacagg tgtttagggg	540
gccagggcct gtgcgccaag ggttgaagaa taaagagtaa ttcttttttc ccccttttta	600
aggggggncgc gagtccccct ccccccggc cgtggtaagg gccccctt gctccgtaag	660

gggccctcct ttggnaaaac aactcctttt ttcttttttt attttgtccc ccccnccca 720
 ataatttaaa nncctccctg ntcgcccccg cccccgctt tttttttttt tttttctnaa 780
 acccccacc ccccccccc cccttnnttt gtttccgctt ttattccaag aaaat 835

<210> 50
 <211> 645
 <212> DNA
 <213> Homo sapiens 3.E.04

<400> 50
 gcggccgccc gcttgacgtg tacggcgctg atcacctacg cttgctgggg gcagctgccg 60
 ccgctgcctt gggcgctgcc aaccccgctg cgaccggtgg gcgtgctgct gtggtgggag 120
 cccttcgggg ggcgcatag cgccccgagg ccgccccctg actgccggct gcgcttcaac 180
 atcagcggct gccgcctgct caccgaccgc gcgtcctacg gagaggctca ggccgtgctt 240
 ttccaccacc ggcacctcgt gaagggggccc cccgactggc ccccgccctg gggcatccag 300
 ggcacactg ccgaggaggt ggatctgcgc gtgttgact acgaggaggc agcggcgccg 360
 gcagaagccc tggcgacctc cagccccagg cccccgggcc agcgtgggt ttggatgaac 420
 ttcgagtgc cctcgactc cccggggctg cgaagcctgc aagtaacctc ttcaactgga 480
 cgctctcta cggggcgga cggacgtct ttgtgcctta tggctacctc taccacagaa 540
 gccaccccg cgaccgcct cagcctggcc ccgactgtc caggaaacaa gggctggtgg 600
 catgggtggt gagccacttg ggacgagcgc caggcccggt tccgt 645

<210> 51
 <211> 1021
 <212> DNA
 <213> Homo sapiens 3.E.50

<220>
 <221> n
 <222> (1)..(1021)
 <223> a or g or c or t

<400> 51
 gcggccgccc gacggggaga tgcggccccg gtattgatgt cgaaaatgat ggataacgcg 60
 ggaatggcaa atatactatt tgtctaattg ctcggcaatt aaattcccct gtaaattgacc 120
 catgcctcat ttcactccta tctatggaat tttgattgaa ttcgtcagct ctaattgaaa 180
 aatactgcac tttaatgtct gcattgcagt ttcaggacga gagggtttt aatgagacag 240
 tgcccccatt acccggaat atttgagact tttattcgga atttaaagcc aggagattgc 300
 tcgactgagc cctgagattt cctctcctgt atccacgtcc atccatctcc agacgcgatt 360

taataaacgc acttaaggat aaatgcgccc cgcacctcg cgccaacgtg ttaccccacg	420
ggcgcccctc ctcggaataa gggacggcgg aggcggggga ggcgggggag ttggggggct	480
cagaagggtcc tgggtccctcc ccggcccaag tttccctgcc ctccctgcca ccctgggtccc	540
caggcactgt cgcggacccc agactccgcc ttccctaggc caaacctagg cgacctccct	600
ggactaggag gcctgggtgc ctgccaccgc cgcaccggaa gaagggactc gcgcactcgg	660
agaaggggccc gggccccgac gcgctttata tgcaaattggc gaggcgaagc catccctgag	720
aaatagctac ttgctgaagc tatntactag attgaaatga gttaagagaa acattttaagt	780
cgtgcaacga gataattggg ccgattaact ggggatgttt gctctttcaa aaaaaaaaaa	840
aaaaaaaccg ccgaggagga gagagcagta agccgcgttg attgagccca ctgtcaagac	900
cgaattccga tgcggggacgg tctcggggac tcgaagagac ccacggagga ctgagaggct	960
ttcgccggcc gcgcatttct tttcaggcat ccaccggcca gggcctagaa gtccgaaagg	1020
c	1021

<210> 52
 <211> 518
 <212> DNA
 <213> Homo sapiens 3.E.55

<400> 52	
gcggccgcag gaaccacgat gagaggcagg agctgctcct ggctgagggg cttcaaccac	60
tcgccgagga ggagcagagg gcctaggagg accccggggc tggaccaccc gccctggcag	120
ttgaatgggg cggaattgc gggggcccacc ttagaccgaa ggggaaaacc cgctctctca	180
ggcgcattgt ccagttgggg ccccgcggtt agatgccggc aggccttccg gaagaaaaag	240
agccattggg tttttagtaga ttggggccct ctttttagtga tactggattg gcgttggttg	300
tggctgttgc gcacatccct gccctcctac agcactccac cttgggacct gtttagagaa	360
gccggctctt caaagacaat ggaaactgta ccatacacat tggaaggctc cctaacacac	420
acagcgggga agctggggcc agtaccttaa tctgccataa agccattctt actcgggcga	480
cccctttaag tttagaaata attgaaagga aatgtttg	518

<210> 53
 <211> 498
 <212> DNA
 <213> Homo sapiens 3.E.57

<400> 53	
gcggccgccc ccacggctcc accctctcgg cggggccgca gccatctggg gccctgcca	60
gtagcggccg ccttccgctc agcctctggt ccagggcgag cctggcgagc cggcgaagca	120
ccggcgggga ggaggactag aacaggagga ggggcacggc ggattgaagc gagctgggct	180

gtgagcaagg gacaccacaca gcctggagaa acagccccgc tctcttgccg gctgtctgct 240
ccagccgcta ctgggggctc taagcagcgc gatgctgctt cgcttcttct aggcggcggc 300
cggcggaggc tttccgcagc cgcttggccg gcgcggccc ctattccgtt ggcaagtccc 360
ttgtctatcc cggagggcgc acccggacgc tcgagccgga gcgagcgcga agtccgaagt 420
ccgccccag agccgccaac ttccctgtga gccctctcc ccgcgcagc ctgcgccaga 480
cctgggagcg atgcgcc 498

<210> 54
<211> 471
<212> DNA
<213> Homo sapiens 3.E.59

<400> 54
gcggccgccc gggcccgcgg gcgggggat cgccggggg gacccgcggg gtgaccggcg 60
gcaggagccg ccaccatgga gttccgccag gaggagtctt ggaagctagc gggtcgtgct 120
ctcgggaagc tgcaccggtg agcctggcgg ggggtccggg agaagagtgg gaggatctga 180
ggaggatgct aattcccacc tgggcgcaga ctgacagatg aacgggagat accccggcat 240
gggggtccac ccattctgtc agttttctgc cgtgggctcc gacggcgctg ttctccctgg 300
tcgagccttg tccattatcc tgttctttt tctgcacccc accccaccg gctccactct 360
ctctggtgct gtaaagtcc ctctccggg tctctggctc ctccccacc acttctgggt 420
ctctgtcccc gtctctttt ggatgtctct gcccttttc tctctgggtc t 471

<210> 55
<211> 971
<212> DNA
<213> Homo sapiens 3.F.16

<220>
<221> n
<222> (1)..(971)
<223> a or g or c or t

<400> 55
gcggccgccc tgggcctgca aaacttccaa agtagcagcc tgtttctcct cgtctccctt 60
ctcctgggta ccagcgcgc cgcttcccc agaaagggcg aggggtgggg gcagggtctc 120
ctcgggaggt ggccaagcgc cgggacgcgc tcccagcgtt actcaggaca cttgggattt 180
ggcctgcagc ccccttcccc atccctggcc tggtgcggg gtcccttgct cccctctgct 240
gctgctcctg ccccatcaag tcgaaaatct gaggggtggga tgggggtgggg gaccaggggg 300
taccctccca ggccgctccg cagcaggccg aggtggagac cctgcccggg aggcgagtc 360

ttgtgcccac agctcggagc cagcagcggg gtgacaaaaa agataaagtt ggtgaatgat	420
aaagaccgta ttttccacgc tttgggtgcg ggaccagatg atctagaaaa tgagctgaaa	480
tggattcagc ctccgagcct gttgtgagag cagctgattc ccccatctcg ggccagatgg	540
ctgctgaaca cagatttgca ttcattttcg cttaatatcg tccaaaatag tggggcagct	600
gcatttggtg tcaaaaaggt ttaaaacccc ttttctttct ggggcaggat cgttacctta	660
tgtgatgggc ttatagaact tttttttcct ctttagtcaa cagtatcaga tttagaagga	720
tttgttttta aaccttctaa tttggtaatc agatttaaat cgcttgggcg cgtgtaatct	780
gaattaaaga tactgtaaat gattntaagc atgatacttt cgttagcgca aggaaggggc	840
acctctagca caggctggac attttaggaa gtgtgctata aaggagcatt gttcctatct	900
caacttaatc ttccgaaaag gctttggtat tctgcataac gctgctggcg ttgcctggtg	960
agcccgagag t	971

<210> 56
 <211> 550
 <212> DNA
 <213> Homo sapiens 3.F.2

<400> 56	
gcggccgcac gcgggtgcta atttgcacac atcaagactg aagtgtagtg aggaaacggt	60
gagtttctgt tttcaaacct ttaacttcgt aattagagat ttaacaactt gaaggggggc	120
ggggagaggc gggggaggag gtgggcagaa ggaataaaac tccatctaaa attcctaata	180
gcaattcctt agaattataa actgcgagat gatcagaagt gacatctttg ccttctttga	240
aggctctctt ctctaagtta ctaataatga taatgcacgt tcgggtacag aaatatgagc	300
caagaactca agtctgcaat gaaggagtgg acatgacagc gtaagaggga gcatcattgt	360
ttgatctatt ttaacctttt ccgtctcaaa gatacgatgg tgcttctctc aggaagaaaa	420
gcctgtaagc tcaaacaaga gctcccctgg aacagaagac actggagacc gtaagaggtg	480
ggaggttga agggggaaaa ggatagaaaa actgcctgtt ggggtattatg ctcaccacat	540
gggtgacggg	550

<210> 57
 <211> 870
 <212> DNA
 <213> Homo sapiens 3.F.50

<220>
 <221> n
 <222> (1)..(870)
 <223> a or g or c or t

<400> 57
 ttagactctc actgggcagg tctgctgtcc cctctgtctc cgcaggactg gagccaccga 60
 gctcgcgcct tcttctcggg gtgcgatttc tctcctcttt tggactcaag atcaatgctt 120
 cccggccggc gcagatcaca cagcaggacc ccaggggaga ctgtggcctt cttcccgctt 180
 cccaattccc caagaccgcc tctagaggct gctgtgtccg gagaactccg agcattttct 240
 ggacacagat tgcctaacag aggaacaggg gttaggtggg gagcggctgg ccggcccaaa 300
 cacagcagcc ccaagctggc tcccaagcct gggctctcca cccccgtcc cctcctctct 360
 tgagcacagt taggccaac acccctgtcc cccaaaacac ctctaccct cctcccccc 420
 cagcccccat cttcaggaac atcacagggc tcacactcac taaccgcgga gagcacatgc 480
 aggccggagc cctcagcccg gcagctctcg gaccctgccc agctcgacgc ggactcatgc 540
 agaagaggac attccgcagg taggtacaat ccagcgcctg gggcctgggg cgtccggggg 600
 gcggcctttg agcttccccg ataccgctcg cctgtctccg gagctgttcg gccgacggct 660
 gcccgntcg tgcactttca gtangggccc gctgactcta ctgcccttgg gctaggccta 720
 ccgngnatgc ccagactcct tgggacgctg gaccgcngc gcgggcggac acgcannagc 780
 tccgctctnc gcccggaatc gttgagacgg aatctcagcg gatcccgcgat cgccgagcgc 840
 cgggncaggg agaaaggccg tgtggcgctn 870

<210> 58
 <211> 848
 <212> DNA
 <213> Homo sapiens 3.F.72

<400> 58
 gcggccgcgc cgtcgcgcgc gcccggcagg actgagcgca cggagcggcg gaactcctcg 60
 ttctccacg tgtagagcag cggattgagc gcggacaggg cgagcacag gagccagctg 120
 gccgcctgca ctccccaggg caccggcagc gagaagccgc tggccaggct caccacacc 180
 agtggctgcg tggccagcag gaagacgcag cagagcagca gcaccgacag gccgctgaga 240
 cgccgctgtg ccgcgcgcgg gtgcagcgcg ggcggcaggg gctgggcctg cgccgggtgc 300
 gcggcgccac cggggcccg cgcgtgctgg gcgcccggga aggcggcggc ggcggcggcg 360
 caaccgggca actggtgcaa catgtggaag ttgatcacgc tgaccactt gacacttaca 420
 cacacgcagc gcacgatgcc caaatagcat tgcaacaaca tatctgtctg ctccaacaac 480
 accacagaag ccatcaacac cagataatgg attctcagtg gcacagcacc cagccccagt 540
 gcccaaagcg agaacaacat cactaggccc aaggccagct cccaaaacca caccaacatc 600
 cccccctagt gcacctttta tacaacaccc tgtaaataga caacccccca ataataacca 660
 attaccattt aaagcccccc aacaatttga aaaagaagga caaccgtaat tcccaacccc 720

acacaccacc ccctaaaaaa aaaataatTT tcgccaatac cgtcccaatt tttaaaaaat 780
 ttccccaaaaa cctctaattcc aaaaacccca accccgcctt cttctatatatt tcaaaaaata 840
 cccaaact 848

<210> 59
 <211> 2770
 <212> DNA
 <213> Homo sapiens 3.F.82

<220>
 <221> n
 <222> (1)..(2770)
 <223> a or g or c or t

<400> 59
 atccanatat ttttnaacct ctaacaatga agagtannac acanactcaa ttttanaagg 60
 cacaggacct atgaanacat tttatggtaa aagaaataca aatggccatt tcccacgtna 120
 agatgcatct aacctcaatg gtggtcacag naaaataaat tacaaaaaan aaagttttgt 180
 gtgaccatca gttaggtnaa ttaaattgctt cctactaatc ttttcatgat aagtannaac 240
 atactagcca ggcattgggtg ctcattgcctg tattctcagc atgttgggaa gctgaggcag 300
 aaggatacct taagctcagg agtttgaggc tacaatgagc tatgatcatg cactccagcc 360
 tgggtaacag agagtgagac cctgtttcta aataaataaa taaatgagtg catgagtgaa 420
 catacataca tacatatata cacacggttt tttacatgtt tatagagagt ataatggcc 480
 aatgaccttt taaggcacia ttagcaaata tgtattgagt ggaaagatgc atgttcttgc 540
 atgcaggatt ctacctctg aaatgcatct gataacactg cttgaaaatg tgtgtagaaa 600
 tgcccacact agcatgtttg tgggtgggcat ataaataata gcaaaacaaa acaaaggaaa 660
 aagaaaagta catatatgtg aggaaccctt ttgggttatcc tgggtttttg agataatgtt 720
 catagaagga aagcaaatca aatgaagagc aattgagcag gaaacggggg gaaataccct 780
 cagagtaata agattatctc attacactta agttttgctg atgcttcaag tttcctgagt 840
 aagttatgag aagcatcttt ctctgaaaat cttcttgctg cagaacaaac catgttttagt 900
 gtctgtatat gtctcaactt cctgtcccca cctggcggat gggaaaaagg acacggctcct 960
 tgcttggtgtt ttggagtga aagagcatta aaggcttgc agactttacc aaggattctc 1020
 ctggtctcat ttcagatcca acttccaact ccaggcagcc tctgtgtttt tctttaatgt 1080
 ataatacagga tgtacttcaa tttggactct attgctgttt ggcctgtata tgcagtttca 1140
 agatagcccc atacacctgc ctgcaatgat ccttcaggaa tagaatgggc ttctgagttg 1200
 aggaatttgg gagtatactg agccctttgt gtatttttat taagtttctc tattcatgcc 1260

aggagaaggc tgtggacaaa aagtaaagga ggagacactg gaattgtgat gtccaaagat 1320
 tccaatgttc aaggattatt tgaacccttc acgcctcttt agccaccgcc gccgacagcg 1380
 aagacgcgga gaaaaaagtt ctcgccacca aagtccttgg cactgtcaaa tgggtcaacg 1440
 tcagaaatgg atatggattt ataaatcgaa atgacaccaa agaagatcta tttatacatc 1500
 agactgccat caagaagaat aaccacaga aatatctgcg cagtgttaga gatggagaaa 1560
 ctgtagagtt tgatgtgggt taaggagaga aggggtgcaga agcagccagt gtgactggcc 1620
 ggggtggagt tcctgtggag ggcagtcgtt acgcgctgat tggcgccgtt acagacgtgg 1680
 ctactatgga aagcgccatg gccctccccg ggattacgct gggaggagga ggaagaaggg 1740
 agcggcagca gtgaaggatt tgacccccct accactgata ggcagttctc tggggcccg 1800
 aatcggtgc gccgccccca gtatcgcccc cagtacaggc agcagcggtt cccgccttac 1860
 cacgtgggac agacgtttga ccgtcgctca ccggtcttac cccatcccaa cagaatacag 1920
 gctgttgaga ttggagagct gaaggatgga gtcccagaag gagcacaact tcagggacca 1980
 tttcatcgaa atccaactta ccgccaagg taccatagca ggggacctcc tcgccacga 2040
 cctgccccag cagttggaga ggctgaagat aaagaaaatc agcaagcctc cagtgggtcca 2100
 aaccagccgc ctgttcgccg tggataccgg cgtccctaca attacggcg tcgccacgt 2160
 tctcctaacg ctcttcaca agatggcaaa gaggccacgg caggtgaagc accaactgag 2220
 aaccctgctc catccaccga gcagagcagt gctgagtaac accaggctcc ccaggcacct 2280
 tcaccatcgg cagggtgacc taaagaatta atgaccgttc agaaacaaag caaaaagcag 2340
 gccacagcct taccaacacc aaagaaacat ccaagcaata aagtggaaga cgaaccaaga 2400
 tttggacatt ggaatgtttg ctgttattct ttaagaaaca actacaaaaa gaaaatgtca 2460
 acaaattttt ccagcaaact gagaacctgg gaattcctgc acagaagaca agagagcagc 2520
 ctccccagtt tcagcaagcg ctaggtttat atttttttcc tggtttttac tgtttgggta 2580
 atagatattg aaacaagtaa tattaatacc gcatggggag aacccaacc aaagaaatct 2640
 gaaatataaa ataaatgctt ttttttccgt tttgttcat tttggatgct ggcgctaagc 2700
 ctccaagtgt catgattaaa aaaaaatta tgtccttatt tatttctagg atgaggggag 2760
 gataacattt 2770

<210> 60
 <211> 563
 <212> DNA
 <213> Homo sapiens 3.G.46

<400> 60
 gcggccgccg ccttccgcag taatggttgt tcagcgaaca agatccgggc ggaaacagta 60

gataggcggg tgcagcggg cagaacatag gttgccttag agagggtccc cggtgtccc	120
acggcggctc aagtcagagt tgctgggttt tgctcagatt ggtgtgggaa gagcctgcct	180
gtggggagcg gccactccat actgctgagg cctcaggact gctgctcagc ttgcccgtta	240
cctgaagagg cggcggagcc gggcccctga ccggtcacca tgtgggcctt ctcggaattg	300
cccatgccgc tgctgatcaa ttgatcgtc tcgctgctgg gatttgtggc cacagtcacc	360
ctcatcccgg ccttccgggg ccacttcatt gctgcgcgcc tctgtgggtca ggacctcaac	420
aaaaccagcc gacagcagat gtgagcagcg gcacacgggt ccgggcaggg ggcaagggt	480
aaggaaggag tggctagggc aggggcggga accggggtgc ttgaccacac gtgaagactc	540
agaactaacc caggcagcct gga	563

<210> 61
 <211> 4104
 <212> DNA
 <213> Homo sapiens 3.G.78

<400> 61	
gatatctctc tccaagcccc cttcccaact ccatttctgt aggaaagtac agcccctgga	60
attgggttct ggtttcgtt tgggctggag gtgggtggat gggggtcaga gagagaatga	120
ggtggggggg acttcaaggt tctgtccac cgaccagagt ctgaagacta ttcgcctttc	180
ccaacacgga cctccgccc tccaggccc ggactatccc ttcgcggtgt agcggcagcc	240
ggagacctgg ctgaggaggc aaccgcgtag acacctccct gcttagaaaa caaactctga	300
accagaccga tcccagttgg agggttcgaa aatgttccag acagcctgtc gggaggggtt	360
gttgttgctg ttggactaaa tagctattcc tgattgggtca tgtataggggt tttttaaggc	420
gggtgggggg aggagggggg agaggaaagg ctccaaacac ctgcagggtg ggggcggaaa	480
gctgtttgcg attccctgga ctggttggtc ggggacagga ggtaattccc agccattgac	540
ccccatttct ctctctccct cctcttgcg ctgcctcttt ctctccaccc ctatctttcc	600
tggaaactcg ctttgggcgc ggcagatcgc ccaggaccac accgcagcgt aactgcaggc	660
ctctcagcga aaaaggggga aagcaaagac ccgggtgtgc atcctcttcc tcggcttccg	720
cccctttccg gcggagtgga gatcctattc agagggggcg gtctctctaa atatgcccc	780
ggtgagtttt caggggaatg gtgcgggtgg aaacggtgtc taggaaggcc ttgtgttccg	840
gcctgggggtg aggaaggctc aggacagagg agagcccatt ctgagattgg ggggtggggg	900
aggggaggac cagccagagc ttggaatcgg gatctgactg ctgtagctgc ctctgtggca	960
ttcagcggct ttttcccttt tccaccaggg gtaaaaccag ctagttggac ttagtcgtcc	1020
aggcctttcc cattgggtccc ggttctgtgg acgtttccca aggcgggtaa ctttggggcg	1080

gctgtatccg ggtggtacag actgtgcctg gagctccccg aggaggaagg cggcagcctt	1140
cctggctagt gcagtcccag ctcgagtggg ccctgatccc aggcctgagg cctaggggtg	1200
ggaggcagga acaccctct tctccggtag aggcgaggat ggtggtgctg ttccctggtg	1260
ggtttggtac ttgtgcaggc ttggggcttc tccaggggtg tgtgctggtg tgggcccaga	1320
agagagacca gaggctgggt ctaagggcct gaggctgttt tcatctaaga aattctctgt	1380
atgggggatt gggctctgctt gagacctgtc cccaggaaga atctcctggg gtottctgtc	1440
ttgttctggc acaggtggaa atattctggc tgtctggcaa ctgcagatga ggatttctg	1500
ttgggggcta taagcagggt ctccgtagta caaagagaga ggagctgtag tctcaaata	1560
ctctagaacg attcagtcta aaatctccct cctccttcat tctcccaaa taaaaacaaa	1620
caaaatctct cgggcgttcc tttctgtaat ccaaatacag tgatgcagct tagtcgcaa	1680
caaccatcag tgtttgtgag tggcttcttt ggggcatgga cctctggctg gtaatcctaa	1740
accggcagga ttttcctaaa atgtggggag gagccggag aggtcctcca cagatcctgg	1800
gatccaatca tatatttctt acaaggaacc ttggcgatgg gatatttata ggtgtctgga	1860
gaggacattt gtggccaggg tcaattcatc tggaatatgt actcccattg cctctcagga	1920
atccaccgct agagcaggag cctaagaatt aattggaggg taaaaatgtg tcataacaga	1980
gcttgagctc agtctgcaac tgcagtgcac actgtcactc ggtagaagc tggggcttaa	2040
gcatggatca ctgggctcac accggtgtgt caggacggag agcagtgagg tagggaacca	2100
ataccttgaa gcttgatatgt ttcccagggg ttggtatatt tctggcacat ttcgctgctg	2160
ctgggagcaa gaggacctgg ctgatatact tctgggtgat ttccagtggc cttggtgtct	2220
tggtggttgc attctatgga tagagacctt ttgtctccac caaaatcata aactcacttc	2280
caatgaagtg tcagggacct actgccttta cagcttgat acaccaggac ttagggaatt	2340
ttgtggtttc tgtgccagac ctggggggct ggcattccca aagaagggtg acagcagtct	2400
gaatcttgac tctctgtcat cctgggtgtc tagtggcaat tgagccaagc tccagaggag	2460
gctgcagatg atccattctc ccttctgggg tgggagggat ggttcctagg atgactcctg	2520
tccagagcat tgcagtggca gtatgggagc tcaatggctg ctatgtatga tttagatgga	2580
ctctgcatgg gggtaaattg tttttttgta tttgttttct tcttttaaat acccaattat	2640
ataattcaga gagcagaaag cttattttta acaacttatg tgggtgtgat catatatgta	2700
caactcacia ctcacaaaact ctggcccttg agtctcctga tttttctgtt ttggttcttg	2760
ctggtgcca gctctatctg gatgaagcca ggtgatggaa gagccccagc acacctgtgg	2820
gaagtagagt ggctgtggct atctcggagt atgcttggtg ggtcacaagg tggtttctact	2880

gctctgggaa tacaggaggg ttgagcaaag tgagattatt gctctggtct ggctctctca 2940
cagataggct gtgagtgact tgacattcgg ccaggcagtt ttctcactgg cccattctcc 3000
ttgttaataa tgtttacttg aacgtttgca cagcactttc aaatgcataa aggaggtatt 3060
cctccccattt cccaaagaac accaaggcag gagatggcgg tgagggggggc tggaagagtt 3120
caagggcctc atgacatcct gtctgtctct tggatgggag tccagacccc actggcctca 3180
gggaaccctt caaatgccca gctccattct acctcagcca ggctctctct tgagactcga 3240
cctcacttca gagtccagct gagcagaacg aggtggactg tgcagggagg ttgggccagc 3300
accatcttct tcccttggcg acctctcatc tctgtctgag tgggagtaaa gatccgctgg 3360
gcgggcagag gactcacagt ggatttgctc agtgtagaca gacactccct cactccccag 3420
cgggggcgaa tgtgtgtgtg tgtgtgtgtg gagggagctg gttcctcggg attattctct 3480
gccagctctg gcggagtgga tcccagtcct cgtagcctcc actttctaat tccctacttc 3540
catccgcacc gggtttcttg gtgtgtgcct gtaggtgggc tgggaatatt gctgagaggc 3600
caagggaggt tcctaaagca acgaaccctt gcctgacaga ttccccgcta aaaccaaaga 3660
gcacgatccg gaatttggtc cctcctcttc cctttaggcc tgagaaaggg gacagagtaa 3720
tctctttctt gcctccttgt acatttcctt cctcctgatt tccccttctg tgtttctgtc 3780
getggetgta ttccttttct tccggtgtct ctgtcgtctt cctccatctc tgtccttttg 3840
gccctcagtc tctgtgtctc ccaggcaccc ctcccttctc ccaatccaga gaccctcttt 3900
ccctcccacc ctagcccca cgggcctccc gccctagccc cacgtggcgc taactttgtc 3960
tgctcttctt cacgtctcgg tgcgtgagtt cctctctctg cccttctccc ctttacccca 4020
gcccacgtcg gtgggtcagg ggcggtcgtc agagcgggca tccgcttgtc tgtctgtctg 4080
cccacaggat gaccgagcgg ccgc 4104

<210> 62
<211> 570
<212> DNA
<213> Homo sapiens 4.B.44

<400> 62
gcggccgcct gtctgggcgc cgcgctcctg ctctatgcg ccgcgccccg ctccctgcgc 60
ccgggtgagt gcccgcgggc cgagccgcgc accccaacc aaacctggct cctcgcgctt 120
tccaccgcgg cctgaccctt cgacagcgcg ggggacacct gttgtctcct tcctggctgg 180
ggctaggggt ggcgggcagg ggcgctggtg cggcacagaa aggctctaga cgcggggcgc 240
agcaaaggct cttgctcctc ctocggagtt acctccccac tcccagagcg gtgactgttt 300
tgagtcccac agccggtgcc tggagaccgg ggtcagttgt ggggggtaga ggacaattgg 360

ccaatccggg aaggccatct cccttacott caccoccttc ccttgcgcac ccacggccc 420
 ctggacatga gcgctgctgg gcgcatgcgc ataggagggg aagcttgggc cactcggtcc 480
 ggtcccttgg ttgtcctact gtgcagtggg tgccactccc tgctccaccc tgaatccac 540
 actgggtagg gcttgggact cctgtgcacc 570

<210> 63
 <211> 535
 <212> DNA
 <213> Homo sapiens 4.B.56

<400> 63
 gcggccgcgc tttctccatg gccccggcct cggcgcgcct ggctccggct cgggggtccg 60
 gcacggcagt ctcagtgcgc ggtcgccagg gcgcgcgtcc caccocggct cggcttgggg 120
 gtggccccgc gcctccgccc cgcacgcagc tagctggttt ttaaattgct aatctcatta 180
 acggcgcgcc cgtccgagag gcgaggctgg taaatggatg acggcgagcc ccaccccgcc 240
 cgatcgtcgc ggccgggaag gcacccgaga ttgcagagga cagggcggag tccctgggg 300
 tctccgggt cggcggggcc tttcttcagg ctgcggaact cctcgaagtg ggcgccttc 360
 ctgggccact cacctgtcat ttatcgagcg cctactgtgt gccaggcatt gtctggggac 420
 acggctgtga accacttccc agctccgtct tggagctgac attctggtag agggaaacac 480
 ttgaattgga ctgcatgaaa tgccccattt tcaaccattt ttaatttat agaaa 535

<210> 64
 <211> 737
 <212> DNA
 <213> Homo sapiens 4.C.05

<400> 64
 gcggccgccc ggccgggtta aggcctctca gccaaaggcc cggccagctc actgccaggt 60
 cgggtcagcg cctgcgcgcc aggtccggcc ttggataccc tctgccgcca cgcgtcggtc 120
 cggcctctac gcccgcctgg ccctctgcgc gcgccgccga cgcgcaggt ccgggcctcg 180
 gtgactgccg gaggggcgcg gcgccccgcc tctgtcacc atggccaccg caaccccttc 240
 caccgcctca cggccggccg gcattccaatc acaggcgagc gttaccgatg ccggggcggg 300
 gcaagacagg gagaggaagt cccggaaggg agtgccgagg gatgcggcgc ttcggcgagc 360
 acccgttgtg tgggaactcc gtctcaagtc gccccattg tacggatgaa ggaatcgaag 420
 ccacgagcca gaatttcctc actcgcaact cgagaataaa ttgcgcctcc ctgagtgtgg 480
 aggattaaat aagtagttta aggcgtgttt aaagagcgtc tgtaagttgc caagtcgctg 540
 gagagccagt cccttatccc ttgaaccagg tgatgctgac gtctgatttc aagacagttc 600
 ctacccctcg tggaaggaaa gcccacgcgc aagaagtcga tgtcctgtaa ttacgttat 660

aatcttcgca tcataaagat tactcggcag taattggttt cttgactaat tataaccagat 720
gagaattgaa gactatt 737

<210> 65
<211> 684
<212> DNA
<213> Homo sapiens 4.C.25

<400> 65
gcggccgcca taggaaacac ctggcagtta gttcctcaaa aggttaagcc cagaactccc 60
gtaagaaccc gcaattccac tccttagtat agacccgaga gaaaacatgc gtccgtccac 120
gcaaaaatct gcacacgaat gttcacagaa gcacagga taacagtcga aatgtagaga 180
caacccaaat gtccatatgg atgaactaac tgtggtccat ccatgaccgt aatggaacac 240
gaccataacc aggtgtgaag ttcagctgtg acagggatga ccctcgaaca cggcagcgtt 300
ggtaaaacaa gcccgatgca gaacagcacg attctattta tgcgcctgcc cacaagaggc 360
acaccccggg aaagaaagca gatcagcact tcccaggaac cgggacgcag ggacgcaggg 420
agggaggggac tgctgaagat gcacggcggt tcttttggga tgaagaacag gttctaaaat 480
cgactgtggt gatggctgcg taaatcagtg aatacactaa aaaccttact gaactgtata 540
ttattttattt atttattgaa acagagtctc gctttctcgc ccaggctgga gggcaatcgc 600
accatctcgg ctactgcaa ccttcgcctc ccgggttcaa gggattctcc tgcctctgcc 660
tcccagtag ctgggactac aagc 684

<210> 66
<211> 1012
<212> DNA
<213> Homo sapiens 4.C.42

<220>
<221> n
<222> (1)..(1012)
<223> a or g or c or t

<400> 66
gcggccgagg cggcagcggc tgcggggagc tccagcagcg gcggcgaggc cggcggcggc 60
agcggcagcg gcagcagcag cagcgacacg tccagcaccg gcgaggagga aaggatgcgg 120
cgcctcttcc agacgtgcga cggcgacggg gacggataca tcagcaggta cgcggggagg 180
tacgaggaaa ccgacaggag cgagatcagt ccctccgcgc gcccttgacc cctgctctgc 240
cccctcgccc caacttgagg caagttgctc agaagctcgc gggaaaagtt ggccgcgact 300
ccgagagcgc gtagccggct cggccacgaa ggccgagggg actgctctgt tcgccttgcg 360

ggggtgccag ttggtccaac ttttcccagc gctgtctttg tctagggcgtt gggagacatc 420
 tccttaggat gcgcactctt ccgggggctc ggagtgttct tccctgtggg aaaaggagtt 480
 ctggccgctt gtcccaggta ggaggggctg ccccacagcc tcggggtcct gggcatcaag 540
 atgccgcagc acggggcagc gatctgcccg gcggcttggg ggacacccca gggccgcacc 600
 gggaggagat gagctaagcg acagcctcgg acagggaaat aacctgtgaa gaaactttct 660
 tgtgccgcag aacccatgaa ttccaaactt cagagcccaa gaatgggtat cgtttgccac 720
 ccagtattga tttaaacgca gtagcctgag aggaacgaag cgctcaggag caaactaggg 780
 ctagaccgga ctntactccg gctctgtgcg ctgaccaggt gagcttcggc gtgggtccgg 840
 gcgcctcgng cctcactaca acaacttttg ggtgttgctt cgatccccga cttctacaga 900
 gcngattaag cttctgctcc ngctgncaat atactctgcc aattggacta acttgngtga 960
 gaagatccac ttctgatgct ttgatgtgca cgctgaatgg ttcngatga tg 1012

<210> 67
 <211> 595
 <212> DNA
 <213> Homo sapiens 4.C.9

<400> 67
 gcggccgcct tgaaggcgct ggacgggatg gtgctgaagt cggatgaagga gccccggcag 60
 gtgagctcgc gggccgccag cccgctgccc acgcagtagt ggaagaggcc gaagtagcca 120
 ggcttggggg tgctcacgct gtcgcccacc cagtagggct ggatgaagac caccacgttg 180
 atgatggcga agcagatggg gaagatggcc cacagcacgc cgatggcccg cgagttccgc 240
 atgtagtgct cgtggtagag cttggaggcc tcctgcgagg gcagcatggg gcccgagggc 300
 ggggccggcg gcggcgggcg ctggcggggg ccgccggccc gggacggagc gccgggctgc 360
 cgggcgggag ctggggacgc acgcgagaag cggccctgag tcaaggaaacc cgcgagggcg 420
 gggcctgggg cagagctggg ggcgtctggg agctgctaag ggagagagga aggggtcatg 480
 agagtgttga ggccgtgtct agggggactg gcaaaggctc cctactgggg ggcctaggaa 540
 ggggccatga gaaagttggg gggcgccctag gatggggata tgagacctga agtgc 595

<210> 68
 <211> 1955
 <212> DNA
 <213> Homo sapiens 4.D.07

<220>
 <221> n
 <222> (1)..(1955)
 <223> a or g or c or t

<400> 68

atatctatcc	atatctatac	ctacatctac	ctgtatgtgt	gtagtgtata	tatatacata	60
ttatatgtgt	gtatatatgt	acatatatac	atttaaacia	aaatttctcc	ttcgtcctcg	120
aagcaaacia	accagcacc	tcgagtgtcc	gccaggaggc	gcagggggca	gcgtgggacc	180
tgcggtacct	ccacggttgt	agaggtgtag	agggatgccg	cagcgacgga	accgggcttc	240
ttttttaaag	aatcaatgtg	aggggaagggt	gcagagccgc	gttatttcag	ggagacattg	300
tcgcactccc	cctcccacgt	gtaggtagca	tctgggggtgc	gtgcgccttg	ttcgcagacc	360
ccatggagag	acgctggcgg	cggcagatgg	ggctcctttc	acggttgacg	ccggcagtaa	420
cccgaccccc	ccggcgacga	gactgaagaa	gcgcakggga	cagcggcgag	ctgcgaacia	480
aagcccttgg	cgcgggggcg	aagcccakga	cgcggtgtga	gtaaaccggc	tcgggtaccg	540
ggagctgcgg	gaacctgggc	ggccagggtc	tttgcactcc	aggagcccac	ccactgggat	600
gctgtggggg	aactntcgga	gggcacccga	rggcgggtat	ctgaaccccg	actgggggtg	660
atggtatctt	tagcacattc	agacttgag	gagawycgk	gcggtctgag	artatccagg	720
caccttctcc	atccccagca	aaacamccgg	tgggggtggw	ggtggggggcg	gaggcggcgt	780
gcagagccct	cagtaagccc	tgccagagct	gctggagcaa	gaatccatca	cccctcccgg	840
agaggccttt	ggggacttct	cccagccctt	taatcaccgg	ggggccttgc	gaccgagtct	900
cctttggcag	gggaaatcaa	ccataaactt	cttyccytag	gcaaattggg	tcctttggga	960
tgaacaggcc	tcttgctttt	ttgttctctg	aaagctgcat	ccccagtagc	ccgcctaagc	1020
tacaaaciaa	tacgctaata	ctcccgggaa	tcctccagcg	cctccctctc	tagctcctgc	1080
ctgcacctgg	atcttttcat	cttaacttgc	agcagaaagg	ggatgcatct	agcgggctag	1140
gcgcccagag	gagcctcgcc	acaggcctcc	accccgcat	ccgggggctg	aggagagacc	1200
aggctgctct	ctgaacacga	gtgtccgccc	caccccmatc	ccsgtyytgg	cgctcagcct	1260
gggctttccg	acatcggttt	tatgattttac	gtyccaccaa	agcctctgag	cctaataccga	1320
aagcggatta	agttgggatg	gggtgactat	ggatgaggag	gggggaagag	ctctcagacg	1380
tattcctcga	tgtccctcct	tgtgatctgc	agagattcca	acaaaggacg	gggctcagcc	1440
atggtggacc	cagtgcctga	agaagagaag	gcaggagcgg	aaccggcgga	ctctggaggg	1500
gacgaggccg	tggcgctcgt	gccccctgat	tcccagggcg	cacaggagcc	cgcagcctcc	1560
tcggcctcgg	cctcggcctc	cgcgggcggtg	ccccgcaagg	cagaagtccc	atgtgcagcc	1620
gcagaaggcg	ggcgggcgga	gcagtccccg	ctgctgcacc	tcgacctctt	caacttcgac	1680
tgcccagagg	cggaggggcag	ccgctacgtg	ctgaccagcc	cccgctcgct	agaggcctgc	1740
gcccgcctgtg	cgggtcaagcc	ggtggagctg	ctgccacggg	ccctggccga	cctggtgcga	1800

gaggctccgg gccgctccat gcgggtggcc accggcctgt atgaggccta cgaggcggag 1860
 cggcgcgcca agctgcagca atgccggggc gagcgcgacc gcatcatgcg cgaggagaag 1920
 cggcgtcttt tcacgccttt gagccccgcg gccgc 1955

<210> 69
 <211> 1888
 <212> DNA
 <213> Homo sapiens 4.D.08

<400> 69
 gcggccgcca gctcacaaag gatagggagg gatattgctc ttggcatttg atgggaagca 60
 tctgctgcat cccattgggg tgttgcccag gatggatttg aaaagagttg gcaggaaggc 120
 tgagctctgt gctcacaacc tggcttggtg gtggccgagg agcttggcag gagcagagtg 180
 caggacctgg gaactggggg ttggtgcatg tgtgcacgca cgtgtgtgtg tgtgtgcgtg 240
 cgtgctgggt gggtagggag gaagctgtga aaccacatcc cctcctctct gctgctgtgt 300
 tgctgtgtgt ttcagcagca cgtgggtgtc accacacttc ctagcagggt tcaacctcca 360
 agactgttct gggctcttct cccagttggc tgagttggag gtgggagtcc caactgtccc 420
 ctgtggcttc cagagtggga ccttgctgtg ggataggctg gccaatggtg ctccctcccc 480
 tgtgaccctt ctgttgggtg ggtcacgagg aaggactgtg ggtgttggcc acagacaggt 540
 ggacatgtgg caaggacacc ttgggacctt ctttctgacg ccccttgaag ggggcacttt 600
 ctcagctttg agatgagtct ctgtggatgt ggggaagttca ctatctcaag agcagcagcc 660
 ttggaaaatc caacacagaa ccccgagtag gggcggaag gggtcctgtc ccgctcactg 720
 gctgcctggc agagttctgc acaaggaagc gcctgtgttg ctgtgggcgg aggaatggac 780
 tgagggctac attcgttcc tgttgccgt gtaactgctt atcacaaact cagtggctta 840
 aagcaacaga ggctccttcc tttacagtgc taagggtcag aagccgatca gtctcaccgg 900
 actaaagtca aggtgttggc agaatccatt cctgcctctt ccagcttttg gtgggaggct 960
 ctgctggagt tccttggctt gcggctgcat cctccagcc tctgcctcca tcctcctaca 1020
 gcctcctcct tctctgcagt cagatctccc tctgccttcc tctttttttt ttttgagacg 1080
 gagtcaccca ggctggagtg cagtggcaca atcttggctc actgcagcct ccgcctcctg 1140
 ggttcaagcg attctcctgc ctcagcttcc cgagtagctg ggattacagg catgtgctac 1200
 tacacctggc taatttttgt attttttagta gagacagggg tttgcatgt tggccaggct 1260
 ggtcttgaac tcctgacctc aggtgatctg cctgcctcag cctcccaaag tgctgggatt 1320
 gcagccatga gccatcacac ctggcctgcc tccctcttaa aggacgcttg tgatttgggg 1380
 cccacctggg taatctcttc atctcaacat cttcagttac atctacagag tcctgtttgc 1440

cacatgaggt aacacagttt gggaagggag agttattcag cctaccctag gggcctgtgg 1500
 tgtatctcag ggcccttctg attttaagat ataaagcaag aaaacaaact ggctcaaggg 1560
 gaaaaaagga cacgttgaat tctgttgctt taaatgtata tttttttatt gtgctaaaat 1620
 gcacagaaca taaaatttgc cattagtaac actgagtaca ttcacagtgt cgtgcaacca 1680
 tcagcactgt ctacgcccag aactttttca tcaccccaaa gggaaacccc gtatccatga 1740
 aggactcact ccccatctgc cctctccagc ccttggcagc caccagaatg ctttctgtct 1800
 ccataaattc atttttaata agtgcaattc tgtgtgactt taaaataaat aaacatgagc 1860
 acgatgagtt gcttattgga aggatatc 1888

<210> 70
 <211> 994
 <212> DNA
 <213> Homo sapiens 4.D.12

<220>
 <221> n
 <222> (1)..(994)
 <223> a or g or c or t

<400> 70
 gcggccgcta ggaaaaggct cagctccggc cgctccgatt agccgtggcc ttgctctgcg 60
 agcagataaa cgtgacctcc gtggcctgtg gccagcctcg gccctctgga ggcggggctg 120
 tgtgcgggccc tccccctccc agcagggctg agctcagaag cagcaggcag ccggaagggc 180
 tgggcagtc cgcacctgt cctgtgccca gtctggtggg tgttgtgtgt gcaggggtggg 240
 cgtgccggga ccctctggcg tggggctgtc tggcaaaggg cgagggggga gggggctgtg 300
 cttcagcata gaaggggaag gcgtgtccag aagaggggaac agaagagggt ccagaggccg 360
 aaccagaaca cgtcccttca ctgatggaaa cttcccaccg cgctcgaatc aattcccaat 420
 tgctcgactc ctgcacctc ccgggaggtc ctgtagaggc agcgctccct ccagacctca 480
 cccgccggcc tgttcctgcc acagggtctt gcccttcttg agctctccgc ccggactctc 540
 atccccgact ctctcccca tctccttcca aagccagttc tttctcatta ctcagggtctc 600
 tgctccaatg ccacctctc ggagggggcca cctcatctc tgaacggcgc ccacccctcc 660
 ctcttttctc ggngccagct ccattntccc cttctccttt ntcaccacgc ccacaactta 720
 gaggcgcgtg tcccgctccct agaactgctg cggncacagg actnctggcc cttngcatag 780
 gctggcacgt ggcacgttcg cccagcctc gtacgcattt tgatggagag ttggaccaga 840
 gagggcgcgg agcatgaatc tctgaagagc tgaggagccc aaatcagaag ctggtgagtg 900
 agtttaatct gacttggagc atggagttat acgggagctg cttccagaag ccagctctg 960

cactgctacc atatatggca cggacgcttt agct

994

<210> 71
<211> 677
<212> DNA
<213> Homo sapiens 4.D.13

<220>
<221> n
<222> (1)..(677)
<223> a or g or c or t

<400> 71
gatatctttg ttgcattgag acaggaaagc tattttaaga tgggtgtggtg aaaaaggata 60
aaagctcctt actcaagctc tagcttatct aactctcagt caataggtaa caaacacccc 120
aagaagctgt taactgcaag ctctatttc agagggttag ggacttcccc agatccccgc 180
ctgtacagtt agacttaaac tccaacctac atttaccctt tcctcacttt aatgctaaaa 240
attactcctg ggggtggagat ttaaaatgct aatgctacat atgatgtatg aaaaagcata 300
ttggggccact gtgcaagcac tagaaaaact cctcctatag gtgccctgat gntaacccctc 360
ccctatagaa agaccctata aaactgacct acacactatc ctcagagcag tccgttcctt 420
tgcctttctt ggtgctgact cccttgcgca caagctgaat acactttcct ttgctgctat 480
gtttggtgat ctctgttaat ctctatcatg ggagatcata agaatccagg gcaacagtaa 540
cagcttctga gtttttaaatt taaaaataac agtaataata tccttaaatt tttaaaatgt 600
aggacactaa acaagtaaaa tctaaatcca gactacatct gacctcaaag ttcattgggct 660
tctcacttcc ctggcca 677

<210> 72
<211> 435
<212> DNA
<213> Homo sapiens 4.D.47

<220>
<221> n
<222> (1)..(435)
<223> a or g or c or t

<400> 72
gcggccgcgt nccctctcgc ccgnaaagag gactggagaa ggggctgggg tggaggtntt 60
ctctgtgtgt gtctanggtt gngggcagga gaggttaatt ctattaagan ntcacatc 120
anccngtgtg cacttttcgc tcgacancgg ntctnctac ttnanagcaa gtctggncca 180
gctgggatcc gaccagaaac cgcaagcgna ggagacgcat gancgnaggc tgagcgctaa 240
ctgaaggcnc gacctgagcc ctgcagcctg ctggggagct gcgcaaccac ggacagcagt 300

tcggcaatac acggcctggn ctgcatggcc cccgtcacca cctcacgtgg gaagccagca	360
ctgctgccgc cagccctgcc gctgccctca gactnncaag gcgnccaggg tcttcccaac	420
gcgcctgccc cacac	435

<210> 73
 <211> 2343
 <212> DNA
 <213> Homo sapiens 4.E.53

<400> 73	
tggccagggtg aggtcaggct ctgtttcttc cgagctacca tctctacct gattcctcac	60
accttttttct tgtagggcg agctaagaga cagagagaga gagagagaga gagagagaga	120
gagagaagcg actgaaacag agagtaaatt ctagtcttc ctttttagtc tctttcttc	180
tgccctttgc tctgctagtt tatctgcgtc tttctcttc tcgcgtgca agagtggaaa	240
actcgtgctc agttctaggc aaacattaac cccgggcgac gtttccaagc gggagacaaa	300
ctctagagag tgagaagcga gatgcgaggg caccaagggc aagaaggggg ctcggggtac	360
gccacgttg cgggacgcg ccgccgcctc cctctgctgc gcggcctgcg ccgggagcct	420
gggtgggggcg gcaagacgac agaccccgcg cccgggcctc ccaccagtga ccacctccct	480
cgcagcttg gctgatectc cagacagcat gcaacggtgg ggagggaggt cccctgactg	540
ggcgggggac ctacggctg ctctgaaact ccgaacacct gaagaggagg cgcggaaggt	600
ccagccgccc aagactcgca ctttccctc ctccgcagcc cgggcagggt accgtcctgg	660
gcctgggtga gcgcggagg gatccgggcg ggagctgagc tcggttccc aggcctgaca	720
agtggccgcg tggcacgacc aacccgggc acagggtg ggctgctccc caagggtggg	780
aatttaattc tcacattttc gactaccct gacggagctg gacgcgggaa gcgggaaaga	840
cccgttctg tttgcagtgc ccgaggggca ggacacctac cagaagggct ctatcacagt	900
gggtgttaggc cgggcgcagt ggctcacacc tgtaatcca gcactttagg aggccgaggc	960
gggaggatcg cttgaacca ggaggcagag gttgcagtga gccaagatcg cccactgca	1020
ctccatcccg ggcgacagag ctgtcttgaa aaaacacaca aaaaacaaaa aacagtgggtg	1080
ttagagggat gggattatag gtgacatgac tttcgttttg aactttcctt aaccttgag	1140
gggcagccgt gccctgaaaa cgctgtgat ttggagtaga gggccaggc gcagtgtggt	1200
gagtgacct aggcagggtca ctagtcttt ttcagccttc actgaatcct ctcttacacg	1260
gggatgttac cccaggtct ccgtgtcttt caggagaaa ttagttcatg agttagatgg	1320
tgcactatca atcatccttt tattagacag aaacaataag tttgaggaag aggacgtcta	1380
ccttacaggg ggtttaattt tcagcttctt tgagataaaa ttcattgaac ggtgttttac	1440

gtgcgcgcct tttccaacag accccacgcc tattcccagc gccagaggcg gacaaccgct	1500
ttactgagat acagagacag gtacttcctg aggcacttca gtccagttcc actgggttta	1560
ctacaactaa taatgactgt ttctgtttac taggtattag gcgatgtgtt ttaagtaaat	1620
gaattgtctc taatcctcac aactctaaag caagttaggc gtcacccgca ttttacaat	1680
catagcgccc tgctcaccat atctggaatc ttgcctcgcc ccgagggttc taattttcac	1740
tttagagagc tgagcaagat gattgcccag cgctaactcc gtgaaatccc tgggactgaa	1800
aatcacaggt aactcgccag agtttttcaa ttttaggcct aggagattat gcaaagattt	1860
ccttcaagta aacgctgttc tctggggcct ctgggatcta cagtcggaga aggggaataa	1920
gtcccgggcc ggtgggggat ggggtgggtgc agtttcctaa atagaggaaa gccactttca	1980
ttcaaagggc tgtggaactc tggctagagg tgggtttcct tgcagttaat catctgcaag	2040
gctcttttga tgctgatc cagaaaccca gaactcacac ttagggtcac aaaatccagg	2100
gcatttattt gccgagcccc atggatgtta tccctatgga tgcaccccgcc ccctgtccgt	2160
tctcctttgg agcagaacga aaccatttcc agagcttttg caggaagtct tcaggccctt	2220
gcgtccggcc ccttttagaca tcaaagcccc ccctgagagc aaaggacttt gaaagatagg	2280
aaaagctcag gatccttate gcgtctctgc tccctcccga cctagtcgta aattccgagc	2340
ctc	2343

<210> 74
 <211> 507
 <212> DNA
 <213> Homo sapiens 4.F.15

<400> 74	
tacgactcac tataggcgca attggagctc cacgcggtgg cggccgcggg cagtgcggac	60
caggcggggg ccctgtggct gccggccaca tcccggagca acagcagaaa caacggcagc	120
agcagcagca gcagctgggg cccgggtccc gggctgggtcc gagcggggac atgagccatg	180
gcgtggtgag ggcggcaaag ggtcgaagtc caggaggagg aaggcgagcg ctggcgacc	240
ggaggctgcg gactgacctc gcggcagtag ggcgcgcggg gagagcccg gcagcagggc	300
gctggatacc gaggtccgcg cggggcgagg ggcttagcgg agcaggcacc cgggcgcgcg	360
gtccgtgggt accggtggcc cgagcccccg gccagcggtc acagccgtcc ggagcagcgc	420
agagccgagc cgagcccag tcggcgcgct gccttggcgg actcgcgctg cgaaagtgtg	480
tagccactg cgcgcccgcc ccggctg	507

<210> 75
 <211> 446

<212> DNA
<213> Homo sapiens 4.F.17

<400> 75
gcggccgcac acacgagggc ccgtcgcgcc ccccgccctg ccccgccctg ccctccacgt 60
ccctgcaccc ccgagtcgca ctaagaaccc agtccccgat cggtttcctc tacgccgtct 120
gagcagaaga gagtgggaac cggggtgacg gataaggggg gggcgccac gcgacgtcgg 180
ggtgcatggg agcgcgcggg aggcgctagt gggcgacgg ggcgtgaggg ggacacagcg 240
cgggcgtggg gatggccact gcgcggggag ggttctgcct ggagaaggag ggatgggagg 300
aggttggggg agcagggcgc gtggaggagg gaggttgac gtgtgtacag cgcctgggga 360
cctcgctggc cccttggtgc cccaggact ctgaggcttc tcctttcggc ttgaaatgtt 420
tttcccttcc tgcttttcaa atctgt 446

<210> 76
<211> 424
<212> DNA
<213> Homo sapiens 4.F.22

<400> 76
gcggccgcct tgaaggcgct ggacgggatg gtgctgaagt cgggtgaagga gccccggcag 60
gtgagctcgc ggcccgccag cccgctgccc acgcagtagt ggaagaggcc gaagtagcca 120
ggcttggggg tgctcacgct gtcgcccacc cagtagggct ggatgaagac caccacgttg 180
atgatggcga agcagatggt gaagatggcc cacagcacgc cgatggcccg cgagtccgc 240
atgtagtget cgtggtagag cttggaggcc tcctgcgagg gcagcatggt gcccggaggc 300
ggggccggcg gcggcgggcg ctggcggggg ccgccggccc gggacggagc gccgggctgc 360
cgggcgggag ctggggacgc acgcgagaag cggccctgag tcaaggaacc cgcgagggcg 420
gggc 424

<210> 77
<211> 558
<212> DNA
<213> Homo sapiens 4.F.6

<220>
<221> n
<222> (1)..(558)
<223> a or g or c or t

<400> 77
gcggccgcag ctcaccactg gcctagagat gccctttgcg aggcggcagc aactgacaag 60
atggtcgcgg gtcgcccgt ccggagccgc ccaccagggt gccaggagga ggcgggagcg 120
gggatcaagc ttatcgatac cgtcgacctc gagggggggc ccggtaccag cttttgttcc 180

ctttagtgag ggtaatttc gagcttggcg taatcatggt catagctgtt tcctgtgtga	240
aattgttatc cgctcacaat tccacacaac atacgagccg gaagcataaa gtgtaaagcc	300
tgggggtgcct aatgagtgag ctaactcaca ttaattgcgt tgcgctcact gcccgctttc	360
cagtcgggaa acctgtcgtg ccagctgcat taatgaatcg gccaacgcgc ggngagaggg	420
ggtttgcgta ttgggcgctc ttccgcttcc tcgctcactg actcgctgcg ctcggtcggt	480
cggtgcggc gagcggatc agctcactca aaggcggtaa tacggttatc cacagaatca	540
ggggataacg caggaaag	558

<210> 78
 <211> 865
 <212> DNA
 <213> Homo sapiens 4.F.69

<400> 78	
gcggccgcag cgagttttct ggcagcgcta gcgccgcggg gcctgggttc ccgggttcgg	60
gtctccgccg gctccgggct cgccccgcg agttggccgc accgttcccc cgccccggg	120
gcagccgctc ctccgggagg ctccggcagg gaccttcgcc ccggcccccg agcggcagtg	180
cggtccagc tggaggcctg gcccggaag caaagtgaag ggacagaggg ctcttctctc	240
gccagccgcc cgccgcgctt ttcccagctc aggcggcgcg cccgcggcgc ggagggagcg	300
aaagagtcgg ggctgcccc ctccaccgcc cgcattctcg ccgccgcacc cgggtccgcc	360
ccgggaggcc ccgcgggagg gaacccccgg cccgctgggc gcttccgcac tgacgccttg	420
gggcgcgcgc cccccgcccc ttactaccgc tacaccgct gggccccga ccccgctccc	480
gggctgctgc cagcgccgct tttccccgta gaaacttcgg agacaccggg gaagctgctc	540
tttgagttg gggaaactta ggaagaatgg gaaaagccga ggaagtcggg gaggaccccg	600
cagttgcctt gccctcggcc gaaattctct tgcaattgga cgggaagcct gccacgcccc	660
gagagccacc cgggtggcacc ccgttgggga cctgcggctg ccctaggctt gagctggcga	720
ccaacggcgc ataccgccgg caccctagg ggaccgtgcc cggccccggt tgggggctcc	780
taacgccagg cttgtgagct atagggtgga gagtgggccc gctcttaagg ggaaaaattt	840
gcggcctttt accaggcaca gccag	865

<210> 79
 <211> 983
 <212> DNA
 <213> Homo sapiens 5.D.9

<400> 79	
gcggccgcag ccagcgccgc ccctcccggc cgggcgggccc caaaagccc tttctgtcac	60

cgcaccaggg cgcgaccggg tgatgcattt ccacaccagc ccgcccacac ctccatgggt 120
 ttggagctcc cgggcaggcg gtggaaactt ggcgaccgt gccactctc cggcgccgct 180
 ccgacagccc gacgggtccc gcggccagga agccactcgg cgcctctcgc cgtcactcga 240
 cccccggccc ctttcggact ccgatcctcc cgtccccagg ccacacggcg cggaaagggg 300
 atgccgagcg ggacgcgcac gaccagggcg ccaggaacga gggcgctgga ggagactccg 360
 ggcagggacc ggggtcccag gggccggg cggggctcaa caccacccg atgggggtgcg 420
 ggcccagcgg ggcccggggg tgggagtagg ggcggcgggg gcccgcgag gaggagtggg 480
 gataggccgc gcagggggtg cccgggaccc cggcgcaag ctgggaaaga ggcacgcggg 540
 ggcggcgcg cggggccggg acaggcgccc gtcctcacct gccgggcagg tgtcccgcg 600
 gcgagtcgc cgcggttgc tccgaggtg aactgtcgtg gtccacggcg catggcgcg 660
 tgaaggcagc ggccagcagc ttcataaggt cggcgccggg gcaggtgccg gggccgggtc 720
 ggaggccacg cgggggccct gggctggggg cggggcgact agcgggctgc gagcgggttc 780
 cacgcgcgc gttcaacggg ctgcaccgc gccgcaccgt gccaacactt cgggcgggccc 840
 ccgctgaggc tccggttgcc cgcactagga ggcgagggcc cccgcgtgca agccgcccgc 900
 ggcgggcccc ggttgccacc ggccccagcc atgggtggg tccgggttgc tttccccccc 960
 tgccccctag ggaattgagc cga 983

<210> 80
 <211> 432
 <212> DNA
 <213> Homo sapiens 5.E.2

<400> 80
 gcggccgctg gtgacctccg cccgcggtca ctgcagccc agccttggcg cgtttgcgca 60
 actgcttttg tcccagacct tcattctggg cgcagtcccc tctcccagtc cccctgccgc 120
 ggcgcctgga actctcctgg tggctgtaag attttcctac cgttaggtcg tctgtggcga 180
 ccgccaggcc tgccccacat cgctagccgc cctgtctacc cctcagctc ccagccacta 240
 aactcgtgg acaaccttac gctagtaaca gtttttgagt ctgagactca tctgtgaaag 300
 ggcagtcata tttgaggact ccaaattggg tgcaagtgcg aaaccacat gcgatatttg 360
 gttgctattg cccacctcag cctgtggcca atgtgtctct gtaggaacag cactagattc 420
 tttggggttt tt 432

<210> 81
 <211> 746
 <212> DNA
 <213> Homo sapiens 5.E.25

<220>
 <221> n
 <222> (1)..(746)
 <223> a or g or c or t

<400> 81
 gcggccgcgg gggcgctcagg tccttgcgcc tcctcctcgg gctcttcccc cagcctctgc 60
 ggggcgtcct ctcccacctc cggggccac tcctcccccg gagagccccg gggcgcatcc 120
 tcaaaagcat cctcctcacc ctctcatcc gtgtccccag cccctcgac gggggctccg 180
 gccgcttctt cccccggccc ggcctcgga aatgggaaag ccgtggagga gggcgagtct 240
 ttggccgcgg gttgcgctgc cgggagactg ggcgcctcgg agaccgggag gccgccgggg 300
 gacggcggtt gctggggctc ccggggctcg gcggccaggc tctcgggcag gtcggagagc 360
 gcggacagcg cctgctcggg gtccggactg cccggggcct cccagcccc gccgctcggc 420
 cccagcagga accggtccag gccaggaag gccccgggct gaggggagac ggcagtgggg 480
 ggcgctgcag gctcctcggc gccctggagc tgctgctgct gctgctgttg ctggagctgg 540
 agctggagct gctgctgctg ctgctgctgc aggcggatcg cctgctggat gtctgaaagc 600
 aaatcctctt gctccgtagc cgaatggaag ctatagatgt ccgtgtccga gcccgagctg 660
 gtcctttgtc catcctgcgc cctgctgca gttncacat cctcggcgat cggccggccc 720
 ccgaccctag cctcggcagg cccagg 746

<210> 82
 <211> 617
 <212> DNA
 <213> Homo sapiens 5.E.4

<400> 82
 gcggccgcgg gccggtgttt caggcagctc ttgggcgcgg gcgggctcgg ggcgggcgcc 60
 gtggaggggct cgggtccaat tctctcgggc tcgggtcccc ctctctctc gggctccgtc 120
 tccgcttctc tctcgggctc aggcgcgggc cctggggggc ccttctctc atccgggagc 180
 acgggcggcg tcgggtccgc ttccttcggg aactgctgt ctggcccgtc gcgagcagag 240
 ggcgcctctg aggtggcggc ggggtcagtc tcggggggag tcgtgtcccc ctgagggatg 300
 gcggtgggaa acgggctcgc gacgtcttcg ggagcacaga ccacctctc cgccttgtcc 360
 gtggccgggg cacacggggc tgcggggggc gcctccccat cctgctttcc gccgtcggga 420
 ccgggattcg gggggccctc cggcggggac gggggctcca cgcggagagt gggggccgac 480
 tcgggctcgg cgagctccgg ggtggccggg cggcttgagg ggtcctcccc ggggacgcc 540
 cctcctcca cgtggtccgt gaggcgggag gagtgtgca ggcgggcgcg tctggcacgg 600
 gcccctccgg gtggcgg 617

<210> 83
 <211> 1840
 <212> DNA
 <213> Homo sapiens A.2.F.45

<400> 83
 ggcgcgccga ggcgcgaggcg cggagaggcg cggcgctctt ggggagacgc ggcgcagggc 60
 atagacgtac gccggcgccct ccccgaggcg gaggggtcgc tgggcgggcg ggagtgaggc 120
 gcggcgcccg cgagagacg cagtcgctg ggctgagggt ggcggggagt gttgcagtcg 180
 tacattcgcg cgccgccggg cggggagcgc gggggtggcg cgggtgcaggc gcagagacac 240
 acgtacccgg cggcgagag acgagtggaa cctgagtaat ctgaaaagcc cgtttcgggc 300
 gcccgctgct tgcagccggg cactacagga ccagcttgcc cacgggtgctc tgccattgcg 360
 cccctactg gcgactagga caactacagg gccctcttgc ttacagtgtc gtccagcgcc 420
 ccctgctggc gccggggcac ggcagggtc tcttgctcgc agtatagtgg tggcatgccg 480
 cctgctggca gctaggaaca ttgcagggcc ctcttcctca cattgtagtg gcagcacacc 540
 cgctgctgg cagctgggca cactgccggg ccctcttgct cgcattgtcg tggctgcacg 600
 ccacatgcag gcacatgggg actacgcagg gccctcttgc tcccgggtgtg acggctggcg 660
 tcccatattg gccacctcct gcaccactta aagtcagagc gccagttatt aatccccatc 720
 agttctgtaa attaaaactg aaaaggagct attactgcgg agagctgatg tcccagttat 780
 taacttgga gacagctttt caccaagagg cagtacaaag atggaagata acttcattga 840
 aaagaaatac agtgtaaaga gcttattgta caaaaatagg gaggagtagg ctgatactgc 900
 atgaaaacag cctaagagtc ctgtgcaggg atttttattt tggacttctt cacattccta 960
 cctctgtctc aagtctccgc ctgttttctt tggttttcct gctactgcct taggtccccg 1020
 acttgcccca cttagccttg tgggacctcc tcaattgatt gaggtacatg tgtggtgatc 1080
 aatccgaatc cactctggca ccagcctcct tcccaccata ccaggcaggc tgacagcggc 1140
 cacgtttgta tctactgcag ctgcctcttt tgaatgtctt tctctgcctt aatctgtact 1200
 tatggtgcca ggtttctctt aagaatgtcc cctttgtcct tcttatcagc atgtagctag 1260
 caatattctg acatttttat tgcagaatga atgatgattg gggcttcttt tttttttttt 1320
 tttttgagac ggagtctcac tctgtcacc caggcagact gcggactgca gtggcgcaat 1380
 ctgggtcac tgcaagctcc gcttcccggg ttcacgccat tctcctgcct cagcctcccg 1440
 agtagctggg actacaggcg cccgccaccg cgccagctaa ttttttgtat ttttagtaga 1500
 gacgggggtt caccttggtta gccaggatgg tctcgatctc ctgacctcat gatccaccg 1560
 cctcggcctc ccacagtgtc gggattacag gcgtgagcca ccgcgcccat ccgattgggg 1620

catcttaaga gaagttctag ggtgtttctg cgtaggtacc ttttctccct cctaaccaca 1680
attgacaagt gcccatccac tccagcacta gagatgctac taatatgtgc atttttggtg 1740
gtccctccag gtgagccttc acagactttc ccttttccag gagctcccc cctgttcat 1800
gtctagctag ctatctactc taacagagcc cactatcctg 1840

<210> 84
<211> 3592
<212> DNA
<213> Homo sapiens A.2.F.50

<400> 84
gccgaggagg cggctccgac ccaggtcgtc gcagcagcac aggaagctgt aacacaggta 60
agtgcaggag agcgagagcg tgaaggcgaa gagcagcctg cgcgccctcc gcggctgagg 120
tggccccgcg cggcccagga ccctataggc catggctcca tgggccccg cggggggtca 180
tggtttccga gggggcaccg gcggctgagc tgctgtggcc ctgcggtcgc ctagagggtc 240
cgcgtggcgc tgccacggcc acgcgggtcg ggcgttgggg gcgccgtctt ctccgggggc 300
tgctgaccag ggtgcgcaca gtgccagggg gtcccggggg cagcggctcc tcggggaaca 360
ggcggttgca tttccagcat ctcccggctc taggcgatgg ggctccgggc agccgggcgg 420
ctcgggcgct cccaggctct tacgtgcgcc gggttcggag cgcgccagc gccgaagcc 480
ccattcctga tcctcggagc gccgctcacg aaacgctcgg cggcggcgcg gctgtgcggg 540
ctggcggggtg gaccggacgg tggcgctggc gccggccgcg atctggctct tcgggaaatg 600
ccgagcggag cgcgctgccg gctctattta aggagtggcc tgacgtcagc cgcgcgggtc 660
ccccgagccc gcgccgcgcc cagggacctg gccgcggccc tgcgccccca ctctcttacc 720
cctcccagaa acacagcacg cgggcccctc ccatgcaggc cactccctac ggagccccag 780
gccagctttg gggcggtgaa acgaaggtgt caaggcatag tactcctccg ggaggctgga 840
cacccccacc acgtggcct ctgcacatcc agggacacga atccaggctc agatcgcgcc 900
gacatgcaga ccagacagac ccagacgcag acgcaggcac cctgccctga tgcgcggctc 960
caccaccctg acccgcacac gcacgcacag gcacagaagc acacgcgccc tagcccggac 1020
acacccccac acccacgcgg ggggtggggag gagaagtccc ctaacctggg ccagataca 1080
ccgacaagga cactcccccc gctctcgaca tctcgccaaa tggacacaca cagcccggaa 1140
tcggacaccg agcgcacgca cgcctggac tgggacacgc gctgtagacg ggatgggtgg 1200
aggagccgag cgtgagttag attccgtgac tattcacca gcttcttagc cccagcgcg 1260
ctgactcaca ccccgggcgc tcgctctgtc tcgcacctat gaggcacgcg cgcaccccaa 1320
cccattgtca cccacctct cccggggcct gccggagagc gagccccgga gcggcagact 1380

ccgcgtcagg	agggttcctc	tcttagcagc	cgccgcctag	cggtagactg	ctccccgggg	1440
agctgtccag	ggtaccagag	ggtcgccgag	ggctgagtga	ggagggcttc	ttcacacaga	1500
gacactagga	ggaggaaaca	gagtacaagg	agaacgtatc	caggagcaat	tccacttcga	1560
atgattccta	agtgaatgcc	tacaggacag	ttctcggtga	ccatgtccag	aacaggcata	1620
agtgacgatc	cccagtactt	ccctgagggg	ccacactggt	accttggtatc	agaaccctgc	1680
atcagaacag	gcctaaatgg	ccatggctaa	gaacacggct	gagttgtcct	tcaacagcaa	1740
tgccaatgcc	aattcaccat	gtccgagtgt	tcacaagggtg	agtgcctcc	accaccaccc	1800
agccatagaa	tgtctagatg	accaccatga	ccccaccct	gatcagggtg	taactgactt	1860
ccttcctcag	gctgtaaact	gatcattagg	ttctgtggat	cttagcccaa	accagaaaat	1920
attttgtccc	caaactagtc	ccatccctag	aaaccttaaa	ccaattctac	ggcagataat	1980
aataatagct	gccaactttg	tatcaagcac	ctggcatggg	ttaactgatt	aaatattcac	2040
aacctatgaa	gttggtacca	ttaccctggc	atcactttgc	tgtcttaatt	ctaatagtag	2100
ctagcattta	ttgagtgcct	gttttatggg	agttatgcgc	taatcacttg	acatgcacta	2160
cctcatttat	ctttggagat	aggtattatt	gtaatttcta	atctacaggc	agtgataaga	2220
agatttaaca	aacatataca	cagtaactgg	cagagctggg	attaaaccog	ggcagtcttg	2280
actccaagat	tcaagctctt	agttacagca	ctttgcagct	tcctaacttc	ctttgaccat	2340
tattcatata	attccatcct	aggctcctct	cctggatgta	agctaatttg	tctatgtctc	2400
ttctaaaatc	tcacacctgg	gactgcgcga	ggaatttcag	atatggattg	aaaagttcaa	2460
caggactctc	acctctcttt	tgtaagttct	atctctagta	atgccaccta	agactccatt	2520
atctttttct	tgtggctata	tcacactgct	gacatctcaa	acttgcagcc	aagtaacatc	2580
tctaaatggt	tcttacaagt	gctgctgatt	aaggcacagc	taccccatac	tgtgcttgta	2640
cagtgggcct	ttttggaccc	aatgtgtagg	tccttataga	tttgacttga	ttgcatttca	2700
tcttgtctca	tcagttcgct	gccctagttt	tttttaaagt	tctatttgaa	gtcaaaccac	2760
gaggtagctt	tcatttattc	aaaaagaaaa	agtagaaaga	ttgtatccca	gctttaccct	2820
ttattccagg	tgtactttgg	gcaagtggac	cccctttaag	cctcagggtc	ctcagctgta	2880
aaatggggacg	ctatgattca	ccttaaaagt	ctctcaaagt	ttagatgttg	catgattcta	2940
tgattccatt	acccaaagca	tgaaccactc	acttggcatc	atgtaatttc	cacagttgat	3000
cacaatttaa	ttaattcctc	attctaattg	ttaataaaaa	tgtcaaaaca	aatatactta	3060
aaggagtctt	tcttcttctt	ttgggtgagg	ggaagtgtct	cactctgttg	accatgctgg	3120
catgcagtag	tgcaatcata	gctcatgctg	cagcctccac	ttcctgggct	caagggatcc	3180

tcctgcctca	gcctcctgag	tagctaggac	tacaggcatg	tgccaccaca	cctagctagt	3240
tttttaattt	ttttagaga	tgaagtctta	ctgtgttgcc	caagctggtc	ttgaactcct	3300
gagctcaagt	gacctcctg	cctcagcttc	ccaaagtgt	agaattacag	acatgagcca	3360
caatgcctgg	cctggaagga	gctcttatat	atactttgaa	caattattca	catcatgaac	3420
ctgctatttt	tgtattccat	tgttaaaatt	acaagggttaa	atgtggagtc	atctgctgtg	3480
atcagtacta	tttcccttag	aaaataaaac	atgaatataa	tgattttctca	taattctgtg	3540
cttggtctaa	tttttaaata	atttttaacc	tttgaattca	taaactgtga	ta	3592

<210> 85
 <211> 2722
 <212> DNA
 <213> Homo sapiens A.2.F.67

<400> 85	
cgccgcgcgag	gacactcggg
cgcacacccg	ccgcgctggc
gtccccacc	cccagcccaa
60	
acaaaagaca	agccttgggg
tcgtggcctc	gctgggccc
ggcgccccga	gccggccagg
120	
gcgcctctg	gggcccagagc
tccatggttt	gcctaaggca
tagcttcttg	gcggtaggcc
180	
gcaagcggcg	gggagacgcc
aggcagggt	gggcccgcga
gaggtccgaa	gatgcctcca
240	
gtcgccgccc	cggggaaggc
gcgggcgacc	tctgagtgtc
ccggtaacgt	gtgcctttgt
300	
tccccaaactc	aggtgaaaat
ctggtttcag	aacaaaagat
ccaagatcaa	gaagatcatg
360	
aaaaacgggg	agatgcccc
ggagcacagt	cccagctcca
gcgacccaat	ggcgtgtaac
420	
tcgcgcgagt	ctccagcggg
gtgggagccc	cagggtcgt
cccgtcgt	cagccaccac
480	
cctcatgccc	accctccgac
ctccaaccag	tccccagcgt
ccagctacct	ggagaactct
540	
gcacctcgtg	acacaagtgc
agccagctca	atcaattccc
acctgccgcc	gccgggctcc
600	
ttacagcacc	cgctggcgct
ggcctccggg	acactctatt
agatgggctg	ctctctctta
660	
ctctcttttt	tgggactact
gtgttttgct	gttctagaaa
atcataaaga	aaggaattca
720	
tatggggaag	ttcgaaaaac
tgaaaaagat	tcatgtgtaa
agcttttttt	tgcattgtaag
780	
ttattgcatt	tcaaaagacc
cccccttttt	ttacagagga
ctttttttgc	gcaactgtgg
840	
acactttcaa	tgggtgccttg
aaatctatga	cctcaacttt
tcaaaagact	tttttcaatg
900	
ttatttttagc	catgtaaata
agtgtagata	gaggaattaa
actgtatatt	ctggataaat
960	
aaaattattt	cgaccatgaa
aagcgggaatg	tttctgaaaa
atacttcatt	ctgccccctc
1020	
gataactggc	tagtgaagtt
ttattgaagg	caactaaaga
aggacaagct	ctgcagagat
1080	
ccaacaaggc	aaaaaagaaa
acagaagtcg	gggctctatg
catgcagact	gtatatgtat
1140	
atatgttcaa	tgctataact
tgtgtgtgtg	tgtgcatata
tatatataat	atatatggca
1200	

tgtttatagt actgccatat ctcataattg tttcaggtag aaagtaatgc tgaaataaaa 1260
 atacatccct ctcaccctgt atgtgagtta gaaggcaaca gaaatccctc aataaccctt 1320
 ctgaattcta agctcaaagc aattatcttg gagaagcgcc cccacccatc agcctctgtg 1380
 tagtgccaga gcaattagac aaattaccct tcaaaggagag tttccagaga tgagaaaatg 1440
 aaaaagaaat ctaccctcac acctattaca ttttttaaaa atctaaaatg tttggagcat 1500
 ggcaaagtat agaaccttggt actcttttga gtatgattat aaatgtatcg gctcttttcg 1560
 agagatgaaa acattgcaga tattgtgaag agggaaacttc agggttgggg aaaggaagga 1620
 atgaaagcat tgtggcgccg tgttgatttc attttgtgtg agataatact cttaatat 1680
 ccttcccg cttccttttt tcaggaagga gcttcctctg ttttgctttt acataaaaaca 1740
 gtggcaaaca ggttctaaat gatgcaaaat agaactctgtt tactaggatt tctcctttgg 1800
 gaagccttct ttgggacaga gaggaaggac ttgctgcagc tgtgccctgt gtcccttctt 1860
 tcttcttgca ctctgcagc tagataccaa cagcatgacc agagctatgc actgcaccta 1920
 aagaccagg cctgaattgt aggtgtcttt ctgtctggcc gtccttcagt gggccagact 1980
 ctcttctctt aggatacgaa ggaaaatgtt ggggttgaaa ttacaagatg catgtgaaat 2040
 attttacagc taggaagtca gcagcaataa atgtgacaaa agagccttct taaagtgggg 2100
 gtagattaga gcataaaaaa ttatatcctg tctactgagga tttctcagaa ggctcttcca 2160
 ggggttgagg actagacctg aaaaggcacg ctatgtgcct tgaggggaat ttaccttacc 2220
 tacatgtttc tctctctgtc tctctctctc ctctctctc tctctcttct tctctcattt 2280
 tctctgtctc tctgctgcc tctcctctc cttctctccc tacctccctt ccacctcctt 2340
 tatttttttc gttctcttct cctttacttt ttttctagaa gagttaccag gcccgccagt 2400
 gtggaacagc ttgcttcttg gaggaatcag tattttgacc gctctttaga catatcccg 2460
 agcctggctc cgaggcagaa ctacgcccg cagcctggcc tgtgcacccc tctccggca 2520
 ccccgagcg cgcgactca atatttccgt ctccccagtc cgctccagcc gtactttctc 2580
 ggaaggagca ctgggtgcgg ggaagagggg gcaataggaa ggtttgcggg gggcgggggg 2640
 gggggcgggg agccaaaggg tgccccattt tgttttctgc gctcacagag aataggggga 2700
 ttggggaaga gatgaagata tc 2722

<210> 86
 <211> 3366
 <212> DNA
 <213> Homo sapiens A.3.F.38

<400> 86
 ggcgcgcctc cagttccaag gccgagctca ctttcaacag ctctggaaat atgaatgtat 60

ttttccccc	tttagaagaa	gctatacgag	gaacaacttt	ttgaaatcgg	gagtgtgttt	120
gtagagaagg	agataaggat	tgcatttcgc	ttatTTTTct	acaggtgata	gaagtgtttt	180
gggggtcaga	gtatcctctc	aaggaaaatg	taaaacgtgg	gggctcgcat	tctctatcta	240
agcctttgta	agttaaatta	acaggaccct	taaagtattc	cttatagcta	cagataaaaa	300
attacaggca	atgtttggat	aagggggcaa	ctctccgtgt	ccaaacattt	agagaactgc	360
ctgtgagtgt	acaccgttgt	aatcttattg	ggagcccttt	gtcgaattct	gtatTTTact	420
ttgatgcttt	ttgagtacca	ttcccattgt	ttgggtgtcc	tttaactccg	tttacagcaa	480
tatattaata	aagaggatgc	atatgtcagc	gttatgtatc	cacaagaatt	tggattcctt	540
taaaatcaaa	cggcttggtg	agcaggcaag	cactcaaaac	ccaacagtct	caaacagcaa	600
taataatgtc	agcaaacggc	tgccatgcct	cctTTTTctc	aaatgctggt	tattctaaaa	660
tcaataagtt	aggagataca	ttgcagagaa	acagtcatta	gtggttcagg	gttggcaggt	720
ttgtTTTTca	ggtgtagatg	ttcttgagta	atacctctcc	actgtggact	aaatattagt	780
agattgtcgt	tgtcattttt	ctaatttaat	gcggcagcct	cagggaagta	ctcatccaga	840
caattatggg	gtatcgattt	ttaactttaa	gattaaaaaa	ataccatatt	tcacttgcct	900
tgggactact	tttcttgata	aaaatatatc	tgggaagatg	attttagggc	catgttagcg	960
taggggaggg	gaattaaggc	acaaatgggtg	gttgggttaag	gaaattttat	gaaagaaaat	1020
aaagaaaaca	tgtcagaata	aatcaatcag	aggcacaagt	gagttagagg	aatctgagga	1080
caaccagcat	cttggggatt	cttctgttcc	cgcggttctc	agatatagga	ataagggctc	1140
gagttatgcc	cagaatacat	tcgtctggta	ctggatgtcc	cagtccctta	gctgttccac	1200
gtaatgaaga	agctctaatt	cccgagaact	ttggggctta	tttttaccat	cattgagtct	1260
gcccaggctc	agctctctta	caaagggtata	aatctgaaat	tcatgtatta	atttgaatcc	1320
ccaagatccg	agttatgaga	aagggcaagg	gcaggctcta	ctcctatttt	gtttactttc	1380
accgagttac	tgtgaagtga	ttggaaactt	tcttaacggg	cagagagaga	atacacggaa	1440
actcggatgc	agtaataaag	ttgacatagg	agtcggaaca	gggggctctt	tttgatctc	1500
acctttactg	gggcttgagg	ttgtggaatg	ggtggaagag	taattaactg	aatgaagaat	1560
tttaacgttg	aaaacagagc	ccacagtatt	tttggttata	gtggtgtggt	ctctgcctcg	1620
gcaaagaaac	aaacaccccc	acccatctt	cgcagttctc	ctctctgctg	tagcgacgcc	1680
aggcgctgct	ttccgccggg	taaattagcg	gcgagcctcg	ccagacgctt	tcctccttgc	1740
cttctttcgc	cgaaaggggg	cgcgctcctc	ccaggctgcg	ctggtaccta	tcctgccttc	1800
aaaaattttct	gggttcctgc	aggacagaca	gtaacaaaac	gtgggaaata	atagtttgat	1860
gacacttcag	ggactatagg	aatataaggt	gcacacacat	gcactttaat	ggaacatgt	1920

agacacctgg caggagcatt ggctgcctgc ctctcctcct ttcaaagtag ggtgggtcggg 1980
 gttccagggt ggcaggaggg gagtggggcc agatgaccgt ggatggaatt ggtgggtgct 2040
 aggactgacg cctgggttcc atggcggagg agagggtttg tccccatgga gctgtgtgga 2100
 cttttctgca tatgtacttg aggtcttcaa agaaagaagg gcagatctga gaaatggaga 2160
 agtggctggt attagtgaga tgttgaaaaa ctgccacaga agccctcaca gtgcctggag 2220
 tgtaagaca gaagagaaaa cctggcacca tagagtttta ggccctggga tcagggtaac 2280
 ctttcctcct cacgaaagaa caataactgc cccaaatctt gtgtgagcct gcaacttggg 2340
 tacctaaagc catttccaat ctgcaaactt gactcctggc ctccactgat cctccatttt 2400
 tgggcaagag tttcaagaga ctcacaggac agatgaggat aaatttttaa ccccttctgt 2460
 aaatttaggg attttcgact tcttaccact ccctgacaat ggggggtcaac aaatcaaggc 2520
 acggtgagag taacaaactg gaataatata tattttgtct tcatagcata gatgatggtt 2580
 aatacactt ttccaagata atctgagctg gagtggtcac tagaaacagg agcacaaggc 2640
 cagaactgta aggcaaattg ctttcccaca aacgtttgtc tgagaataag aacattcacc 2700
 ccattcactt aattttctcat catcagtcac gtcattatat tttcaaggac ctcacagtgc 2760
 tggaaagtgg tgtagttata aataagcata aaaacagatg ggtgatccca gtcctctaaa 2820
 tataatcggg gatgccaaat cttttcaaag agaattcata tatacaactt aaaggccaag 2880
 gagcccaatt caatcaaaat ttgagccagg atatgctaag ttcaatcagc ttgaatatgg 2940
 gcaaagtgta agacctagcc agcacttcag atatatacag agaaccacat tttctcaagt 3000
 ttccattggt attttccaca caaatttagt gttagtcttc aaagggtattg ttagatttgg 3060
 tttgggccgg gaggggtggtg agagtcagtg cccagggctc ctgtccttgt ctactccctt 3120
 ttcttttggt ctctctctgc ttcagcagtt tgccgaaaat ctgtgttgca gagaaaattg 3180
 acacctagag gccacagagg tctcctaaat gctgttttct aggatcctca gaaaacaaga 3240
 ggaccgctga gctcaattat atgtaatata cctggatatc ttatgtattt ttcttttctg 3300
 ctaattcatt ttataatagc taagttagag acttcttgga gatttaggtt ttgggggactg 3360
 gatatc 3366

<210> 87
 <211> 638
 <212> DNA
 <213> Homo sapiens A.4.D.30

<400> 87
 ggcgcgctc gcccgagatg cccctgcgtc cgctggcca ggcctggggg ttacccgacc 60
 cgggttctcc ctctgctggc tttgcgccc ttcacacctc tgcggtgggg acggagctgc 120

cgagacaagc agagtgcgaa ctggagaaaag cccagagctc agagctccca ggagcccacc 180
 gtgccccacg gctaggcggt ctctgggtgt ggacggctag cgggtgcatt acttcttaca 240
 aaagtttatt tttgaaagct tctcccttcc ttccttcttc ccttccttcc ttttcttctt 300
 tttttctttg ttttgagtca ggttctcact ctgtcgccca ggcaggagcg cagtggcgct 360
 atctcagctc acggagcctc cacctattgg gctcaagcga tctctccacc tcagcctccc 420
 gagtagctgg gaccacagtc gcacgccacc acgtccggct aattatTTTT ttcgtTTTT 480
 cgtagagagg gagtgtcggt atgtgcccc ggctgggttc aaactcctgg cctcaagcga 540
 tctctccacc tccggcttcc caaagtgtcg ggattcgggg tgtagccac tgtcccgga 600
 tacttctttt ttatcctgtc agaaaaacta tccatggt 638

<210> 88
 <211> 1860
 <212> DNA
 <213> Homo sapiens A.4.D.36

<220>
 <221> n
 <222> (1)..(1860)
 <223> a or g or c or t

<400> 88
 ggcgcgcctg tccccaccta atgccacgat cccccctcc cccaccctnc cgcactgcct 60
 cccttgcgcg tgtaggggag atccctgacc ttgtctgccc agctgcaggc cacttgccca 120
 ggcggccctt cccttggtgc cacctccgc ccagctcacc aggagcgtgt gccctgttgc 180
 tactggcaac tgctgtgcc taaagctcag cccccaaact ggcttaatgc tgattgatgg 240
 tcagaaatag gatattttct ggaacagagc ggagcgtgg tgcaaggccc tctctgtgc 300
 tgagtctag ggacctccg ggtggcaggc ctctctctc ctctcctttt ggccccaccc 360
 acctacact accctcaga gaccaacggg ctcttcggac atctcatct cagggttaagt 420
 gctgagccag caagccagtg ttcgttttct tgctgagtaa caggcagcca ccccggaatt 480
 tctctcttta tcttgaggc ttctgagttt tatgaatgag gcccggttg ctggacgcta 540
 ccacttccct ttttattttc atccccacta acttggtcac tcgttcactc ctcttatac 600
 ataggtaact aaaatagact accctctag taaccagaac tattcctgca aacgcttaca 660
 agagcatttt ccagaaataa atcatttcat atcagtatcc ctctctcagt catttcccg 720
 cttcatgcca cctccctcct aagacacaga attggtcatt tccaccactt taaagacaca 780
 gtctagataa aaagcctgca tttataatgt tctttgcagg agtagctttt gcctattttg 840
 tgggggtttt gtttggtttt tgtttctgt ttgatactcc ctctcaaact gcagcctccc 900

ttcccttttc tgggatggca gcctccttct ctgagccatc ctggactaac attttctgga	960
ctaataaaatt tctgcacctg tctctactcc ttctccttcc cagtctgact gtaaaggacc	1020
agatttcatt atcaaatcaa ttctcttttag aagaactttg ttctgtagca tttctttcca	1080
ggacccaat atttttggca gagtattttc attatttaaa ttgtcgtact tagcttcttt	1140
ttgcctatgg acattacttt ggaaaacat gtgatgtttc tgagtcactg atttgcttct	1200
ccaaacaaaa cttccttcag aggctcccat atgttgggca ccattgtagg cccccggggg	1260
tgggaatgga gcaaagacaa gacccaaatg ggtttcagca ttttaaagcc ccattacag	1320
ctggtttatg gttattgcta tgatggttaa tgtgataaca gcacactaca tttgactagg	1380
actttacagt ttacaaaagg ctttcaaaga cattatctcc attaatcca gcagcaggaa	1440
ttttaaatag caaggattcc accaaaaggc ccagtaatgc tcaccaatcc tgcttaacca	1500
aaaagaaaaa tattgcaaat catcctaaca gctgatggag ctttaaaaca cagaataaac	1560
aattcataag aagcttctga agcttagtta ctggaatgta acttgagaa gataagtga	1620
atgcacgtaa catgtatatt accagaaggg tgtcttgag agaaactcca tctggggct	1680
tcagtggcct ggtgaactgc tggaggtgga ggctttccag ggctctggac tattgcctta	1740
tcctaggatc taaaatggga tgaaagtgtt agcacaaagt tgctgggaga ctagcaaatt	1800
aagcaaatg agtaggcaat gatgttactt tctttagcta caaagcattc ttgagatac	1860

<210> 89
 <211> 2107
 <212> DNA
 <213> Homo sapiens A.4.E.32

<400> 89	
ggcgcgccac aaggcgtgg tgctgcgctg ccacgctgtg ctgctggcgc gggcgacaa	60
ggcgcgcgcc ctggcccgcc tgctccgcc gaccgctg gcggccttca gcgacttcaa	120
gcgcctgcag cgccagagcg acgcgcgcca cgtgcgccag cagcatctcc gcgctggggg	180
cgccgcccgc tgggtgcccc gcgccccact gcgcccctg ctcaatgcc agtgcgccta	240
ccggccgccc cagagcgagc gcagccgagg ggccgcccgc ctcagcagca tccatgagga	300
ggacgaggag gaggaggagg acgacgcgga ggagcaagag ggaggagtcc ccagcgcg	360
gcggccggag gtgctcagcc tggcccgga gctgaggacg tgcagcctgc gggcgcccc	420
ggcgcccccg ccgcccgcgc agccccgccc ctggaaggcc ggccccagg agcgggcggg	480
ccaggcgcg tgagagccga aggacaggac tcgcagcccc aggcccgacc cgccagactc	540
acagcctcca accccggccc tgcccgttc ggctgcccc ggccccggcc cgtgtctccc	600
ccgtggtctc cgtgttgtcc gcccgcgc ctcattttgg ctcaaggatga tgctgatac	660

gcccttggtt attggggggt gttcctctct cccacaccc ggagtttccc gggcctgcca 720
 ttgtggaccc gccccctatg ctttacacct agtctctttg cccacagacc tcctcattcc 780
 ctcccaaaac atcctctcaa gagaaggag gagaagtttc aagaaatcag gaggggtggg 840
 tttggaccct gggcaggggtg gaggcagtga ccttgccctt ggtccctcta gccttcttcc 900
 ctgtgcaaaa aaaaatgacc ctggagaggc attctttagt gagaagaatc tagcggccgg 960
 ggagaattgg ggccggggccg gcggtgggca gagtccgctg ctatacacac agggaggaat 1020
 tctcacgccc aagccccgcc tctctacgcc ttggaggact cctgtgactt cactgctctg 1080
 cctctggaga aactgaggag agtcctaccg acgttcaaac aacagggttag gccaggtaac 1140
 agccctgcac caggccgctg cccacgcctc tgccctggca ccccagggg attccttgcc 1200
 catcccatct ctctgcagac ggatgtgtgt ggccccctcc taggtgcccc acaaccagga 1260
 ccaagatggg gctcccaaag gaggtaagga gaacctttgg caggtgctta ggacactgac 1320
 tacctagaaa gtagacgcag cagagttgct cccaagtcga ggctcctcag agcagggtggg 1380
 tcctgacagc agtggattct cccagcagga tgaggaagga ggggtgtgta accaaccaag 1440
 ggagtgggcc cccacccag gtgtctccgc aagaccacaa aaagcccaa gatctatgtg 1500
 tcaactgatca ttgtaaataa agtggacctg cttttacagc cctgtcacta ctctgtgtt 1560
 gtgtttaatg ccaggcctgc tgggggtgaa aaaatggatt gaagatcaga taagccacag 1620
 gtgagcctgt atagctcccc ctggttacca tcagaaacct gaaagtagtt cttttgagca 1680
 gccagagcca accccaggat taggacggga tctggggact gctgccagga agctgttctt 1740
 taatgtcaga gaaggaggca gtaacttatg ccttgtctga aaatcacatg tgccaggctc 1800
 cctggagggg cgtcggctgt ctgtctcagc ctcccaggat gtctgtacgc ctgggcactc 1860
 agatgcaggt gtctgggaca tttggcaggg agggagcact gggctggggg cttctcataa 1920
 gcatgtattc atatctctga gaaggttcat gtgtatttca gagcatatgg tatagactgt 1980
 gtgtgtgctc tcagggatga gtgcgagcag gttgtaagag aatgtgggtga gcagcccagt 2040
 tttctttcag aggctctgga aaaacctgtc cagaccctgt ggcagtgtga gtcttcagct 2100
 ggatatc 2107

<210> 90
 <211> 498
 <212> DNA
 <213> Homo sapiens A.5.E.28

<400> 90
 ggcgcgccgg agttcgggct gccggctcct tagccgcggg gcgggggaga cgctcgggga 60
 aggggagagg cgcggggcggg tgggaacggg cgggagacga gcggggacgg ggagacgcgc 120

cggaggcccg gagcccgcgc atgctcagtg cgcggccgga ggaggcgagc gctggggacg	180
cagcacctgc cccgcgcggc cgagaggcgg cagccccagg tcccagcgc gcgaaattag	240
taaagggcgc ctggcccgat tctcaggcaa gaggagatta tcagccgat tcccgtgcgg	300
ggacgtaggg gttgcgttgt tcagcggcca gggatgcgc gaggcgatgt ctctccctt	360
tacaaccgca gtatcggggc acgaggaggc gcgaccttcc tgggtacca aacctctggc	420
ctccgggaga cgcggaattc gggggatcgt taaggcgccc tggccaggga aacagatgct	480
tctgcgtctg ggctgaaa	498